Children's anti-inflammatory reliever study, United Kingdom (CARE-UK)

Submission date	Recruitment status Recruiting	[X] Prospectively registered		
27/01/2024		☐ Protocol		
Registration date 19/11/2024 Last Edited	Overall study status Ongoing Condition category	Statistical analysis plan		
		Results		
		Individual participant data		
13/11/2025	Respiratory	[X] Record updated in last year		

Plain English summary of protocol

Background and study aims

This study aims to see if a new type of asthma reliever inhaler reduces asthma attacks in children. Children who have asthma attacks are treated with reliever medications (blue inhalers), which act quickly to briefly open the airways when they become narrow but their effect soon wears off. These blue inhalers don't treat the underlying problem, airway inflammation (swelling). Asthma attacks can be prevented with an inhaled steroid treatment (usually brown /purple inhalers which prevents the underlying airway swelling), a maintenance treatment. This new type of combination reliever inhaler contains a fast-acting steroid working by helping to dampen inflammation and is usually taken regularly/every day. In adults and teenagers, this inhaler has been shown to work well by relieving symptoms and treating the underlying airway inflammation resulting in fewer asthma attacks. This combination inhaler can be used instead of the usual reliever medication on its own for people with infrequent asthma symptoms or as both a regular preventer and a reliever for those with more troublesome symptoms.

Who can participate? Children aged ≥6 to <12 years

What does the study involve?

Children will be randomly allocated to one of 2 groups:

- 1. Usual care, no change in treatment
- 2. The new approach, using the combination inhaler on its own as a reliever when symptoms occur or for both maintenance and reliever treatment

This study will identify children through GP records, and attendance at hospital asthma clinics or they may self-refer through social media adverts. Once they have agreed to take part in the study, the study team will ask that they attend 4 visits in total, some face-to-face, some over the telephone, and some in person at their local study site. The study will see which group has the most asthma attacks requiring a course of steroid tablets. The results of the trial will change National asthma guidelines and findings may be shared by Asthma and Lung UK on their platforms.

What are the possible benefits and risks of participating? Children in both groups of the study will have access to the study team, who will make sure parents and children know how to take their inhalers, have an up-to-date Personal Asthma Action Plan and know when to seek help. It is hoped that those in the intervention group will find it easier to have just one inhaler to use.

If the child is on the intervention arm, using the combination inhaler, there may be a period where the child will need to adjust to the new inhaler. The schools will have to get used to the revised Asthma Action Plan as the number of puffs of the new reliever inhaler will be different to the number of puffs of the blue reliever inhaler. Detailed directions will be sent to the schools about students who are taking part in the trial and their new Personalised Asthma Action Plan. Parents will also be encouraged to review the new Personalised Asthma Action Plan closely with the children's teachers to ensure everyone in contact with the child is aware of the change of plans.

The new combination inhaler contains steroids which can affect growth. The study will measure height at the onset of the study and the end. Some children will be given an electronic monitoring device for their inhalers so the study team can keep track of how much medicine they have taken. For all children, prescription records will be checked so that inhaled steroid doses and courses of oral steroids can be compared in each group. The risk burden is fairly low for these patients and the long-term benefits of controlled asthma, and hence reduction in asthma attacks needing a course of steroids, outweigh the potential side effects.

Where is the study run from? Imperial College London, UK

When is the study starting and how long is it expected to run for? June 2023 to February 2027

Who is funding the study? National Institute for Health and Care Research (NIHR), UK

Who is the main contact? Louise Fleming, l.fleming2@nhs.net

Contact information

Type(s)

Scientific

Contact name

Dr Keith Boland

Contact details

Room 221. Level 2, Medical School Building, Norfolk Place London United Kingdom W2 1PG +44 (0)20 7594 9480 k.boland@imperial.ac.uk

Type(s)

Principal investigator

Contact name

Dr Louise Fleming

Contact details

Praed Street London United Kingdom W2 1NY +44 (0)20 7352 8121 ext 2938 L.Fleming@rbht.nhs.uk

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

Integrated Research Application System (IRAS)

1009041

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

23SM8661

Central Portfolio Management System (CPMS)

59444

Study information

Scientific Title

A novel reliever strategy for children with asthma: children's anti-inflammatory reliever study, United Kingdom (CARE-UK)

Acronym

CARE-UK

Study objectives

To determine the clinical effectiveness of budesonide-formoterol reliever therapy either alone or as part of maintenance and reliever therapy (MART) compared with SABA reliever therapy either used as monotherapy or with the child's usual maintenance treatment in children with asthma aged ≥6 years - < 12 years old.

- 1. To determine the clinical efficacy of budesonide-formoterol reliever therapy either AIR alone or as part of MART compared to usual care
- 2. To determine the safety of budesonide-formoterol reliever therapy either AIR alone or as part of MART
- 3. To determine the cost-effectiveness of budesonide-formoterol reliever therapy either AIR alone or as part of MART compared to usual care

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 10/09/2024, Wales REC 5 (Castlebridge 4, 15-19 Cowbridge Road East, Cardiff, CF11 9AB, United Kingdom; +44 (0)2922 940910; Wales.REC5@Wales.nhs.uk), ref: 24/WA/0046

Study design

Randomized controlled open-label parallel-group study

Primary study design

Interventional

Study type(s)

Efficacy, Safety

Health condition(s) or problem(s) studied

Paediatric asthma

Interventions

Randomisation

If the child is eligible for the study they will be randomised. A computer-generated randomisation list will be created and allocation concealed using an online system (OpenClinica). Stratified block randomisation will be used. The randomisation will be stratified by:

a. Site

b. Current GINA treatment step (see 'Dosing Regimen' below for more details)

The study is open-label. Blinding is not being performed in order to maintain the potential real-world advantage of ICS-formoterol as a reliever.

Control (Standard Care) Arm

Patients in the control arm will continue with their current standard treatment as prescribed by their usual healthcare provider. This will be one of either:

- SABA: A short-acting bronchodilator that provides immediate relief for asthma symptoms.
- ICS: An inhaled corticosteroid that can be used as maintenance therapy.
- ICS-LABA: A combination of an inhaled corticosteroid and a long-acting bronchodilator that can be used as a controller therapy or for reliever therapy.

Intervention Arm

Patients in the intervention arm will be issued two budesonide-formoterol (Symbicort®) 100/3 MDIs and a compatible spacer device such as Aerochamber Plus. One for home, maintenance and reliever doses, and the other to be kept at school for reliever use.

Dose Regimen

In clinical practice, treatment is stepped up and down depending on clinical control in terms of symptoms and risk of an asthma attack. Stepping up and down will be determined by the patient's usual healthcare provider. For those in the intervention arm participants will increase or decrease the number of maintenance budesonide-formoterol puffs per day. The control arm will continue as per usual clinical care whereby the ICS dose is increased or decreased and additional controllers such as LABAs are added or removed depending on clinical control.

The starting dose for budesonide-formoterol will be based on the dose of usual care inhaled corticosteroid ((ICS) low, medium or high) prescribed in the past 4 weeks as per current Global Initiative for Asthma (GINA) guidance:

For paediatric low-dose, moderate-dose and high-dose for budesonide-formoterol:

Beclomethasone dipropionate (ICS):

Standard particle CFC free inhalers: 100 - 200mcg/day; >200 - 400mcg/day; >400 - mcg/day Extra-fine particle CFC free inhalers: 50 - 100mcg/day; >100 - 200mcg/day; >200 - mcg/day

Budesonide:

Metered dose and dry powder inhalers:100 - 200mcg/day; >200 - 400mcg/day; >400 mcg/day

Fluticasone propionate:

Metered dose and dry powder inhalers: 100mcg/day; 150 – 200mcg/day; >200 mcg/day

Intervention Type

Drug

Phase

Phase III

Drug/device/biological/vaccine name(s)

budesonide-formoterol (Symbicort) [budesonide, Formoterol fumarate dihydrate]

Primary outcome(s)

Rate of severe asthma attacks: The rate of severe asthma attacks per patient per year, defined as worsening symptoms leading to an urgent medical review resulting in the prescription of systemic corticosteroids, measured at 52 weeks

Key secondary outcome(s))

- 1. Rate of total asthma attacks: The rate of total asthma attacks per patient per year, defined as worsening symptoms leading to an urgent medical review, whether or not systemic corticosteroids are prescribed, measured at 52 weeks
- 2. Time to first severe attack: Time to first severe asthma attack measured in weeks from baseline over the 52-week study period
- 3. Proportion of participants with severe attacks: Proportion of participants experiencing at least one severe asthma attack, measured at 52 weeks
- 4. Days lost from school: Number of days lost from school due to asthma, measured annually
- 5. Days lost from work for childcare: Number of days lost from work due to caring for a child with asthma, measured annually
- 6. Change in Childhood Asthma Control Test (cACT): Change in cACT scores measured from baseline to 52 weeks
- 7. Change in Paediatric Asthma Quality of Life Questionnaire (PAQLQ): Change in PAQLQ scores measured from baseline to 52 weeks
- 8. Growth during the study period: Change in height from baseline to study completion at 52 weeks
- 9. Total ICS dose: Total inhaled corticosteroid (ICS) dose as measured by an electronic monitoring device or prescription records at 52 weeks
- 10. Total systemic corticosteroid dose: Total dose of systemic corticosteroids prescribed over 52 weeks

- 11. Change in Child Health Utility (CHU9D): Change in CHU9D scores measured from baseline over the course of the 52-week study
- 12. Quality Adjusted Life Years (QALYs): QALYs calculated as the area under the CHU9D curve over 52 weeks
- 13. Cost of asthma-related healthcare: Total cost of asthma-related healthcare utilization measured over 52 weeks

Completion date

28/02/2027

Eligibility

Key inclusion criteria

- 1. Clinician diagnosis of asthma
- 2. Children aged ≥6 to <12 years
- 3. Washout period of 6 months post previous IMP studies
- 4. Prescribed asthma medication in the past 6 months (SABA, ICS or ICS+LABA)
- 5. Parent or caregiver able to understand the study requirements and willing to provide informed consent

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Child

Lower age limit

6 years

Upper age limit

12 years

Sex

All

Total final enrolment

0

Key exclusion criteria

- 1. Other chronic airway diseases including but not limited to bronchiectasis, cystic fibrosis, sickle cell disease
- 2. Already using ICS-formoterol as a reliever
- 3. Children on step 5, very high dose treatment (e.g. high dose ICS/LABA, prescription of biological therapy such as omalizumab)

Added 19/06/2025:

4. Any known or suspected contraindications to the medications

Date of first enrolment 01/12/2024

Date of final enrolment 27/02/2026

Locations

Countries of recruitment

United Kingdom

England

Scotland

Wales

Study participating centre Great Ormond Street Hospital for Children

Great Ormond Street London England WC1N 3JH

Study participating centre The Royal London Hospital

Whitechapel London England E1 1BB

Study participating centre Kings College Hospital

Mapother House De Crespigny Park Denmark Hill London England SE5 8AB

Study participating centre St Georges University Hospital Blackshaw Road

London England SW17 0QT

Study participating centre Evelina London Children's Hospital

Westminster Bridge Road London England SE1 7EH

Study participating centre Leicester Royal Infirmary

Infirmary Square Leicester England LE1 5WW

Study participating centre Nottingham Children's Hospital

Queen's Medical Centre, Derby Rd Nottingham England NG7 2UH

Study participating centre Addenbrookes Hospital

Hills Road Cambridge England CB2 0QQ

Study participating centre Jenny Lind Children's Hospital

Norfolk and Norwich University Hospital, Colney Lane Norwich England NR4 7UY

Study participating centre Oxford Children's Hospital

John Radcliffe Hospital, Headington Oxford England OX3 0AG

Study participating centre Birmingham Children's Hospital

Steelhouse Ln, Queensway Birmingham England B4 6NH

Study participating centre Staffordshire Children's Hospital at Royal Stoke

Royal Stoke University Hospital, Newcastle Road Stoke-on-Trent England ST4 6QG

Study participating centre Royal Alexandra Children's Hospital

North Dr, Brighton and Hove Brighton England BN2 5BE

Study participating centre Bristol Royal Hospital for Children

Paul O'Gorman Building Upper Maudlin Street St Michael's Hill Bristol England BS2 8BJ

Study participating centre Alder Hey Children's Hospital

E Prescot Rd Liverpool England L14 5AB

Study participating centre Royal Manchester Children's Hospital

Manchester Royal Infirmary, Oxford Rd Manchester England M13 9WL

Study participating centre Great North Children's Hospital

Victoria Wing, Royal Victoria Infirmary Newcastle upon Tyne England NE1 4LP

Study participating centre Leeds Children's Hospital

Leeds General Infirmary Clarendon Wing Leeds England LS1 3EX

Study participating centre Sheffield Children's Hospital

The University of Sheffield, Clarkson St, Broomhall Sheffield England S10 2TH

Study participating centre Southampton Children's Hospital

Southampton General Hospital Tremona Road Southampton England SO16 6YD

Study participating centre Noah's Ark Children's Hospital for Wales

University Hospital of Wales, Heath Park Way Cardiff Wales CF14 4XW

Study participating centre Westburn Medical Group

Foresterhill Health Centre Aberdeen Royal Infirmary Foresterhill Road Aberdeen Scotland AB25 2XE

Study participating centre Royal Hospital for Sick Children (Glasgow)

1345 Govan Road Glasgow Scotland G51 4TF

Sponsor information

Organisation

Imperial College London

ROR

https://ror.org/041kmwe10

Funder(s)

Funder type

Government

Funder Name

National Institute for Health and Care Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet		11/11/2025	11/11/2025	No	Yes
Study website			13/11/2025	No	No