

Effectiveness of 3D-guided piezocision-assisted orthodontic surgery (a minimally invasive surgical technique) in accelerating the correction of severely crowded teeth

Submission date 04/04/2021	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 07/04/2021	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 29/03/2023	Condition category Oral Health	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Orthodontic treatment uses appliances (usually braces) to correct the position of teeth. The length of comprehensive orthodontic treatment ranges between 18-30 months, depending on treatment options and individual characteristics. In addition, orthodontic treatment time ranges between 25-35 months for extraction therapies. Reducing orthodontic treatment time is one of the main goals for orthodontists due to problems such as root resorption (shortening of the roots), periodontal (gum) disease and caries (tooth decay) that are associated with prolonged treatment time. Many techniques have been introduced to speed up orthodontic tooth movement, both surgical and non-surgical. The surgical approach is the most clinically applied and most tested with known stable results. The invasiveness of these procedures might have limited their widespread acceptance among orthodontists and patients. Therefore, more conservative flapless corticotomy techniques have recently been proposed. These procedures can be accomplished in a reasonably short period that might cause less pain and discomfort, and may improve patient acceptance. Although various techniques of flapless corticotomy have been reported to be successful in practice, scientific evidence for their effectiveness so far has been limited to a few small studies. This study aims to provide evidence about the effectiveness of 3D-guided piezosurgery with a flapless technique to align crowded anterior (front) teeth.

Who can participate?

Healthy adults with severely crowded teeth

What does the study involve?

Participants are randomly allocated to one of two groups. The first group receive conventional orthodontic treatment without any surgery, whereas the second group receive piezocision-assisted orthodontic treatment. The duration of orthodontic treatments differs from one patient to another, and they will all be followed up until the end of the alignment phase.

What are the possible benefits and risks of participating?

The main benefit of participating is faster orthodontic treatment reducing problems such as root resorption, periodontal disease and caries. The risks are limited as 3D-guided surgical cuts will avoid harming teeth and vital structures.

Where is the study run from?

Damascus University (Syria)

When is the study starting and how long is it expected to run for?

August 2019 to August 2021

Who is funding the study?

Damascus University (Syria)

Who is the main contact?

Omar Gibreal

omar.gibreal@gmail.com

Contact information

Type(s)

Scientific

Contact name

Dr Omar Gibreal

Contact details

Damascus University

Mazzeah Highway

Damascus

Syria

97009

+963 (0)992271037

omar.gibreal@gmail.com

Additional identifiers

EudraCT/CTIS number

Nil known

IRAS number

ClinicalTrials.gov number

Nil known

Secondary identifying numbers

UDDS-OMFS-08-2019

Study information

Scientific Title

Evaluation of the efficacy of 3D-guided piezosurgery (a minimally invasive surgical technique) in accelerating orthodontic alignment

Study objectives

3D-guided piezosurgery accelerates orthodontic alignment

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 15/08/2019, scientific research committee in the faculty of dentistry at Damascus University (Mazze Highway, Damascus, Syria; +963 (0)113341864; manager@hcsr.gov.sy), ref: 505

Study design

Single-center interventional double-blinded randomized controlled clinical trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details to request a participant information sheet

Health condition(s) or problem(s) studied

Anterior teeth severe orthodontic crowding

Interventions

With the aid of a computer-generated list of random numbers, the recruited patients will be assigned to two parallel groups with a 1:1 allocation ratio. The first group receive conventional orthodontic treatment, whereas the second group receive piezocision-assisted orthodontic treatment. The allocation sequence will be concealed using sequentially numbered, opaque, sealed envelopes.

Piezosurgery will be performed on the anterior lower segment of the dental arch in order to accelerate the correction of mandibular anterior crowding. The device tip will be used to create small vertical incisions into the cortex of the dentoalveolar process of the lower anterior teeth guided by a 3D surgical guide. For the control group, traditional orthodontic treatment will be provided to the patients without any surgical interventions.

The duration of orthodontic treatments differs from one patient to another, they will be followed up until the end of the alignment phase.

Intervention Type

Procedure/Surgery

Primary outcome measure

1. Duration of tooth alignment measured at the end of treatment by calculating the time required time (in days) to achieve complete alignment of lower anterior teeth from the first day of treatment and up to 120-150 days of observation
2. Change in tooth alignment measured using the Little Index of Irregularity at 30 days after the onset of treatment
3. Change in tooth alignment measured using the Little Index of Irregularity at 60 days after the onset of treatment
4. Change in tooth alignment measured using the Little Index of Irregularity at last assessment when a complete alignment is achieved; this is expected between 90 to 120 days after the onset of treatment

Secondary outcome measures

1. Pain measured using the visual analogue score (VAS) at baseline, 12, 24 and 48 hours
2. Discomfort measured using the visual analogue score (VAS) at baseline, 12, 24 and 48 hours
3. Swelling measured using the visual analogue score (VAS) at baseline, 12, 24 and 48 hours

Overall study start date

15/08/2019

Completion date

01/08/2021

Eligibility

Key inclusion criteria

1. Adult healthy patients, male and female, age range 15-27 years
2. Severe crowding ≤ 7 (Little's irregularity index)
3. Permanent occlusion
4. All mandibular teeth exist (except third molars)
5. Good oral and periodontal health:
 - 5.1. Probing depth < 4 mm
 - 5.2. No radiographic evidence of bone loss
 - 5.3. Gingival index ≤ 1
 - 5.4. Plaque index ≤ 1

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

50

Total final enrolment

30

Key exclusion criteria

1. Medical problems that affect tooth movement (use of corticosteroid, NSAIDs)
2. Patients have anti indication for oral surgery (medical, social, psychological)
3. Presence of primary teeth in the mandibular arch
4. Missing permanent mandibular teeth (except third molars)
5. Poor oral hygiene or current periodontal disease:
 - 5.1. Probing depth ≥ 4 mm
 - 5.2. Radiographic evidence of bone loss
 - 5.3. Gingival index > 1
 - 5.4. Plaque index > 1
6. Patient had previous orthodontic treatment

Date of first enrolment

01/12/2019

Date of final enrolment

01/03/2020

Locations

Countries of recruitment

Syria

Study participating centre

Damascus University

Faculty of Dentistry

Mazzeah Highway

Damascus

Syria

97009

Sponsor information

Organisation

Damascus University

Sponsor details

MazzeH Highway
Damascus
Syria
-
+963 (0)114442727
info@damascusuniversity.edu.sy

Sponsor type
University/education

Website
<http://damasuniv.edu.sy/>

Funder(s)

Funder type
University/education

Funder Name
Damascus University

Results and Publications

Publication and dissemination plan
Planned publication in a high-impact peer-reviewed journal

Intention to publish date
07/10/2021

Individual participant data (IPD) sharing plan
The data-sharing plans for the current study are unknown and will be made available at a later date.

IPD sharing plan summary
Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		11/08/2022	15/08/2022	Yes	No
Results article		28/03/2023	29/03/2023	Yes	No