

DElayed COrd CLamping versus early cord clamping in preterm infants born between 24 and 34 weeks

Submission date 06/10/2013	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 10/12/2013	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
Last Edited 10/12/2013	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Prematurity is responsible for more than half of all neonatal deaths. Advances in neonatal care have dramatically improved survival of extremely premature infants but there remains a significant risk of handicap and disability in survivors and an associated social and economic burden. Although all births before 37 weeks of gestation are defined as preterm, most damage and death occurs in infants delivered before 34 weeks.

The optimal timing of cord clamping in preterm infants is still the subject of continuing debate, not only due to the potential risks (hyperbilirrubinemia, polycythemia and respiratory grunting) but also in order to hand neonates over the neonatal team for resuscitation.

There is a strong association between hypotension and intraventricular hemorrhage, patency of the ductus arteriosus and necrotizing enterocolitis. It seems that a decrease in blood volume at the time of delivery would negatively influence blood pressure. Furthermore, about 30-40% of very low birth weight (VLBW) infants experience some delay in motor functioning sometime during childhood. Hypovolemia secondary to early cord clamping, might be disruptive to the developing brain resulting in a motor delay.

The aim of this study is to help decide the best time to clamp the umbilical cord in preterm babies below 34 weeks, either early (within 10 seconds) or delayed cord clamping (45-60 minutes), analyzing any differences in blood test the first week of life and any neurosensory disability at the age of two-three years old.

Who can participate?

Women who are expected to give birth below 34 weeks of gestation.

What does the study involve?

Pregnant women will be randomly allocated to one of the two groups: cord clamping within 10 seconds or cord clamping 45-60 seconds after birth.

A blood test is taken 48 hours and 7 days after birth.

Around the age of two-three years old, the parents will be contacted to book an appointment with the pediatrician for a neurological follow up.

What are the possible benefits and risks of participating?

Risks to participants are minimal because the intervention on the babies it is unlikely to cause any damage. All the babies will be born in a very safety environment with paediatric assistance at birth-time and will be moved to the Neonatal Intensive Care Unit immediately, in case they need it.

Where is the study run from?

Department of Obstetrics and Gynecology-Neonatal Intensive Care Unit, Clinic University Hospital Virgen de la Arrixaca, Murcia (Spain)

When is the study starting and how long is it expected to run for?

The study started in February 2011 and will continue to recruit women until September 2014.

Who is funding the study?

Sistema Murciano de Salud, Spain

Who is the main contact?

Catalina De Paco Matallana

Katy.depaco@gmail.com

Contact information

Type(s)

Scientific

Contact name

Dr Catalina De Paco Matallana

Contact details

C/Alhelies 4. Edif. Al Andalus 3E

El Ranero

Spain

30009

+34676672617

katy.depaco@gmail.com

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

N/A

Study information

Scientific Title

Randomized study of delayed cord clamping versus early cord clamping in preterm infants born between 24 and 34 weeks

Acronym

DECOCLA

Study objectives

To evaluate the effect of timing of umbilical cord clamping on neonatal venous haematocrit, maternal postpartum haemorrhage and neurosensory disability at two-three years of age.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics Committee CEIC Clinic University Hospital Virgen de la Arrixaca; 21 December 2010

Study design

Randomized trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Diagnostic

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Topic: Reproductive Health, Disease: Reproductive Health & Childbirth, Paediatrics

Interventions

Participants are randomized to two groups:

1. Early cord clamping: clamping of the cord within ten seconds after birth
2. Delayed cord clamping: clamping of the cord 45-60 seconds after birth

Intervention Type

Other

Phase

Not Applicable

Primary outcome measure

Evaluation of neonatal hemoglobin, hematocrit and bilirubin levels within the first 7 days after birth

Secondary outcome measures

1. Neonatal hemoglobin, hematocrit and ferritin at six months of life will be evaluated by blood sampling
2. Neonatal complications (Intraventricular hemorrhage, necrotizing enterocolitis, retinopathy, sepsis, respiratory problems, days on ventilation or oxygen, need for phototherapy, transfusions) and days in the neonatal intensive care will be evaluated by medical history review
3. Cardiac output in the first week after birth will be measured by echocardiography
4. Blood loss in the mother (blood test 48 hours after birth)
5. Neurodevelopmental assessment of newborns at the age of two-three years in both groups of the study will be test by Bayley Scales of Infant Development

Overall study start date

01/02/2011

Completion date

01/09/2014

Eligibility

Key inclusion criteria

Women will be eligible for the study if they are likely to have a live birth before 34 weeks gestation, regardless of mode of birth or whether cephalic or breech presentation, singleton or dichorionic pregnancies.

Participant type(s)

Patient

Age group

Adult

Sex

Female

Target number of participants

100

Key exclusion criteria

1. Major fetal abnormalities (defined as those that are lethal or require prenatal or postnatal surgery)
2. Fetal growth restriction
3. Monochorionic twins
4. Maternal pathology (pregestational Diabetes, severe cardiopathy, etc)
5. Infectious disease, hypertension and/or preeclampsia, obstetrics complications (abruptio, etc)

Date of first enrolment

01/02/2011

Date of final enrolment

01/09/2014

Locations

Countries of recruitment

Spain

Study participating centre

C/Alhelies 4. Edif. Al Andalus 3E

El Ranero

Spain

30009

Sponsor information

Organisation

Clinic University Hospital Virgen de la Arrixaca (Spain)

Sponsor details

Carretera Cartagena-Madrid S/N

El Palmar (Murcia)

Spain

30120

+34 968395588

katy.depaco@gmail.com

Sponsor type

Hospital/treatment centre

ROR

<https://ror.org/058thx797>

Funder(s)

Funder type

Government

Funder Name

Murcia Health Service [Servicio Murciano de Salud (SMS)], Murcia (Spain)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration