

# Group intervention to reduce intimate partner abuse by men in substance use treatment

<b>Submission date</b> 15/01/2020	<b>Recruitment status</b> Stopped	<input checked="" type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
<b>Registration date</b> 20/01/2020	<b>Overall study status</b> Stopped	<input type="checkbox"/> Statistical analysis plan
		<input type="checkbox"/> Results
<b>Last Edited</b> 06/10/2021	<b>Condition category</b> Mental and Behavioural Disorders	<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

## Plain English summary of protocol

### Background and study aims

Intimate partner abuse (IPA) includes physical, sexual, psychological and financial abuse and/or controlling behaviours by a current/ex-partner. IPA impacts negatively on victims' health. Men who use substances are more likely be abusive towards their partners than men who do not. This study will recruit 378 men from substance use treatment in England, Wales and Scotland who have been abusive towards a female current or ex-partner in the past year. Men will be recruited by chance to the ADVANCE group intervention which aims to improve relationships and substance use treatment. The aims are to find out whether this new intervention is any better than usual substance use treatment with a keyworker, to compare the costs, to explore whether the intervention can be delivered in substance use treatment, and to find out whether men find the intervention acceptable, attend sessions, and stay in the study.

### Who can participate?

Men who have perpetrated abusive or violent behaviour towards a current or former female partner in the last 12 months.

### What does the study involve?

Male participants are randomly allocated to the intervention group or the control group. The intervention group attend 13 weekly 2-hour group sessions, along with substance use treatment as usual. The control group receive substance use treatment as usual (e.g., sessions with keyworker, group attendance and/or opiate therapy). The men's' current/ex-partners are offered support for IPA and are invited to take part in the study by providing information about their partner's behaviour and their own well-being. The man's keyworker and the women's support worker share information that relates to women's safety and risk. Women are updated about their current/ex-partner's overall progress in the intervention. Data on IPA, substance use, emotional well-being, quality of life and service use are collected from both men and women at the start and end of the intervention.

### What are the possible benefits and risks of participating?

The study will help to show whether the ADVANCE intervention helps men who use substances reduce their abusive and violent behaviour towards female partners. It is hoped that if this happens, this will improve the wellbeing of their partners and children. To ensure that women

and their children are safe, staff from the substance use treatment service and the women's support service worker will talk to each other on a regular basis to share information that relates to the women's safety and risk. Participating in the study could improve relationships and the wellbeing of women and children in the future. All participants are given the opportunity to get support for their relationship and are provided with a range of national and local contact numbers and services that will be able to help.

Where is the study run from?

King's College London (UK)

When is the study starting and how long is it expected to run for?

February 2020 to September 2021

Who is funding the study?

National Institute for Health Research (NIHR) (UK)

Who is the main contact?

Prof Gail Gilchrist

Gail.Gilchrist@kcl.ac.uk

## Contact information

### Type(s)

Scientific

### Contact name

Prof Gail Gilchrist

### ORCID ID

<https://orcid.org/0000-0002-5616-6283>

### Contact details

National Addictions Centre

King's College London

4 Windsor Walk

London

United Kingdom

SE5 8BB

+44 (0)7708255859

Gail.Gilchrist@kcl.ac.uk

## Additional identifiers

### Clinical Trials Information System (CTIS)

Nil known

### Integrated Research Application System (IRAS)

271242

### ClinicalTrials.gov (NCT)

Nil known

**Protocol serial number**

RP-PG-1214-20009, IRAS 271242, CPMS 44092

## Study information

**Scientific Title**

A randomised controlled trial of the ADVANCE integrated group intervention to reduce intimate partner abuse perpetration by men receiving treatment for substance use, compared to substance use treatment as usual

**Acronym**

ADVANCE

**Study objectives**

The ADVANCE integrated intervention to reduce intimate partner abuse (IPA) by men in treatment for substance use + substance use treatment as usual (TAU) will be more effective and cost effective than substance use treatment as usual only

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Approved 07/02/2020, Yorkshire and the Humber – Sheffield Research Ethics Committee (Barlow House, 3rd Floor, 4 Minshull Street, Manchester, M1 3DZ, UK; +44 (0)207 104 8208; nrescommittee.yorkandhumber-sheffield@nhs.net), ref: 19/YH/0431

**Study design**

Interventional multi-centre parallel group individually randomised controlled trial

**Primary study design**

Interventional

**Study type(s)**

Treatment

**Health condition(s) or problem(s) studied**

Intimate partner abuse and substance use

**Interventions**

Intervention condition:

The group intervention consists of 13 x 2-h group sessions + substance use treatment as usual. This includes a pre-group individual motivational and preparation session followed by 12 core group sessions.

Control condition:

Substance use treatment as usual (TAU) will be permitted for both treatment arms. This is

usually individual sessions with a keyworker or attendance at groups. It may also include opiate substitution treatment. Men randomly allocated to the control condition will receive only TAU. They will not receive the intervention at the end of the trial.

#### Randomisation methods:

Male participants will be allocated to the intervention group + TAU or TAU only (ratio 2:1) within service strata via an independent online system based at the King's Clinical Trials Unit (King's CTU) at King's College London. Allocation will be at the level of the individual participant and stratified by service. Randomisation will take place after a set of 24 men has been recruited in a service. The whole set will be randomised at the same time (16 men to the intervention arm and 8 to the control arm). A minimum set of 18 men will be required for randomisation to take place.

#### Delivery of intervention:

All sessions will take place in the substance use treatment service and will be facilitated by two facilitators (one male and one female): a substance use worker and a worker with experience of IPA where possible. The group intervention will be delivered to closed groups consisting of a maximum of 16 men. Men allocated to the control arm would not be able to join.

#### Follow up study:

To investigate the impact of the ADVANCE intervention measures will be collected at follow-up from female current or ex-partners at 4- and 12-months post randomisation of male current or ex-partner.

### **Intervention Type**

Behavioural

### **Primary outcome(s)**

Total score in self-reported perpetration of IPA by men in substance use treatment measured using the adapted 29-item Abusive Behavior Inventory (ABI) - partner form (perpetration) in the previous 4 months at 12 months post-randomisation

### **Key secondary outcome(s)**

Evaluated for men at 4 and 12 months post-randomisation:

1. Other IPA perpetration
  - 1.1. Abusive Behavior Inventory (ABI) - partner form (perpetration) in the previous 4 months at 4 months post randomisation only
  - 1.2. Controlling Behaviours Scale (partial) in the past 4 months
  - 1.3. Use of social media, stalking, locked in and Using children against partner in the past 4 months
2. Substance use
  - 2.1. Number of alcohol free days in past 28 days assessed using the Treatment Outcome Profile
  - 2.2. Number of drug free days in past 28 days assessed using the Treatment Outcome Profile
3. Mental Well-being
  - 3.1. Depressive symptoms assessed using the PHQ-9 in the past 2 weeks
  - 3.2. Anxiety symptoms assessed using the GAD-7 in the past 2 weeks
  - 3.3. PTSD symptoms assessed using the Primary Care PTSD Screen in the past month
  - 3.4. Propensity for Abusiveness Scale [anger subscale]
4. Self-Control: Brief Self-Control Scale
5. Quality of life will be measured using the EQ-5D-3L

6. Capability will be measured using the ICECAP-A

7. Service use and medication in past 4 months at 4 month follow-up and in the past 8 months at 12 month follow-up

8. Measures obtained to describe the therapy experience:

8.1. Therapeutic alliance

Men in the intervention arm and group facilitators will complete evaluation forms at the end of each session to assess the acceptability of the intervention. At the end of the intervention, men's perception of therapeutic alliance will be assessed using the Working Alliance Inventory – Short Revised and the California Psychotherapy Alliance Scale- Short Form

8.2. Nested process evaluation

A process evaluation will be undertaken to understand the intervention's functioning. Six priority areas will be assessed: context (local factors that influence implementation), fidelity (extent to which intervention is delivered as conceived), dose delivered (amount of intervention offered to participants), dose received (extent of participants' engagement in intervention), reach and recruitment. We will also assess how the intervention was delivered (e.g. the training and resources necessary to achieve full implementation), and the quantity and quality of what was delivered), intervention components (what is to be implemented), mechanisms of impact (the mechanisms through which an intervention will work) and intended outcomes. Treatment integrity (Treatment Design, Training Providers, Delivery of Treatment, Receipt of Treatment, and Enactment of treatment skills) will be measured on a random sample of 10% of video recorded sessions using a pre-determined checklist. All sessions will be video recorded.

9. To investigate the impact of the ADVANCE intervention in follow-up study, the following measures will be collected at follow-up from female current or ex-partners at 4- and 12-months post randomisation of male current or ex-partner:

9.1. IPA victimisation using the Revised Abusive Behavior Inventory (ABI-R) in the previous 4 months

9.2. Other IPA victimisation

9.2.1. Controlling Behaviours Scale (partial) in the past 4 months

9.2.2. Use of social media, stalking, locked in, and using children against partner in the past 4 months

9.3. Substance use

9.3.1. Number of alcohol free days in past 28 days using the Treatment Outcome Profile

9.3.2. Number of drug free days in past 28 days using the Treatment Outcome Profile

9.4. Mental Well-being

9.4.1. Depressive symptoms assessed using the PHQ-9 in the past 2 weeks

9.4.2. Anxiety symptoms assessed using the GAD-7 in the past 2 weeks

9.4.3. PTSD symptoms assessed using the Primary Care PTSD Screen in the past month

9.5. Quality of life will be measured using the EQ-5D-3L

9.6. Capability will be measured using the ICECAP-A

9.7. Service use and medication in past 4 months at 4 month follow-up and in the past 8 months at 12 month follow-up

## **Completion date**

30/09/2021

## **Reason abandoned (if study stopped)**

Pandemic

## **Eligibility**

## **Key inclusion criteria**

Men receiving community substance use treatment:

1. Male participant has perpetrated abusive or violent behaviour towards a current or ex female partner in the last 12 months
2. Contact with current or ex female partner at least once in the past 4 months – in person, or by phone/ text/ email/ social media
3. Plans to stay in current location for the next 6 months
4. Agrees to provide contact details of current and/or ex female partner
5. Ability to understand and communicate in English
6. Able to attend the intervention (believe that the proposed day and time the group is planned to be delivered is suitable)
7. Substance use treatment service assesses as suitable to participate in the trial

Current or ex-female partners of men in the trial:

1. Current or ex-partner participating in the trial
2. Aged 18 years or older
3. Lives in the UK
4. Ability to understand and communicate in English

## **Participant type(s)**

Patient

## **Healthy volunteers allowed**

No

## **Age group**

Adult

## **Lower age limit**

18 years

## **Sex**

All

## **Key exclusion criteria**

Men receiving community substance use treatment

1. Male reporting current order preventing him from contacting current or ex female partner
2. Currently attending an intervention for IPA
3. Previously attended the ADVANCE intervention for IPA
4. Participant is not/ no longer attending the substance use treatment service
5. Other safety concerns that may put the female partner at risk. These will be considered on a case by case basis by the research team and the substance use treatment service e.g. where both participants share a mobile phone number, the male participant has a court case pending for IPA or there is a child protection hearing pending.

Current or ex-female partners of men in the trial

1. Current order preventing her from contacting current or ex male partner recruited to the trial
2. Other safety concerns that may put the male partner at risk. These will be considered on a case by case basis by the research team and the clinical team e.g. where both participants share a mobile phone number, the female participant has a court case pending for IPA or there is a

child protection hearing pending.

3. Female partner discloses that there is an order preventing her male current or ex-partner from contacting her (i.e. contradicting what he has said in his screening interview). In such cases the man would not be withdrawn, unless the clinical team felt there was an increased risk to either party in his continuing in the study.

Male partners and non-English speaking female partners will not be eligible to take part in the trial but will be offered support for their IPA victimisation.

If a female partner is excluded because she has a current order preventing her from contacting current or ex male partner – her current or ex male partner will remain in the trial.

**Date of first enrolment**

03/02/2020

**Date of final enrolment**

30/09/2020

## **Locations**

**Countries of recruitment**

United Kingdom

England

Scotland

Wales

**Study participating centre**

**King's College London**

Institute of Psychiatry, Psychology and Neuroscience

National Addiction Centre

4 Windsor Walk

London

United Kingdom

SE5 8BB

**Study participating centre**

**University of Bristol**

Centre for Academic Primary Care Population Health Sciences

Bristol Medical School

Canynges Hall

39 Whatley Road

Bristol

United Kingdom

BS8 2PS

**Study participating centre**  
**University of Edinburgh**  
School of Health in Social Sciences  
8-9 Hope Park Square  
Edinburgh  
United Kingdom  
8HQ 9NW

## Sponsor information

**Organisation**  
King's College London

**ROR**  
<https://ror.org/0220mzb33>

**Organisation**  
South London and Maudsley NHS Foundation Trust

**ROR**  
<https://ror.org/015803449>

## Funder(s)

**Funder type**  
Government

**Funder Name**  
National Institute for Health Research

**Alternative Name(s)**  
National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

**Funding Body Type**  
Government organisation

**Funding Body Subtype**  
National government



## Location

United Kingdom

# Results and Publications

## Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available due to confidentiality.

## IPD sharing plan summary

Not expected to be made available

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Participant information sheet</a>	Participant information sheet	11/11/2025	11/11/2025	No	Yes
<a href="#">Study website</a>	Study website	11/11/2025	11/11/2025	No	Yes