# Nordic Long-term Obsessive compulsive disorder (OCD) Treatment Study

| Submission date               | Recruitment status  No longer recruiting             | <ul><li>Prospectively registered</li></ul> |  |  |
|-------------------------------|--|--|--|--|
| 20/10/2007                    |  | ☐ Protocol                                 |  |  |
| Registration date             | Overall study status                                 | Statistical analysis plan                  |  |  |
| 19/03/2008                    | Completed  | [X] Results                                |  |  |
| <b>Last Edited</b> 10/07/2025 | Condition category  Mental and Behavioural Disorders | Individual participant data                |  |  |

#### Plain English summary of protocol

Background and study aims

Pediatric Obsessive Compulsive Disorder (OCD) is a serious and often chronic disorder involving obsessive and excessive fears, and behaviours (i.e., rituals) that aim to neutralize the fears and dangers. The symptoms lead to impairment and reduced quality of life. There is impressive evidence for the effectiveness of Cognitive Behaviour Therapy (CBT) with exposure to the feared situations and response prevention of the rituals. Moreover, CBT gives good symptom relief in many, better than drug treatments (i.e., serotonin reuptake inhibiting drugs [SSRIs]). SSRI treatment also has an impressive evidence base. However, little is known about what treatments to offer in children and adolescents who do not benefit from CBT. So, our study aims at investigating whether continued CBT or a switch to sertraline (an SSRI) is best in these non-responding children and adolescents. However, we are also interested in investigating whether regular child and adolescent clinicians can be taught such CBT and be as efficient as psychotherapists working in specialized OCD clinics.

#### Who can participate?

Children and adolescents, aged 7-17 years, with moderate to severe OCD.

#### What does the study involve?

Following a thorough baseline diagnostic work-up, patients participate in 14 sessions of CBT. Assessments are made also at the 7th and 13th weeks of the therapy. Youngsters who do not benefit from CBT are then randomly allocated to either sertraline or to continued CBT.

#### What are the possible benefits and risks of participating?

The participants receive state-of-the-art CBT and drug treatment which increase the chance of symptom relief, while the risks are small.

#### Where is the study run from?

Three Scandinavian countries (Sweden, Norway and Denmark) contribute. The Center for Child and Adolescent Mental Health, Eastern and Southern Norway (RBUP) is the data center.

When is the study starting and how long is it expected to run for? The study started in 2008, has now stopped inclusion, and is currently working on a long-term follow-up scheme to study whether gains from therapy are durable.

#### Who is funding the study?

All participating centers finance their own contribution with the aid of the local hospital or grants from research foundations. At RBUP, grants from research foundations have covered costs associated with graduate students. RBUP and the Norwegian Research Council have contributed to an electronic data capture system.

Who is the main contact? Dr Tord Ivarsson tord.ivarsson@r-bup.no

# Contact information

#### Type(s)

Scientific

#### Contact name

Dr Tord Ivarsson

#### Contact details

Gullhaug Torg 4B Oslo Norway 0484 +47 (0)22 58 60 00 tord.ivarsson@r-bup.no

### Additional identifiers

#### Protocol serial number

NordLOTS protocol 1.2

# Study information

#### Scientific Title

Nordic Long-term Obsessive compulsive disorder (OCD) Treatment Study

#### Acronym

**NordLOTS** 

#### **Study objectives**

Children and adolescents with obsessive compulsive disorder (OCD) who do not respond to a course of cognitive behaviour therapy (CBT) will benefit equally from sertraline and from continued CBT. Identification of CBT versus sertraline responders is possible. Non responders to CBT and sertraline will benefit from aripiprazol augmentation.

#### Ethics approval required

#### Old ethics approval format

#### Ethics approval(s)

- 1. Denmark, Institutional Review Board (IRB) (Den videnskabsetiske komité for Region Midtivlland), ref: 20070140
- 2. Sweden, IRB, 04/02/2008
- 3. Norway, IRB, 10/03/2008

#### Study design

Randomized active controlled trial with three steps:

- 1: Open uncontrolled
- 2: Randomised and controlled
- 3: Open uncontrolled

#### Primary study design

Interventional

#### Study type(s)

Treatment

#### Health condition(s) or problem(s) studied

Obsessive compulsive disorder

#### **Interventions**

Step 1: Cognitive behaviour therapy (CBT)

Step 2: Sertraline plus CBT support (less intensive CBT) or intensive CBT

Step 3: Sertraline plus CBT support plus aripiprazol

Non-responders to CBT are randomised to continued CBT or sertraline with CBT support. CBT plus sertraline non-responders are treated un-controlled with aripiprazol. Outcome is studied for 36 months.

#### Dosing schedule of sertraline:

Week 0: no dose given

Week 1: 25 mg for 3 days, then 50 mg (range: 25 - 50 mg)

Week 2: 75 mg (range: 50 - 75 mg)

Weeks 3 - 4: 100 mg (range: 75 - 100 mg)

Weeks 5 - 7: 150 mg (range: 75 - 150 mg)

Weeks 8 - 12: 200 mg (range: 75 - 200 mg)

Weeks 6 - 12: 200 mg (range: 75 - 200 mg)

#### Dosing schedule of aripiprazol:

Week 0: no dose given

Week 1: 2.5 mg for 7 days (range: 2.5 mg)

Weeks 2 - 4: 5 mg (range: 2.5 - 5 mg)

Weeks 5 - 7: 7.5 mg (range: 2.5 - 7.5 mg)

Weeks 8 - 12: 10 - 20 mg (range: 2.5 - 20 mg)

Weeks 12 onwards: 2.5 - 20 mg

#### Intervention Type

Drug

#### Phase

Not Applicable

#### Drug/device/biological/vaccine name(s)

Sertraline, aripiprazole

#### Primary outcome(s)

- 1. CYBOCS
- 2. Clinical Global Impression Scale
- 3. Clinical Global Improvement Scale
- 4. Children's OCD Impact Scale

Outcomes measured (approximately) at weeks 0, 7, 13, and months 6, 12, 24, 36.

#### Key secondary outcome(s))

- 1. Screen for Child Anxiety Related Disorders Revised (SCARED-R)
- 2. Mood and Feelings Questionnaire (MFQ)
- 3. Children's Global Assessment Scale (CGAS)
- 4. Child Behaviour Checklist (CBCL)
- 5. Family Accomodation Scale (FAS)
- 6. Need to add another treatment

Outcomes measured (approximately) at weeks 0, 7, 13, and months 6, 12, 24, 36.

#### Completion date

31/12/2013

# Eligibility

#### Key inclusion criteria

- 1. Patients 7 17 years of age
- 2. Moderate-severe obsessive compulsive disorder according to Diagnostic and Statistical Manual of Mental Disorders Fourth Edition (DSM IV). Severity is defined by Children's Yale-Brown Obsessive Compulsive Scales (CY-BOCS) scores of 16 or above

#### Participant type(s)

Patient

#### Healthy volunteers allowed

No

#### Age group

Child

#### Lower age limit

7 years

#### Upper age limit

17 years

#### Sex

Αll

#### Total final enrolment

269

#### Key exclusion criteria

- 1. Co-morbidity has not higher treatment priority (e.g. psychosis, anorexia nervosa, severe depression with suicidality, an autistic disorder or Asperger's syndrome)
- 2. Pervasive developmental disorders (PDD) not otherwise specified (NOS) is allowed if Clinical Global Impression (CGI) score for the PDD is less than or equal to 3 and CGI for the PDD NOS is less than or equal to CGI for the OCD
- 3. Mental retardation (intelligence quotient [IQ] less than 70)
- 4. Patients have not been treated with selective serotonin reuptake inhibitor (SSRI) or CBT for their OCD during the last year
- 5. If the patient is of non-Nordic ethnicity both the patient and one parent must speak a Nordic language

#### Date of first enrolment

01/01/2008

#### Date of final enrolment

31/12/2013

#### Locations

#### Countries of recruitment

Denmark

Norway

Sweden

# Study participating centre Gullhaug Torg 4B

Oslo Norway 0484

# Sponsor information

#### Organisation

The Centre for Child and Adolescent Mental Health, Eastern and Southern Norway (R.BUP) (Norway)

#### **ROR**

https://ror.org/042s03372

# Funder(s)

#### Funder type

Research organisation

#### Funder Name

The Centre for Child and Adolescent Mental Health, Eastern and Southern Norway (Regionsenter for Barn og Unges Psykiske helse [R.BUP]) (Norway) - for the Nordic coordination

#### **Funder Name**

The participating clinics finance their participation from local funding agencies

# **Results and Publications**

Individual participant data (IPD) sharing plan

#### IPD sharing plan summary

Not provided at time of registration

#### **Study outputs**

| Output type                   | Details                           | Date<br>created | Date<br>added  | Peer<br>reviewed? | Patient-<br>facing? |
|-------------------------------|-----------------------------------|-----------------|----------------|-------------------|---------------------|
| Results article               |                                   | 29/10/2014      |                | Yes               | No                  |
| Results article               |                                   | 01/03/2015      |                | Yes               | No                  |
| Results article               |                                   | 01/11/2019      | 21/04<br>/2020 | Yes               | No                  |
| Results article               |                                   | 01/04/2021      | 19/04<br>/2021 | Yes               | No                  |
| Results article               | Quality of life results           | 01/04/2021      | 22/04<br>/2021 | Yes               | No                  |
| Results article               | Relapse rates following remission | 03/12/2023      | 11/12<br>/2023 | Yes               | No                  |
| Results article               |                                   | 09/07/2025      | 10/07<br>/2025 | Yes               | No                  |
| Participant information sheet | Participant information sheet     | 11/11/2025      | 11/11<br>/2025 | No                | Yes                 |