

Randomised comparison of intermittent urethral and indwelling suprapubic catheterisation in the management of voiding after urogynaecological surgery

Submission date 28/05/2009	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 09/07/2009	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 01/05/2012	Condition category Urological and Genital Diseases	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Mr Paul Hilton

Contact details
Directorate of Women's Services
Level 3, Leazes Wing
Royal Victoria Infirmary
Newcastle upon Tyne
United Kingdom
NE1 4LP

Additional identifiers

Protocol serial number
N/A

Study information

Scientific Title

Intermittent urethral versus indwelling suprapubic catheterisation in the management of voiding after urogynaecological surgery: a randomised single centre controlled trial

Study objectives

The aim of this study was to investigate the hypothesis that intermittent catheterisation (IC) is associated with a more rapid return to normal micturition following urogynaecological surgery by undertaking a randomised comparison of IC with suprapubic catheterisation in women undergoing surgery for urodynamic stress incontinence or utero-vaginal prolapse.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Newcastle and N. Tyneside Local Research Ethics Committees approved on the 20th January 2004 (ref: 2003/155)

Study design

Single centre randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Urodynamic stress incontinence, utero-vaginal prolapse

Interventions

All women electively admitted for surgery for urodynamic stress incontinence or pelvic organ prolapse were approached with a view to randomisation. A trial information leaflet was provided and those agreeing to participate completed a trial consent form in addition to their surgical consent. They were randomised into one of two groups using opaque sealed envelopes, opened prior to surgery by the consenting surgeon. No blinding of patient, surgeon, nurses nor outcomes assessor was feasible. The two randomisation groups were as follow:

Group 1: bladder drainage by a suprapubic catheter inserted in theatre. The catheter was left on free drainage for 48 hours post-operatively before commencing clamping

Group 2: catheterised intermittently post-operatively

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Length of hospital stay, measured from day of admission to day of discharge with a range of between 2 - 19 days.

Key secondary outcome(s)

1. The time to resume normal voiding (defined as voided volumes greater than 200 ml and post-void residual volumes consistently less than 100 ml), recorded within the time of the hospital stay
2. The number of episodes of urinary tract infection (UTI) (defined by catheter-specimen urine [CSU] or mid-stream urine [MSU] showing a single bacterium growing at a colony count greater than 100,000 colony forming units per ml), recorded within the time of the hospital stay
3. Patient experience of catheterisation as determined from a questionnaire given to patients at the end of their hospital stay, recorded within the time of the hospital stay, prior to discharge

Completion date

01/07/2004

Eligibility

Key inclusion criteria

All women electively admitted for surgery for urodynamic stress incontinence or pelvic organ prolapse. No age limits.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Other

Sex

Female

Key exclusion criteria

1. Women undergoing surgery where post-operative catheterisation is not routinely employed
2. Women requiring continuous post-operative bladder drainage, e.g. following repair of vesicovaginal fistula, urethral diverticulectomy, augmentation cystoplasty and operative bladder injury

Date of first enrolment

01/04/2004

Date of final enrolment

01/07/2004

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
Directorate of Women's Services
Newcastle upon Tyne
United Kingdom
NE1 4LP

Sponsor information

Organisation
Newcastle upon Tyne Hospitals NHS Foundation Trust (UK)

ROR
<https://ror.org/05p40t847>

Funder(s)

Funder type
Other

Funder Name
Investigator initiated and funded (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary
Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/10/2010		Yes	No