# Umbilical cord blood transfusion for children with severe anaemia (Wazo Geni study)

Submission date	<b>Recruitment status</b> No longer recruiting	Prospectively registered	
01/07/2007		☐ Protocol	
<b>Registration date</b> 05/07/2007	Overall study status Completed	Statistical analysis plan	
		[X] Results	
Last Edited	Condition category	Individual participant data	
22/12/2015	Haematological Disorders		

#### Plain English summary of protocol

Not provided at time of registration

### Contact information

#### Type(s)

Scientific

#### Contact name

Dr Oliver Hassall

#### Contact details

KEMRI/Wellcome Trust Research Laboratories PO Box 230 Kilifi Kenya 80108

00

ohassall@kilifi.kemri-wellcome.org

## Additional identifiers

#### Protocol serial number

SSC 1215

## Study information

#### Scientific Title

An unmasked, single arm trial to assess harm, safety and efficacy of umbilical cord red blood cell transfusion for children with severe anaemia in a Kenyan hospital

#### **Study objectives**

Please note that as of 16/02/2009 this record was updated to include an amendment of the protocol from packed cord red blood cell transfusion to umbilical cord blood transfusion. All updates can be found in the relevant field under the above update date. The initial title (one title only) at the time of registration was: 'The safety and efficacy of packed cord red blood cell transfusion in children with severe anaemia in a Kenyan hospital'. Please also note that the anticipated end date has also changed; the initial end date at the time of registration was 31/03/2008.

#### Current hypothesis as of 16/02/2009:

Transfusion of umbilical cord red blood cells is safe and efficacious in the management of children with severe anaemia requiring blood transfusion in a Kenyan hospital.

#### Initial information at time of registration:

Transfusion of packed cord red blood cells is safe and efficacious in the management of children with severe anaemia requiring blood transfusion in a Kenyan hospital.

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

Approval received from:

- 1. KEMRI/National Ethics Committee on the 6th June 2007 (ref: SSC 1215)
- 2. Liverpool School of Tropical Medicine Research Ethics Committee on the 18th April 2007 (ref: 07.15)

#### Study design

Phase 1b, open-label, non-randomised, non-controlled, single arm trial

#### Primary study design

Interventional

#### Study type(s)

Treatment

#### Health condition(s) or problem(s) studied

Severe anaemia

#### **Interventions**

Amended as of 16/02/2009:

Umbilical cord red blood cells from a maximum of two cord blood donations to provide a minimum quantity of haemoglobin equivalent to 20 ml/kg of adult-donated whole blood. Transfusions will be over 3 - 4 hours, with or without frusemide according to current international guidelines.

There will be one transfusion episode per child. If further transfusions are required the child will receive standard management. Follow-up is for 28 days.

#### Initial information at time of registration:

A volume of packed cord blood cells from up to two cord blood units to be transfused to provide a minimum quantity of haemoglobin equivalent to 20 ml/kg of adult-donated whole blood.

Transfusions will be over 3 - 4 hours, with or without frusemide according to current international guidelines.

There will be one transfusion episode per child. If further transfusions are required the child will receive standard management. Follow-up is for 28 days.

#### Intervention Type

Drug

#### **Phase**

Phase I/II

#### Drug/device/biological/vaccine name(s)

Umbilical cord blood transfusion

#### Primary outcome(s)

Amended as of 16/02/2009:

- 1. Serious adverse events
- 2. Suspected unexpected serious adverse reactions
- 3. Adverse events

Initial information at time of registration:

- 1. Serious adverse events
- 2. Suspected unexpected serious adverse reactions

Monitored throughout in-patient stay with formal assessments at the following times:

- 1. Pre-transfusion
- 2. During transfusion
- 3. Two hours post transfusion
- 4. 24 hours post transfusion
- 5. At hospital discharge
- 6. 28 days post transfusion

#### Key secondary outcome(s))

Rise in haemoglobin at 24 hours and 28 days.

#### Completion date

17/06/2008

## **Eligibility**

#### Key inclusion criteria

- 1. Children aged 12 years or less, either sex
- 2. Children with severe anaemia for whom a blood transfusion is indicated (aged less than three months, Haemoglobin [Hb] less than or equal to 10 g/dL; aged greater than three months, Hb less than or equal to 4 g/dL)

#### Participant type(s)

**Patient** 

#### Healthy volunteers allowed

#### Age group

Child

#### Upper age limit

12 years

#### Sex

Αll

#### Key exclusion criteria

- 1. Coma (Blantyre Coma Scale less than or equal to 2)
- 2. Prostration (if unable to sit when well, inability to take enteral feeds; if able to sit when well, inability to sit unsupported)
- 3. Uncompensated shock
- 4. Compensated shock (capillary refill time greater than 3 seconds; temperature gradient)
- 5. Respiratory distress (deep breathing)
- 6. Neonatal hyperbilirubinaemia requiring exchange transfusion
- 7. Any other marker of clinical severity considered to preclude the child from recruitment into the study
- 8. Enrolment in another intervention trial
- 9. Children for whom informed consent to enter the study is not possible or not given

#### Date of first enrolment

25/06/2007

#### Date of final enrolment

17/06/2008

## Locations

#### Countries of recruitment

Kenya

#### Study participating centre KEMRI/Wellcome Trust Research Laboratories

Kilifi

Kenva

80108

# Sponsor information

#### Organisation

Liverpool School of Tropical Medicine (UK)

#### **ROR**

https://ror.org/03svjbs84

## Funder(s)

#### Funder type

Charity

#### Funder Name

The Wellcome Trust (UK) - Training Fellowship (grant ref: 073604)

## **Results and Publications**

Individual participant data (IPD) sharing plan

#### IPD sharing plan summary

Not provided at time of registration

#### **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/03/2015		Yes	No