Temporary occlusion of the hypogastric artery to reduce blood loss associated with laparoscopic myomectomy

Submission date	Recruitment status No longer recruiting	Prospectively registered		
27/06/2023		Protocol		
Registration date	Overall study status	[X] Statistical analysis plan		
03/07/2023	Completed	[X] Results		
Last Edited	Condition category	Individual participant data		
07/09/2023	Surgery			

Plain English summary of protocol

Background and study aims

Uterine leiomyomas, or uterine fibroids, are benign tumors in the pelvic region that are commonly found in women. For women of reproductive age experiencing symptoms from these tumors, laparoscopic myomectomy (LM) is the preferred treatment option in which myomas are surgically removed through small incisions in the abdomen. In this study, the authors propose that combining LM with blocking blood flow of arteries in the pelvis, or temporary occlusion of the hypogastric artery (TOHA) using vascular clips, can reduce uterine blood flow during surgery and effectively decrease surgery-related blood loss.

Who can participate?

Adult women aged 18-49 years old diagnosed with intramural uterine leiomyomas larger than 4 cm in diameter, who preferred LM and wished to preserve fertility.

What does the study involve?

Participants were randomized into two parallel groups:

- 1. Patients undergoing standard LM
- 2. Patients undergoing LM with TOHA clipping.

What are the possible benefits and risks of participating?

The main benefits regard reduced surgery-associated blood loss. There are no foreseen risks associated with the intervention.

Where is the study run from?

Ob-Gyn Clinic of Emergency Municipal Clinical Hospital in Timisoara, Romania. The clinic is affiliated with "Victor Babes" University of Medicine and Pharmacy, Timisoara, Romania.

When is the study starting and how long is it expected to run for? January 2018 to December 2021

Who is funding the study?
"Victor Babes" University of Medicine and Pharmacy, Timisoara, Romania

Who is the main contact? Prof Laurentiu Pirtea (Principal investigator), pirtea.laurentiu@umft.ro Dr Cristina Secosan, secosan.cristina@umft.ro

Contact information

Type(s)

Principal investigator

Contact name

Prof Laurentiu Pirtea

ORCID ID

https://orcid.org/0000-0002-4398-2492

Contact details

Victor Babeş University of Medicine and Pharmacy Timişoara Piata Eftimie Murgu 2 Timisoara Romania 300041 +40756162282 pirtea.laurentiu@umft.ro

Type(s)

Scientific

Contact name

Dr Diana Lungeanu

ORCID ID

https://orcid.org/0000-0002-0274-1377

Contact details

Victor Babeş University of Medicine and Pharmacy Timişoara Piata Eftimie Murgu 2 Timisoara Romania 300041 +40722488775 dlungeanu@umft.ro

Type(s)

Scientific

Contact name

Dr Cristina Secosan

ORCID ID

https://orcid.org/0000-0003-1355-3790

Contact details

Victor Babeş University of Medicine and Pharmacy Timişoara Piata Eftimie Murgu 2 Timisoara Romania 300041 +40762647210 secosan.cristina@umft.ro

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

Nil known

Study information

Scientific Title

Investigation of bilateral Temporary Occlusion of the Hypogastric Artery in women undergoing laparoscopic myomectomy to minimize consequences of blood loss

Acronym

TOHAs

Study objectives

Combining laparoscopic myomectomy with a bilateral temporary occlusion of the hypogastric artery (TOHA) using vascular clips minimizes uterine blood flow during surgery and can significantly reduce surgery-associated blood loss.

Ethics approval required

Ethics approval required

Ethics approval(s)

approved 10/12/2018, Human Ethical Committee of "Victor Babes" University of Medicine and Pharmacy (Piata Eftimie Murgu 2, Timisoara, 300041, Romania; +40 256466001; cecs@umft.ro), ref: 44/10.12.2018

Study design

Proof-of-concept single-center interventional single-blinded randomized controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Laparoscopic myomectomy in adult women

Interventions

Combining laparoscopic myomectomy with temporary bilateral occlusion of the hypogastric artery.

Study arms:

There are two arms that include participants who are randomly allocated to the following interventions:

LM – laparoscopic myomectomy with no hypogastric artery clipping

LM + TOHA – laparoscopic myomectomy with temporary occlusion of the hypogastric artery

Method of randomization:

Randomization was single-blind and performed with the R package "blockrand" version 1.5.

Intervention providers:

All surgical procedures are performed by a single and highly proficient surgeon with over a decade of experience in laparoscopic surgery.

Modes of delivery

Prior to the surgery, neither Gonadotropin-Releasing Hormone (GnRH) agonists nor any intraoperative hemostatic drugs, such as vasopressin injections, are employed. Before initiating
surgical procedures, general anesthesia was induced concomitantly with orotracheal intubation.
Abdominal access was obtained via a direct trocar entry technique. CO2 insufflation was used to
create a pneumoperitoneum with a low intra-abdominal pressure of up to 12 mmHg. A 10 mm
trocar was positioned on the median line, 8 cm away from the umbilicus, while two 5 mm trocars
are situated on each side of the lower abdomen. The patient was then reclined in a
Trendelenburg position. The parietal peritoneum was incised below the lumbo-ovarian ligament.
The ureter and anterior trunk of the hypogastric artery are identified in the area where these
two structures run parallel. The ureter and hypogastric artery are identified using blunt
dissection. A metallic clip was then placed on the anterior trunk of the hypogastric artery,
cranially to the uterine artery emergence.

The uterine wall was incised, and the myoma/s are located and then removed using traction and countertraction maneuvers. Next, in-bag morcellation was used to remove the leiomyoma/s. The uterine wall was sutured with a double-layer stitch, utilizing a 29 mm, 3/8 circle needle, and Vicryl 2.0 thread. Once the uterine wall had been closed, hemostatic clips are gently removed by traction using atraumatic Johan fenestrated forceps.

Location where the intervention occurred

All surgical interventions are performed within the facilities of the Ob-Gyn Clinic of Emergency Municipal Clinical Hospital in Timisoara, Romania. The clinic is a tertiary referral hospital and is affiliated with "Victor Babes" University of Medicine and Pharmacy, Timisoara, Romania.

Intervention Type

Procedure/Surgery

Primary outcome(s)

Total hemoglobin concentration (g/dL) measured using a blood gas analyzer system (carried out in the CO-oximetry module) within the hospital laboratory before and immediately after the surgery

Key secondary outcome(s))

- 1. Required iron perfusion measured using the quantity of iron administered by intravenous infusion after surgery decided by the consultant based on ferritin concentration $< 100 \ \mu g$.I-1 after surgery
- 2. Required postoperative blood transfusion measured using blood testing and determined by the consultant based on a hemoglobin level of 70-80 g.I-1 measured after surgery
- 3. Anemia measured using blood testing and diagnosed based on hemoglobin less than 120 g/L (12 g/dL) as a dichotomous variable after surgery
- 4. Overall operative time measured using medical records as the time elapsed from the initial incision to final skin closure after surgery
- 5. Twelve-month post-operative spontaneous pregnancy measured using medical records as any pregnancy recorded within 12 months after laparoscopic myomectomy, without assisted reproductive measures at one timepoint at the end of the study
- 6. Hospital re-admission rate of participants in the trial who return to the hospital within seven days of discharge after laparoscopic myomectomy measured using medical records at one timepoint at the end of the study
- 7. Reports of death, including in-hospital death, or within 14 days of discharge after laparoscopic myomectomy, measured using medical records at one timepoint at the end of the study

Completion date

31/12/2021

Eligibility

Key inclusion criteria

- 1. Aged between 18 and 49 years old
- 2. Preference for laparoscopic myomectomy and their desire to preserve fertility
- 3. Intramural uterine leiomyomas greater than 4 cm in diameter

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Upper age limit

49 years

Sex

Female

Total final enrolment

60

Key exclusion criteria

- 1. Patients who did not agree to the enrollment or did not pass inclusion criteria (such as: age over 50, no preference for fertility preservation, personal option for hysterectomy)
- 2. Other types of myomas (such as submucosal or sub-serosal location), or intramural myomas under 4 cm which did not have an impact on the uterine cavity
- 3. Suspected of malignancy

Date of first enrolment

01/01/2019

Date of final enrolment

31/12/2020

Locations

Countries of recruitment

Romania

Study participating centre Emergency Municipal Clinical Hospital

Strada Hector 2A Timisoara Romania 300231

Sponsor information

Organisation

Victor Babes University of Medicine and Pharmacy Timisoara

ROR

https://ror.org/00afdp487

Funder(s)

Funder type

University/education

Funder Name

Victor Babeş University of Medicine and Pharmacy Timişoara

Results and Publications

Individual participant data (IPD) sharing plan

The dataset generated during this study will be made available upon institutional contact and reasonable request from Dr Cristina Secosan, secosan.cristina@umft.ro. These datasets will include individual patient data meta-analysis (IPD meta-analysis).

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		22/08/2023	07/09/2023	Yes	No
Participant information sheet	Informed consent [English]		03/07/2023	No	Yes
Participant information sheet	Informed consent [Romanian]		03/07/2023	No	Yes
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Statistical Analysis Plan			03/07/2023	No	No