The potential role of B-Type natriuretic peptide (BNP) and high-sensitivity troponin t(hs-tnt) prescreening by GPs to better target rapid access chest pain clinic referrals

Submission date	Recruitment status No longer recruiting	Prospectively registered		
29/12/2009		Protocol		
Registration date 06/04/2010	Overall study status Completed	Statistical analysis plan		
		[X] Results		
Last Edited 13/02/2017	Condition category Circulatory System	[] Individual participant data		
13/02/201/	Circulatory System			

Plain English summary of protocol

Background and study aims

At present the most accurate way of telling whether chest pain is coming from the heart or from any other cause such as muscular strain is by doing a blood test 12 hours after the chest pain starts. This test is called the Troponin T test. There is a newer blood test that may be able to provide doctors with the same information much quicker so that a decision on your care can be made faster and more accurately. There is evidence from previous studies that this may be so in patients with other heart conditions. This test is called the B-Type Natriuretic Peptide (BNP) test which is the name of a hormone that is produced by the heart muscle. The aim of this study is to assess the accuracy of this test in patients who present to hospital with chest pain.

Who can participate?

Adults aged between 20 and 93 presenting to the rapid Access Chest Pain Clinic with heart-related chest pain.

What does the study involve?

Participants provide background information and then have a blood sample taken while they are having their routine blood tests when they are admitted to hospital. Participants are followed up for 12 months in order to find out if they are admitted to hospital again for chest pain.

What are the possible benefits and risks of participating?

There are no direct benefits involved with participating. There is a small risk of discomfort or bruising from the blood test.

Where is the study run from? Ninewells Hospital (UK)

When is the study starting and how long is it expected to run for? February 2004 to October 2009

Who is funding the study? Chest Heart and Stroke Scotland (UK)

Who is the main contact? Dr Jacob George j.george@dundee.ac.uk

Contact information

Type(s)

Scientific

Contact name

Dr Jacob George

ORCID ID

http://orcid.org/0000-0001-8154-8278

Contact details

Department of Clinical Pharmacology Centre for Cardiovascular and Lung Biology Mailbox 2, Level 7 Ninewells Hospital Dundee United Kingdom DD1 9SY +44 (0)1382 632180 j.george@dundee.ac.uk

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers GEO003

Study information

Scientific Title

The potential role of B-Type natriuretic peptide (BNP) and high-sensitivity troponin t(hs-tnt) prescreening by GPs to better target rapid access chest pain clinic referrals: a prospective observational longitudinal cohort study

Study objectives

The aim of this study is to determine how accurate a pre-screening test for BNP and high-sensitivity troponin t(hs-tnt) would be in safely identifying that a potential RACPC referral was unnecessary.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Tayside Local Research Ethics Committee, 04/08/2005, ref: 05/S1402/49

Study design

Prospective observational longitudinal cohort study

Primary study design

Observational

Secondary study design

Cohort study

Study setting(s)

GP practice

Study type(s)

Diagnostic

Participant information sheet

See additional files

Health condition(s) or problem(s) studied

Cardiac chest pain

Interventions

625 patients attending Rapid Access Chest Pain Clinics in Tayside were recruited into the study over a two year period. Each patient had demographic data, a near patient BNP and high-sensitivity troponin t(hs-tnt) level (Biosite®), clinical and laboratory measurements documented. The initial diagnosis was considered positive for ischaemic heart disease (IHD) if the patient had a positive troponin T or was referred for angiography. All patients were then followed up one year after their initial RACPC attendance using hospital medical records, primary care records and telephone calls to document further RACPC attendance, hospitalisations for chest pain and survival data.

Intervention Type

Other

Phase

Not Applicable

Primary outcome measure

Negative predictive value of BNP compared to final clinical diagnosis and troponin T result is measured by a near-patient assay at 12 months.

Secondary outcome measures

Clinical event rate such as further admissions for chest pain, referral for angioplasty or coronary artery bypass grafting and death are assessed continuously for 12 months.

Overall study start date

01/02/2004

Completion date

01/10/2009

Eligibility

Key inclusion criteria

- 1. Patients presenting to the rapid Access Chest Pain Clinic with Cardiac Chest Pain
- 2. Aged 20 93 years, both male and female

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

625

Key exclusion criteria

Patients with non-cardiac chest pain

Date of first enrolment

01/03/2007

Date of final enrolment

01/10/2008

Locations

Countries of recruitment

Scotland

United Kingdom

Study participating centre Ninewells Hospital

Department of Clinical Pharmacology

Dundee United Kingdom DD1 9SY

Sponsor information

Organisation

University of Dundee

Sponsor details

Nethergate Dundee Scotland United Kingdom DD1 4HN

Sponsor type

University/education

Website

http://www.dundee.ac.uk/

ROR

https://ror.org/03h2bxq36

Funder(s)

Funder type

Charity

Funder Name

Chest Heart and Stroke Scotland

Results and Publications

Publication and dissemination plan

Planned publication in a peer reviewed journal.

Intention to publish date

31/12/2012

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Dr Jacob George (j.george@dundee.ac.uk)

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/02/2012		Yes	No
Participant information sheet	version V1	30/12/2009	13/02/2017	No	Yes