A randomised, double-blind, placebo-controlled study of Glypromate® in patients undergoing coronary artery bypass graft surgery

∐ Protocol
Statistical analysis plan
☐ Results
☐ Individual participant data
Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Alan Merry

Contact details

University of Auckland Mercy Hospital 98 Mountain Road Epsom Auckland New Zealand 1031 +64 (0)9 623 5700 a.merry@auckland.ac.nz

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

Neu-GPE-CABG-001

Study information

Scientific Title

Acronym

SNUG (Studying Neurons Using Glypromate®)

Study objectives

Study is designed:

- 1. To determine the pharmacokinetics of Glypromate® in patients undergoing Coronary Artery Bypass Graft (CABG) surgery with/without valve replacement/repair
- 2. To show that Glypromate® use is not associated with major adverse events when compared to placebo in people undergoing CABG surgery

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Health condition(s) or problem(s) studied

Coronary Artery Bypass Graft (CABG) surgery

Interventions

This phase two study will be conducted in two stages:

In stage 1 (conducted at the Principal Investigator's site only), patients will be randomised in a 1: 1 fashion to receive intravenous (IV) Glypromate® 1 mg/kg/hr for four hours or 3 mg/kg/hr for four hours. Two to four patients are expected to be enrolled into this open-label stage of the study.

In stage 2, participants from five centres will be randomised in a 1:1:1 fashion to receive IV Glypromate® 1 mg/kg/hr for four hours or IV Glypromate® 3 mg/kg/hr over four hours or IV Placebo (normal saline) for four hours. The Glypromate®/Placebo infusion will commence at the start of chest closure.

Participants will be observed from randomisation through to discharge or day 14, whichever comes sooner.

Intervention Type

Drug

Phase

Phase II

Drug/device/biological/vaccine name(s)

Glypromate®

Primary outcome measure

To determine the pharmacokinetics of Glypromate® in patients undergoing CABG surgery to assess dose-response relationships.

Secondary outcome measures

To monitor the safety profile of Glypromate® treatment compared to placebo in patients undergoing CABG. Data will be collected through to discharge or day 14 whichever comes first.

Overall study start date

26/09/2005

Completion date

28/02/2006

Eligibility

Key inclusion criteria

Participants must meet all of the following criteria:

- 1. Be at least 60 years of age
- 2. Be scheduled for non-urgent, on-pump CABG surgery
- 3. Be willing to provide written informed consent
- 4. Be agreeable to be undergo all study tests (collection of blood for PK assessment)

The Glypromate®/placebo infusion will be commenced at the start of chest closure providing the following criteria are met:

- 1. The patient has been successfully weaned off the bypass pump
- 2. The patient does not have an Intra-Aortic Balloon Pump (IABP)
- 3. The anaesthetist has assessed the patient as having no contraindications to receiving Glypromate®/placebo medication

Participant type(s)

Patient

Age group

Senior

Sex

Both

Target number of participants

30

Key exclusion criteria

A participant will be ineligible if he/she meets any of the following criteria:

- 1. Body weight less than 55 kg or more than 120 kg
- 2. Scheduled to undergo a significant concomitant surgical procedure (e.g. carotid endarterectomy, aortic root repair or replacement, Deep Hypothermic Circulatory Arrest [DHCA] or pulmonary resection)
- 3. Has a pre or perioperative mechanical assist device or IABP inserted for shock/low output syndrome
- 4. Renal insufficiency (serum creatinine greater than 0.17 mmol/l) or renal failure requiring dialysis
- 5. Chronic hepatic failure and/or cirrhosis
- 6. History of significant haematologic or coagulation disorders, including thrombocytopenia (platelet count less than 50,000), known hypercoagulable state, or recurrent deep vein thrombosis
- 7. Current participation or participation within the seven days prior to the start of this study in another investigational drug or device study
- 8. History of or any current condition that in the investigator's opinion would interfere with study participation or evaluation of results

Date of first enrolment

26/09/2005

Date of final enrolment

28/02/2006

Locations

Countries of recruitment

New Zealand

Study participating centre University of Auckland Auckland New Zealand 1031

Sponsor information

Organisation

Neuren Pharmaceuticals Limited (New Zealand)

Sponsor details

P.O. Box 9923 Newmarket Auckland New Zealand 1031 +64 (0)9 367 7167 ext 89771 mscott@neurenpharma.com

Sponsor type

Industry

Website

http://www.neurenpharma.com

ROR

https://ror.org/0503fq502

Funder(s)

Funder type

Industry

Funder Name

Grant from a New Zealand Government Agency, Foundation for Research Science and Technology.

Funder Name

Grant type: Technology for Business Growth

Funder Name

Grant title: Implementation for Phase II for Glypromate

Funder Name

Study is also internally funded by Neuren Pharmaceuticals Limited (New Zealand)

Results and Publications

Publication and dissemination planNot provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summaryNot provided at time of registration