

Periodontal Intervention for Smoking cessation (PARIS): Combining individual-based smoking cessation counseling, pharmacotherapy, and dental hygiene intervention

Submission date 01/02/2010	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 10/03/2010	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 22/02/2013	Condition category Mental and Behavioural Disorders	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Prof Jacques Cornuz

Contact details
Dept Ambulatory Care and Community Medicine
Lausanne University Hospital
44, Bugnon Avenue
Lausanne
Switzerland
1011

Additional identifiers

Protocol serial number
N/A

Study information

Scientific Title

A pilot study combining individual-based smoking cessation counseling, pharmacotherapy, and dental hygiene intervention

Acronym

PARIS

Study objectives

The present study assessed the feasibility and acceptability of integrating dentists in a medical smoking cessation intervention.

Ethics approval required

Old ethics approval format

Ethics approval(s)

The Lausanne University ethics committee approved on the 2nd of October 2007 (ref: 230/07)

Study design

A non-controlled feasibility pilot study

Primary study design

Interventional

Study type(s)

Other

Health condition(s) or problem(s) studied

Tobacco Use Prevention

Interventions

Smokers willing to quit underwent an 8-week smoking cessation intervention combining individual-based counseling and nicotine replacement therapy and/or bupropion, provided by a general internist. In addition, a dentist provided a dental exam combined with oral hygiene treatment and information about chronic effects of smoking on oral health during two visits, at week 1 and week 2.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Feasibility and acceptability of the dentist intervention, assessed by a hetero-administered questionnaire for the patient at 8 weeks.

Key secondary outcome(s)

1. Smoking cessation rates, assessed by hetero-administered questionnaire and carbon monoxide expiration test at 8 weeks and 6 months.
2. Changes in orodental status, assessed by an oro-dental exam by the dentist at the 2nd visit

Completion date

01/01/2009

Eligibility

Key inclusion criteria

1. Age between 18 and 70
2. Currently smoking for ≥ 3 years at least 10 cig./day
3. Motivation to quit smoking of 6/10 or more on the Likert Scale (from 0 to 10)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

1. Current pharmacological use to quit smoking
2. Presence of an unstable or life-threatening medical condition
3. Current psychiatric illness
4. At risk alcohol consumption
5. Illegal drug consumption, such as cannabis
6. Meets American Heart Association's criteria for antibiotic-prophylaxis before dental intervention (Wilson, et al., 2007)
7. Long-term bisphosphonate treatment
8. Recent oral hygiene intervention (< 6 months)

Date of first enrolment

01/10/2007

Date of final enrolment

01/01/2009

Locations

Countries of recruitment

Switzerland

Study participating centre

Dept Ambulatory Care and Community Medicine
Lausanne
Switzerland
1011

Sponsor information

Organisation

Swiss Tobacco Prevention Fund (Switzerland)

ROR

<https://ror.org/01qtc5416>

Funder(s)

Funder type

Government

Funder Name

Swiss Tobacco Prevention Fund (Switzerland)

Funder Name

Federal Office for Public Health (Office Fédéral de la Santé Publique [OFSP]) (Switzerland) -
Smoking Prevention Fund (Fonds de prévention du tabagisme)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	17/06/2010		Yes	No