

Is ropivacaine (Rop) faster than bupivacaine (Bvp) for extending low dose analgesia for Caesarean section?

Submission date 12/09/2003	Recruitment status Stopped	<input type="checkbox"/> Prospectively registered
Registration date 12/09/2003	Overall study status Stopped	<input type="checkbox"/> Protocol
Last Edited 03/07/2008	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data
		<input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Steve Yentis

Contact details

Magill Department of Anaesthesia, Intensive Care & Pain Management
Chelsea and Westminster Hospital
369 Fulham Road
London
United Kingdom
SW10 9NH
+44 020 8746 8026
s.yentis@ic.ac.uk

Additional identifiers

Protocol serial number

N0060047420

Study information

Scientific Title

Study objectives

Not provided at time of registration

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Not Specified

Health condition(s) or problem(s) studied

Pregnancy and Childbirth: Analgesia

Interventions

Women with labour epidurals are randomised to Rop or Bvp when they require emergency Caesarean section.

Added July 2008: Trial stopped - see reasons/results in Publications field

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

ropivacaine, bupivacaine

Primary outcome(s)

1. Speed of onset of block
2. Extent of block

Key secondary outcome(s)

Not provided at time of registration

Completion date

01/08/2003

Eligibility

Key inclusion criteria

Labouring women with epidurals.

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Female

Key exclusion criteria

Not provided at time of registration

Date of first enrolment

01/02/1999

Date of final enrolment

01/08/2003

Locations**Countries of recruitment**

United Kingdom

England

Study participating centre

Magill Department of Anaesthesia, Intensive Care & Pain Management

London

United Kingdom

SW10 9NH

Sponsor information**Organisation**

Department of Health (UK)

Funder(s)

Funder type

Government

Funder Name

Chelsea and Westminster Healthcare NHS Trust (UK)

Results and Publications

Individual participant data (IPD) sharing plan**IPD sharing plan summary**

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/10/2004		Yes	No