Randomised multicentre study of prosthetic treatment options for shortened dental arch: pilot trial

Submission date	Recruitment status No longer recruiting	Prospectively registered	
07/03/2008		Protocol	
Registration date	Overall study status	Statistical analysis plan	
21/04/2008	Completed	[X] Results	
Last Edited	Condition category	Individual participant data	
20/04/2010	Oral Health		

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Prof Michael Walter

Contact details

Fetscherstr. 74 Dresden Germany 01307 +49 351/458 2706 Michael.Walter@uniklinikum-dresden.de

Additional identifiers

Protocol serial number

DFG WA 831/2-1 to 2-6

Study information

Scientific Title

Acronym

SDAS - pilot

Study objectives

Over the last 20 years a mechanistic attitude correlated with a lack of longitudinal controlled randomised trials regarding the question of prosthetic treatment after tooth loss. The need assessment considered the replacement of all missing teeth by fixed or removable partial dentures or dental implants as a necessity especially in cases of shortened dental arches. Modern prosthetic concepts distinguish between different dimensions of need (normative need, perceived need) being well aware of the fact that perceived need has been under-represented in the past. An innovative sight puts a higher emphasis to the subjective components of need assessment and outcome measurement. Generally three adverse effects of non-replacement of molars were postulated: temporomandibular joint (TMJ) disorders, tooth migration /overeruption, insufficient chewing ability. However, no evidence based on randomised trials has been provided concerning the incidence of the adverse side effects mentioned above, nor is there high-level evidence regarding a benefit of removable dentures for molar replacement. On the contrary removable partial dentures are compromised by a high incidence of adverse side effects such as plaque accumulation and peridontal breakdown.

Among therapeutic alternatives, an approach with a limited restoration goal focused on incisors, canines and premolars (shortened dental arch [SDA] concept) has been described and implemented although discussed controversially. Within this concept, fixed partial dentures are used for tooth replacement of which a superior performance compared with removable partial dentures has been reported. The multicentre study was initiated in 2000 because evidence was lacking concerning the benefit of different therapeutic options regarding the preservation of oral health, oral health related quality of life, patients satisfaction, absence of discomfort, satisfactory chewing ability and aesthetic satisfaction. Public health aspects of the study lie in the fields of health economics, avoidance of over-treatment, and therapy guidelines on a population based level.

Two prosthetic therapy arms will be compared:

- 1. The replacement of posterior teeth at last up to the first molar by removable partial dentures

 2. Prosthetic treatment according to the shortened dental arch concept. To avoid removable
- 2. Prosthetic treatment according to the shortened dental arch concept. To avoid removable partial dentures, posterior teeth are replaced up to the second premolar by fixed restorations, if necessary. Molars are not replaced.

The aim of this trial is to test the hypothesis that the treatment outcome varies depending on the treatment concept (fixed versus removable prostheses) in the therapy of patients with missing molars.

Please note that the pilot phase of this trial has been completed. Details of the full trial may be found at http://www.controlled-trials.com/ISRCTN97265367

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics Committee of the Medical Faculty of the Technical University of Dresden (Ethikkommission der Medizinischen Fakultät der Technischen Universität Dresden). Date of approval 19/04/1999 (ref: EK 260399)

Study design

Multi-centre randomised controlled clinical trial.

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Tooth loss/ molar replacement

Interventions

Control group (Therapy A): The molar replacement by removable partial dentures, carried out using fixed crowns and bridges as anchor for removable dentures

Intervention group (Therapy B): Restorations according to the SDA concept, with only fixed restorations or no restoration at all. The maximum extension reached up to the second premolar, and no molars were replaced. All restorations were made according to a standardized procedure (SOP) given by the study protocol.

Standard gold alloys and dental ceramics were used for fixed restorations, base metal alloys for the removable denture frameworks.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

Further tooth loss. Duration of follow-up: 5 years

Key secondary outcome(s))

The following were assessed at baseline (4-8 weeks after insertion), 6 month, then annually from Year 1 to 5:

- 1. Clinical:
- 1.1. Crown/root caries
- 1.2. Abrasion (Index 0-4)
- 1.3. Interdental spacing in the anterior region (Index 0-3)
- 1.4. Sensibility (+/-)
- 1.5. Periodontitis/Gingivitis: Plaque-index (index 0-3), probing depth (6 point measurement in mm), attachment loss (6 point measurement in mm), gingival index(index according to Sillnes and Loe: 0-3), bleeding on probing (BOP)(+/-), tooth mobility (index 0-3), mucosa lesions (California Dental Association [CDA] Criteria)
- 2. Clinical dysfunction index (Helkimoindex): Muscle pain via palpation (m. masseter pars profunda et superficialis, m. temporalis pars posterior et anterior, m. suboccipitalis, m. sternocleidomastoideus, m. pterygoideus medialis et lateralis, t. temporalis)
- 2.1. Range of movement (mm): maximal opening, maximal lateral movements, maximal protrusion
- 2.2. TMJ function: Description of pain on movement/path of movement, palpation/auscultation

- 3. Technical (according to the CDA criteria): Treatment performance, preparation form, marginal fit, occlusion static/dynamic im µm, proximal contacts (shape/ strength)
- 4. Technical performance (according to the CDA criteria): Evaluation of used materials, prosthesis and bridge design, saddle extension, attachment performance, possibility of dental hygiene
- 5. Aesthetics (according to the CDA criteria): rated by the dentist, surface, color, translucency, contour
- 6. Ridge reduction: (x-ray examination according to Steen)

Subjective:

- 7. Oral health related quality of life (OHIP-Questionnaire): Measure of self reported dysfunction, discomfort and disability attributed to oral conditions
- 8. Dworkin Index Axis II (questionnaire): Assessment of psychological distress and psychosocial dysfunction including questions regarding:
- 8.1. Graded chronic pain severity
- 8.2. Depression
- 8.3. Vegetative symptoms and somatization subscales of the SCL-90-R developed by Derogatis and others
- 8.4. Jaw disability checklist

The CDA Criteria are used according to the Guidelines for the Assessment of Clinical Quality and Professional Performance of the California Dental Association: http://www.cda.org/library/cda_member/policy/quality/quality.html

Completion date

26/11/2007

Eligibility

Key inclusion criteria

- 1. Patients over 35 years of age, male and female
- 2. Those who requested prosthetic treatment with a minimum dentition of both canines and one premolar per side preserved in at least one jaw (Kennedy class I). A dentition including all anterior teeth up to the second premolar on both sides in one jaw was defined as maximum
- 3. Rejection of implant treatment by the patient
- 4. Patients with general health according to American Society of Anesthesiologists (ASA) classification group one ore two
- 5. All abutment teeth must be free of periodontal disease (pocket depth less or equal 4 mm, tooth mobility <= grade 2, mean plaque index <= grade 2, bleeding on probing at all teeth <=25 %) and caries
- 6. Caries free adjacent teeth
- 7. Sufficient treatment of the opposite jaw, extending the dentition depending on the randomized treatment option up to the second premolar or the first molar

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Sex

All

Key exclusion criteria

- 1. Patients with alcohol or drug addiction
- 2. Mentally disordered patients
- 3. Patients with TMJ disorders
- 4. Dysgnathic patients with Angle class II or III
- 5. Patients who have received or need orthodontical treatment
- 6. Patients who have been already sufficiently treated
- 7. Patients who do not accept a removable deture
- 8. Patients who demand the replacement of all molars
- 9. Patients with general health American Society of Anesthesiologists (ASA) classification group four

Date of first enrolment

01/10/2000

Date of final enrolment

26/11/2007

Locations

Countries of recruitment

Germany

Study participating centre

Fetscherstr. 74

Dresden Germany 01307

Sponsor information

Organisation

German Research Foundation (Deutsche Forschungsgemeinschaft) (Germany)

ROR

https://ror.org/018mejw64

Funder(s)

Funder type

Government

Funder Name

German Research Foundation (Deutsche Forschungsgemeinschaft) (Grant ref: WA 831/2-1 to 2-6) (Germany)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created Date added	Peer reviewed?	Patient-facing?
Results article	results	01/11/2005	Yes	No
Participant information sheet	Participant information sheet	11/11/2025 11/11/2025	No	Yes