

APPEL 3: Algorithm to control Postprandial, Post exercise and night glucose Excursions in a portable closed Loop format

Submission date 18/12/2012	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
Registration date 29/01/2013	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
Last Edited 21/01/2019	Condition category Nutritional, Metabolic, Endocrine	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Over the last decade insulin delivery and blood glucose monitoring have evolved facilitating better glucose control. Continuous Subcutaneous Insulin Infusion (CSII) and Continuous Glucose Monitoring (CGM) sensors combined with an insulin delivery algorithm results in a closed loop system: an artificial pancreas.

The aim of this study is to test the efficacy of a portable closed loop system first in the clinical research unit directly followed by testing at home.

Who can participate?

All adults diabetes mellitus (DM) type 1 treated with insulin pump

What does the study involve?

All participants will wear the artificial pancreas that will be compared to regular insulin pump use.

What are the possible benefits and risks of participating?

Benefits: This study will provide insight in glucose metabolism

Risk: hyperglycemia (high blood sugar) or hypoglycemia (low blood sugar)

Where is the study run from?

Academic Medical Center Amsterdam

When is the study starting and how long is it expected to run for?

Study started in January 2012 and will run till March 2013

Who is funding the study?

Academic Medical Center Amsterdam

Who is the main contact?

AC van Bon

acvanbon@rijnstate.nl

Contact information

Type(s)

Scientific

Contact name

Dr Hans DeVries

Contact details

Academic Medical Center (AMC)

Meibergdreef 9

Amsterdam

Netherlands

1100 DD

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

NL37969.018.11; U1111-1125-4086

Study information

Scientific Title

APPEL 3: Algorithm to control Postprandial, Post exercise and night glucose Excursions in a portable closed Loop format

Acronym

APPEL3

Study objectives

The aim of this study is to test the efficacy of a portable bi hormonal closed loop system first in the clinical research unit directly followed by testing at home compared to open loop.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethical Committee of the Academic Medical Center Amsterdam, 18 October 2011, ref number 2011_292

Study design

Non randomized controlled trial

Primary study design

Interventional

Secondary study design

Non randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information

Health condition(s) or problem(s) studied

Diabetes mellitus type 1 treated with insulin pump

Interventions

All participants have type 1 diabetes and are treated with an insulin pump. All participants start with open loop: the participants will wear CGM for 60 hours. This is followed after several weeks with the closed loop: the participants will wear the artificial pancreas.

Open loop arm: glucose control in daily life and measured via continuous glucose monitoring (CGM) for 60 hours

Closed loop arm: glucose control by artificial pancreas for 60 hours, the first 12 hours are for tuning and the last 48 hours participants are sent home. The closed loop contains of two CGM, two subcutaneous pumps for insulin and glucagon administration and portable computer with the algorithm.

The closed loop contains the algorithm, insulin pump, glucagon pump and CGM. These are similar to the software used in APPEL 2 (MEC09/263) [registered at CCMO <http://www.ccmo-online.nl>, ref: NL29814.018.09. The hardware of the closed loop system is changed compared to APPEL 2. The personal computer is changed in a small portable computer that contains two systems: a controlling system and an operating system. The interface is a wireless connection. Due to this connection, the glucose concentrations, insulin administration, glucagon administration and heart rate are read out continuously and visually during the whole experiment.

Insulin pump

The D-Tron+ (Disetronic Medical Systems, St. Paul, MN) is a CE marked infusion system. A plastic reservoir filled with short-acting insulin is connected to a catheter that is inserted into the subcutaneous tissue. The insulin catheter needs to be changed every three days. The device is driven by the algorithm to deliver insulin. Insulin will be administered if the glucose rises above 6.5 mmol/l. Depending of the rise in glucose concentration, an insulin bolus will be given.

Glucagon pump

The D-Tron+ (Disetronic Medical Systems, St. Paul, MN) is a CE marked infusion system. A plastic reservoir filled with glucagon is connected to a catheter that is inserted into the subcutaneous tissue. The device is driven by the algorithm to deliver glucagon. Glucagon will be administered when glucose falls below 6.5 mmol/l to regulate the glucose levels between the normal range. First a bolus glucagon is injected depending of the decrease in glucose concentration followed by glucagon delivery according to the formula. Below glucose concentration of 4.5 mmol/l, glucagon is administered in a rescue bolus.

Glucose sensor

The Continuous Glucose Monitoring System Gold (Medtronic Minimed, Sylmar, CA) is a needle-type sensor that quantifies interstitial glucose concentrations via the glucose oxidase reaction with the enzyme immobilized on a subcutaneously inserted electrode. Due to connection to the computer glucose values can be monitored in real time.

During closed loop two CGM sensors will be used: one primary and one back up in case of failure of the first sensor.

Algorithm

The algorithm is designed and patented by Inreda Diabetic B.V., Goor, The Netherlands. Both study devices will be connected via a personal computer embedded in a control unit (Robopump®, Inreda Diabetic B.V., Goor, The Netherlands), on which the algorithm is programmed. The most important assumption is that the algorithm makes use of the current blood glucose values of the patient. In the control unit blood glucose values are programmed between an upper and lower limit. These programmed values are continuously compared to the current, i.e. measured values which the glucose sensor transmits per time interval for programming.

The two D-Tron+pumps are controlled by the control unit. One pump contains a reservoir of short acting insulin analogue (NovoRapid, Novo Nordisk) and the other pump contains glucagon (Glucagen, Novo Nordisk).

The control unit has two operating ranges:

At glucose values below the programmed lower limit a sound signal will be generated. The patient is hereby warned that the lower limit is being approached or has been exceeded. He/she will be advised to eat and thereby correct the low glucose level. In the case of a further decrease of glucose levels, glucagon will be injected in accordance with the set values and a curve to be programmed according to: $\text{ml(E) / fall mmol/l / time unit}$

At blood glucose values above the programmed upper limit insulin will be injected, as according to the set values and a curve to be programmed according to: $\text{ml(E) / rise mmol/l / unit of time}$ (see figure 2), and a (different) sound signal is generated, whereby the patient is alerted that the upper limit is being approached or has been exceeded. If the fall in the blood glucose values starts at a determined speed (preferably adjustable) the supply of insulin is stopped.

The control unit contains a program that incorporates an adjustable curve corresponding to the amount of insulin or glucagon respectively introduced into the body of the patient and the nominal concentration of the glucose in the blood of the patient as a reaction thereto. The control unit stores in memory information relating to the introduced quantity of insulin or glucagon respectively, and in each successive case compare this information to the concentration of the glucose in the blood of the patient as a reaction thereto. Furthermore, this latter information is stored and used for metering the insulin or glucagon respectively. The algorithm is therefore a learning type, adapting to the specific patient.

Interface

The algorithm is incorporated in a portable computer. The voltage is low power and is provided by a rechargeable battery pack. Due to wireless internet connection, the glucose values, insulin administration, glucagon administration and heart rate can be read out immediately.

Intervention Type

Other

Phase

Not Applicable

Primary outcome measure

To assess the efficacy of the closed loop system, defined as the time spent in euglycemia. Euglycemia is defined as glucose concentration between 3.5 mmol/l -10 mmol/l. The proportions of time spent in eu-, hypo and hyperglycaemia in closed loop will be compared to open loop.

Time points of measurement are the last 48 hours divided in first 24 hours and last 24 hours. Furthermore to analyze for a meal and three hours postprandial
Night: from 00.00 till 7.00 AM

Secondary outcome measures

1. Difference in overall mean sensor glucose comparing closed loop system to open loop
2. Difference in postprandial, post exercise and night sensor glucose excursions measured as area under the curve comparing closed loop to open loop
3. Difference in numbers of hypoglycaemias (glucose at or below 3.5 mmol/l) comparing the closed loop to open loop

Time points of measurement are the last 48 hours divided in first 24 hours and last 24 hours. Furthermore to analyze for a meal and three hours postprandial
Night: from 00.00 till 7.00 AM

Overall study start date

01/01/2012

Completion date

01/03/2013

Eligibility

Key inclusion criteria

1. Diabetes mellitus type 1 treated with continuous subcutaneous insulin infusion (CSII) for a minimum of 6 months
2. Age: 18-75, either sex
3. Willing and able to sign informed consent

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Upper age limit

75 Years

Sex

Both

Target number of participants

16

Key exclusion criteria

1. Impaired awareness of hypoglycaemia according to Gold [7] and / or Clarke [8] method
2. Body mass index (BMI) > 35 kg/m²
3. Glycated haemoglobin (HbA1c) > 11.0%
4. Use of heparin, coumarin derivatives or oral corticosteroids
5. Skin condition prohibiting needle insertion
6. Pregnancy and/or breastfeeding
7. Any condition that the local investigator feels would interfere with trial participation or the evaluation of results

Date of first enrolment

01/01/2012

Date of final enrolment

01/03/2013

Locations**Countries of recruitment**

Netherlands

Study participating centre

Academic Medical Center (AMC)

Amsterdam

Netherlands

1100 DD

Sponsor information**Organisation**

Academic Medical Center (AMC) (Netherlands)

Sponsor details

Meibergdreef 9
Amsterdam
Netherlands
1100 DD
-
acvanbon@rijnstate.nl

Sponsor type

Hospital/treatment centre

Website

<http://www.amc.nl>

ROR

<https://ror.org/03t4gr691>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

Academic Medical Center Amsterdam (Netherlands)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date**Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/03/2014	21/01/2019	Yes	No