

A group treatment for antenatal anxiety

Submission date 21/09/2023	Recruitment status Recruiting	<input checked="" type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 12/02/2024	Overall study status Ongoing	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 16/07/2025	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Individual participant data <input checked="" type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Up to a quarter of women struggle with serious anxiety symptoms during pregnancy. Many women will not improve without treatment, with symptoms continuing postnatally and for some, leading to depression. High anxiety during pregnancy is related to negative outcomes in childbirth and baby development. Babies of mothers who struggle with anxiety during pregnancy can have more problems with their emotions, behaviours and cognitive development. If the mother's anxiety continues after the baby is born it can make it difficult for the mother to bond with the baby and provide the kind of care the baby needs. Together, the impact of untreated perinatal mental health problems cost over £8 billion each year in England. Despite the fact that anxiety problems are more common during pregnancy and have many negative effects, there is a lack of high quality research about the best way to help mothers. In the NHS most mothers with antenatal anxiety (65%) are seen in primary mental health care (IAPT), but IAPT lacks treatments that specifically address antenatal anxiety.

We will test whether CALM 1) improves anxiety in pregnant women during pregnancy; 2) has sustained benefits in the postnatal period, up to a year after the baby is born; 3) improves the relationship the parent has with the child and how the child develops and 4) is cost-effective for the NHS to deliver. We will also interview parents who completed CALM to see which parts helped them the most, and we will ask care providers about the best ways to ensure CALM fits within the NHS.

Who can participate?

Any woman or birthing person over the age of 18 who scores 8 and above on the GAD-7.

What does the study involve?

Eligible participants will be randomised to either receive the intervention and treatment as usual or solely treatment as usual. Those in the intervention will receive a 4-session group based online intervention delivered by healthcare and psychological practitioners. The intervention includes partners or a close support and the group encourages peer support. All participants will be asked to complete questionnaires at 6 timepoints (at the start, 22 and 32 weeks pregnant, 2/6/12 months post birth).

What are the possible benefits and risks of participating?

There is a lack of information about the effectiveness and long-term benefits of a group

treatment for antenatal anxiety compared to usual care. Regardless of the group you will be randomised into, your input in the clinical trial will contribute to the evidence around what treatment is effective in targeting pregnancy-related anxiety. If you do decide to take part, as a thank you gesture we will offer you a voucher.

We do not expect there are any disadvantages or risks to you. All the sessions will be arranged at a time to suit you and your partner. You may feel anxious before or tired after taking part in the sessions or while completing the questionnaires, but we will do everything we can to minimise or prevent this.

Where is the study run from?

The Universities of Exeter, Manchester, Cambridge, and Birmingham are leading the recruitment. This trial also has academics from Imperial College London, University of Bath, Anna Freud Centre and Kings College London. It is sponsored by Devon Partnership NHS Trust.

When is the study starting and how long is it expected to run for?

September 2023 to August 2027

Who is funding the study?

National Institute of Health Research - Health Technology Assessment Programme (UK)

Who is the main contact?

Dr Antoinette Davey, antoinette.davey@exeter.ac.uk

Contact information

Type(s)

Public

Contact name

Dr Antoinette Davey

ORCID ID

<https://orcid.org/0000-0002-5118-6912>

Contact details

Devon Partnership Trust
Wonford House
Dryden Road
Exeter
United Kingdom
EX2 5AF
+44 (0)7828843266
antoinette.davey@exeter.ac.uk

Type(s)

Principal Investigator

Contact name

Prof Heather O'Mahen

ORCID ID

<https://orcid.org/0000-0003-3458-430X>

Contact details

University of Exeter
Faculty of Health and Life Sciences
Psychology Department
Washington Singer Labs
Perry Road
Exeter
United Kingdom
EX44QG
+44 (0)1392724651
H.OMahen@exeter.ac.uk

Additional identifiers**EudraCT/CTIS number**

Nil known

IRAS number

333463

ClinicalTrials.gov number

Nil known

Secondary identifying numbers

29369, IRAS 333463, CPMS 60522

Study information**Scientific Title**

ACORN-II: A multi-site randomised COntrolled trial to evaluate the impact of a group tReatment for aNtenatal anxiety

Acronym

ACORN-II

Study objectives

The primary objective is to test whether CALM + Treatment as usual (TAU) reduces anxiety in pregnant women relative to TAU alone. We will also assess whether CALM + TAU has sustained benefits up to a year after the baby is born compared to TAU alone.

Ethics approval required

Ethics approval required

Ethics approval(s)

Approved 24/01/2024, South West - Cornwall & Plymouth Research Ethics Committee (2 Redman Place, Stratford, London, E20 1JQ, United Kingdom; +44 (0)207 104 8143; cornwallandplymouth.rec@hra.nhs.uk), ref: 24/SW/0013

Study design

Group sequential multi-centre parallel arm individually randomized controlled trial with internal pilot interim monitoring for futility and economic and process evaluation

Primary study design

Interventional

Secondary study design

Randomised parallel trial

Study setting(s)

Community, Hospital

Study type(s)

Treatment

Participant information sheet

See study outputs table

Health condition(s) or problem(s) studied

Antenatal anxiety

Interventions

Potential participants will be screened for anxiety at first-trimester scanning clinics. If individuals score 8 and above on the GAD-7, they will be eligible for the trial.

Eligible participants who consent will be randomly allocated on a 1:1 basis to CALM + TAU or TAU only using permuted block randomisation which will be stratified by site and anxiety diagnosis pre-pregnancy versus de novo antenatal anxiety. Randomisation will be delivered by a web-based randomisation system (integrated within the trial REDCap system).

The CALM intervention comprises four 2-hour online group sessions delivered to both the pregnant woman/birthing person and their partner and one 2-hour postnatal reunion. It is delivered by a healthcare and psychological practitioner. The intervention focuses on strategies that: (1) reduce avoidance associated with worry, (2) improve the ability to tolerate uncertainty that contributes to anxious worry in the perinatal period and (3) improve relationship satisfaction through joint communication skills about how to manage worry together.

Intervention Type

Behavioural

Primary outcome measure

Anxiety will be measured using the Generalised Anxiety Disorder -7 PROM at baseline, 22 & 32 weeks pregnant, and postnatally at 3, 6, and 12 months.

Secondary outcome measures

1. Antenatal anxiety measured using Pregnancy-Related Anxiety Questionnaire – Revised 2 (PRAQ-R2) at baseline, 22 weeks and 32 weeks
2. Depression using Patient Health Questionnaire-9 (PHQ-9) at baseline, 22 & 32 weeks pregnant, and postnatally at 3, 6, and 12 months.

3. Relationship adjustment measured using Revised Dyadic Adjustment Scale (DAS) at baseline, 22 and 32 weeks pregnant.
4. Health related quality of life measured using EuroQol-5 Dimensions-5 Level (EQ-5D-5L) at baseline, 22 & 32 weeks pregnant, and postnatally at 3, 6, and 12 months.
5. Service use measured using Adult Service Use Schedule (AD-SUS) at baseline, 32 weeks pregnant and 12 months postnatal.
6. Infant behaviour and temperament measured using Infant Behavior Questionnaire – Revised (IBQ-R) at 6 months postnatal.
7. Perceptions of childbirth measured using Maternal Perceptions of Support and Control in Birth (SCIB) at 3 months postnatal.
8. Parent and infant interaction measured using 10-minute videoed interaction and coded using the well-validated NICHD parent-infant sensitivity, intrusiveness scales at 12 months postnatal.
9. Social, emotional and behavioural problems in the infant measured using Brief Infant Toddler Social and Emotional Assessment (BITSEA) at 12 months postnatal.
10. Child development measured using Ages and Stages 3rd edition at 12 months postnatal.

Overall study start date

01/09/2023

Completion date

31/08/2027

Eligibility

Key inclusion criteria

Pregnant women and birthing persons aged 18+ years who score above clinical threshold (8 or more) (on a standard anxiety screening measure (GAD-7))

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Sex

Female

Target number of participants

484

Key exclusion criteria

1. Actively suicidal
2. Substance abuse
3. Psychosis
4. Have childbirth-related PTSD (i.e., individuals eligible to receive PTSD treatment in maternal mental health services)
5. Receiving current psychological treatment for anxiety at baseline interview

Date of first enrolment

01/05/2024

Date of final enrolment

31/12/2025

Locations

Countries of recruitment

England

United Kingdom

Study participating centre**Royal Devon University Healthcare NHS Foundation Trust**

Royal Devon University NHS Ft

Barrack Road

Exeter

United Kingdom

EX2 5DW

Study participating centre**St Mary's Hospital**

Hathersage Rd

Manchester

United Kingdom

M13 0JH

Study participating centre**Royal London Hospital and Associated Community Services NHS Trust**

The Royal London Hospital

Whitechapel

London

United Kingdom

E1 1BB

Study participating centre**Homerton Hospital**

Homerton Row

London

United Kingdom

E9 6SR

Study participating centre
Birmingham Women's NHS Foundation Trust
Birmingham Womens Hospital
Metchley Park Road
Birmingham
United Kingdom
B15 2TG

Sponsor information

Organisation

Devon Partnership NHS Trust

Sponsor details

Wonford House
Dryden Road
Exeter
England
United Kingdom
EX25AF
+44 (0)1392208866
a.wilkie@nhs.net

Sponsor type

Hospital/treatment centre

Website

<https://www.dpt.nhs.uk/>

ROR

<https://ror.org/04fkxrb51>

Funder(s)

Funder type

Government

Funder Name

Health Technology Assessment Programme

Alternative Name(s)

NIHR Health Technology Assessment Programme, HTA

Funding Body Type
Government organisation

Funding Body Subtype
National government

Location
United Kingdom

Results and Publications

Publication and dissemination plan

The substantial output from the research will be evidence of whether or not the CALM intervention is a clinically and cost-effective intervention for pregnant women with anxiety that can be delivered jointly by maternity and mental health providers.

The outputs from the project will include:

- Open access academic papers in high-impact journals
- New mental health intervention: updated 5-session CALM clinician and therapist manuals
- CALM intervention Implementation guidance for maternity and perinatal mental health services, including implementation top tips.
- Therapist training guidance (for perinatal training programmes) with pre-post assessment tools
- Therapist CALM competence assessment
- Policy documents for: NHSE perinatal and maternity teams and NICE

Intention to publish date
31/08/2027

Individual participant data (IPD) sharing plan

The data-sharing plans for the current study are unknown and will be made available at a later date.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Participant information sheet	version 1	04/12/2023	29/01/2024	No	Yes
Participant information sheet	version 1.0	07/11/2023	29/01/2024	No	Yes