

# GREAT-2: a trial of gremubamab compared to placebo in participants with bronchiectasis and chronic *Pseudomonas aeruginosa* infection

<b>Submission date</b> 15/11/2022	<b>Recruitment status</b> No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
<b>Registration date</b> 26/05/2023	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 04/07/2025	<b>Condition category</b> Respiratory	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Patients with bronchiectasis often get chest infections which are difficult to treat causing coughing, sputum (phlegm) production, breathlessness and tiredness. Approximately one third of people with bronchiectasis become infected with bacteria called *Pseudomonas aeruginosa* (P. aeruginosa). P. aeruginosa can often become resistant to antibiotics. The purpose of this trial is to test whether an intravenous infusion (drip) containing a new drug called gremubamab can reduce the amount of infection with P. aeruginosa.

The purpose of this trial is to test whether a new drug called gremubamab, given intravenously, can reduce the amount of infection with P. aeruginosa in people with bronchiectasis. Whether gremubamab can help reduce the number of bronchiectasis exacerbations and improves quality of life will also be examined. The safety of gremubamab use in people with bronchiectasis will be assessed.

Gremubamab is a type of drug called a monoclonal antibody which is expected to work with the immune system to eliminate the P. aeruginosa infection.

Gremubamab is a new medication which is being developed by AstraZeneca. It has been used in a few trials already, in healthy people (Phase I trial) and people who were on a ventilator in intensive care and developed pneumonia (Phase II trial). Phase I trials look at the safety of new drugs and phase II trials look at how effective new drugs are as well as their safety. This trial is a phase II trial which will look at the safety and effectiveness of gremubamab in people with bronchiectasis.

### Who can participate?

People aged 18 – 85 years with bronchiectasis in the UK and Spain

### What does the study involve?

The health of participants treated with gremubamab will be compared with the health of participants given a placebo. The effects of two different doses will also be compared.

The participant will be in the trial for 6 months and will receive infusions of the gremubamab /placebo at monthly intervals for the first 3 months. The trial is expected to run for a total of 18 months.

What are the possible benefits and risks of participating?

**Benefits:**

Participants will be monitored closely during the trial by the trial team. The tests will give the trial team information about the function of participants kidneys, liver, fitness and general wellbeing. If any of these investigations reveal any new clinically significant abnormality, the trial team will tell participants and either discuss this with their GP (with your consent) or refer them to a specialist clinic at the hospital (whichever seems most appropriate.) The trial may not immediately benefit participants, but if the results of the trial are positive this may improve how people with bronchiectasis are treated.

**Risks:**

Previous trials have shown that there was a low risk of allergic reactions to the gremubamab infusion. There is an extremely small risk of severe allergic reaction. The risk of having an allergic reaction will be reduced by giving the infusion slowly and giving an antihistamine before the trial drug administration starts. Participants will be monitored during all infusions of trial medication. A participant's trial medication will be stopped immediately a participant develops signs of a severe allergic reaction.

A few people in the previous trials also reported headache, indigestion and itch.

If a participant develops any reaction to the infusion the doctor looking after the participant will assess it and discuss with the participant if any treatment is required. The doctor will also decide if it is suitable for the participant to continue with their infusions.

Where is the study run from?

University of Dundee (UK)

When is the study starting and how long is it expected to run for?

November 2022 to October 2024

Who is funding the study?

European Respiratory Society (UK)

Who is the main contact?

Dr James Chalmers, j.chalmers@dundee.ac.uk

## **Contact information**

**Type(s)**

Public

**Contact name**

Dr Gillian Martin

**Contact details**

Tayside Clinical Trials Unit

TASC

Level 3, Ninewells Hospital and Medical School

Dundee

United Kingdom

DD1 9SY

+44 (0)1382 381955

GREAT-2-TM@dundee.ac.uk

**Type(s)**

Principal investigator

**Contact name**

Dr James Chalmers

**ORCID ID**

<https://orcid.org/0000-0001-5514-7868>

**Contact details**

Ninewells Hospital and Medical School

Dundee

United Kingdom

DD1 9SY

+44 1382 386131

[j.chalmers@dundee.ac.uk](mailto:j.chalmers@dundee.ac.uk)

## Additional identifiers

**Clinical Trials Information System (CTIS)**

2022-003215-28

**Integrated Research Application System (IRAS)**

1005993

**ClinicalTrials.gov (NCT)**

Nil known

**Protocol serial number**

1-023-22, IRAS 1005993, CPMS 55567

## Study information

**Scientific Title**

GRemubamab ErAdication Trial (GREAT-2) A phase 2 trial of gremubamab compared to placebo in participants with bronchiectasis and chronic Pseudomonas aeruginosa infection

**Acronym**

GREAT-2

**Study objectives**

Primary objective:

To evaluate the efficacy of gremubamab on P. aeruginosa bacterial burden in sputum at week 12

Secondary objectives:

1. To evaluate the efficacy of gremubamab on P. aeruginosa bacterial burden in sputum
2. To determine the persistent effects of gremubamab on P. aeruginosa bacterial burden following discontinuation of treatment (week 24)
3. To determine if gremubamab can achieve eradication of P. aeruginosa in some individuals
4. To determine the effect of gremubamab on health related quality of life

5. To determine the effect of gremubamab on time to first exacerbation
6. To determine the effect of gremubamab on pulmonary function
7. To assess the safety of gremubamab in patients with bronchiectasis
8. To evaluate the PK of gremubamab

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Approved 11/05/2023, East of Scotland Research Ethics Service (EoSRES, Tayside Medical Science Centre, Residency Block, Level 3, George Pirie Way, Dundee, DD1 9SY, UK; +44 (0)1382 383871; tay.eosres@nhs.scot), ref: 22/ES/0051

**Study design**

Interventional double-blind randomized placebo-controlled trial

**Primary study design**

Interventional

**Study type(s)**

Treatment

**Health condition(s) or problem(s) studied**

Bronchiectasis

**Interventions**

Participants will be randomised using a GCP-compliant Interactive Web-based Randomisation System to one of three treatment arms. Randomisation will be stratified by inhaled antibiotic use. Depending on randomisation participants will receive either gremubamab 1500 mg per dose, gremubamab 500 mg per dose or placebo. Participants will receive trial treatment via intravenous infusion every 4 weeks a total of three times. Participants will be assessed during the treatment period (3 months) and for a 3-month period following completion of trial treatment.

**Intervention Type**

Drug

**Phase**

Phase II

**Drug/device/biological/vaccine name(s)**

Gremubamab

**Primary outcome(s)**

Efficacy of gremubamab on *P. aeruginosa* bacterial burden in sputum measured by change from baseline (day 1 - day 84) in quantitative sputum cultures (colony-forming unit (CFU))

**Key secondary outcome(s)**

1. Efficacy of gremubamab on *P. aeruginosa* bacterial burden in sputum measured by change from baseline to Days 7, 14, 28 and 56 in Quantitative sputum cultures
2. Persistent effects of gremubamab on *P. aeruginosa* bacterial burden following

discontinuation of treatment measured by change from baseline to Day 168 in quantitative sputum cultures

3. Eradication defined by negative sputum cultures for *P. aeruginosa* at the end of treatment (Days 84 and 168)
4. Eradication of *P. aeruginosa* measured by change from baseline to Days 28, 56, 84 and 168 in QOL-B, BIM questionnaire
5. Effect of gremubamab on health-related quality of life
  - 5.1. Measured by change from baseline to Days 84 and 168 in St. George's Respiratory Questionnaire
  - 5.2. Measured by change from baseline to Days 28, 56, 84 and 168 in change from baseline in Quality of Life Bronchiectasis questionnaire
  - 5.3. Measured by change from baseline to Days 28, 56, 84 and 168 in change from baseline in Bronchiectasis Impact Measure questionnaire
6. Effect of gremubamab on time to first exacerbation measured by occurrence of exacerbations (as per EMBARC definition of exacerbation). First event from visit 1 to day 84.
7. Effect of gremubamab on pulmonary function measured by change from baseline to Day 28, 56 and 84 in forced expiratory volume in 1 second (FEV1)
8. Safety of gremubamab in patients with bronchiectasis measured by frequency of adverse events and serious adverse events between groups over 168 days
9. Safety of gremubamab in patients with bronchiectasis measured by safety lab parameters between groups over 168 days
10. Pharmacokinetics of gremubamab measured by gremubumab PK parameters through 168 days post-dose

**Completion date**

31/10/2024

## Eligibility

**Key inclusion criteria**

1. Age  $18 \leq 86$  years
2. Clinical diagnosis of bronchiectasis.
3. Able to and provided informed consent.
4. Previous CT scan of the chest demonstrating bronchiectasis in 1 or more lobes
5. *P. aeruginosa* in sputum, bronchoalveolar lavage or another airway sample at least once in the 24 months prior to screening.
6. A sputum sample that is culture positive for *P. aeruginosa* sent at the screening visit and within 35 days of randomization.

**Participant type(s)**

Patient

**Healthy volunteers allowed**

No

**Age group**

Adult

**Lower age limit**

18 years

**Upper age limit**

86 years

**Sex**

All

**Total final enrolment**

62

**Key exclusion criteria**

1. Known hypersensitivity to gremubamab or any excipient of the investigational product
2. Known clinical diagnosis of cystic fibrosis
3. Immunodeficiency requiring replacement immunoglobulin.
4. Active tuberculosis or nontuberculous mycobacterial infection (currently under treatment, or requiring treatment in the opinion of the investigator).
5. Active allergic bronchopulmonary aspergillosis (receiving treatment with corticosteroids and/or antifungal medication).
6. Recent significant haemoptysis (a volume requiring clinical intervention, within the previous 4 weeks prior to screening).
7. Treatment with long-term inhaled, systemic or nebulized anti-pseudomonal antibiotics which are newly initiated within the previous 3 months prior to screening.
8. Chronic treatment with cyclical doses of inhaled or nebulized antibiotics e.g. 28 days on and 28 days off at the time of screening.
9. Receipt of antipseudomonal antibiotics for an exacerbation during the screening period.
10. Treatment with immunosuppressives within previous 6 months prior to screening.
11. Participants with a primary diagnosis of COPD associated with >10 pack years smoking history.
12. Participants with a primary diagnosis of asthma or asthma which is considered to be poorly controlled at screening.
13. Participants with FEV1 <25% predicted value at screening.
14. Glomerular filtration rate (eGFR) below 25 ml/min/1.73m<sup>2</sup> or requiring dialysis. This will be determined at screening.
15. Use of any investigational drugs within five times of the elimination half-life after the last dose or within 30 days, whichever is longer.
16. Unstable co-morbidities (e.g. cardiovascular disease, active malignancy) which in the opinion of the investigator would make participation in the trial not in the participant's best interest.
17. Pregnant or lactating females.
18. Women of child bearing age or male partners of women of childbearing age and not practicing a method of acceptable birth control

**Date of first enrolment**

10/08/2023

**Date of final enrolment**

24/01/2024

**Locations****Countries of recruitment**

United Kingdom

England

Northern Ireland

Scotland

Wales

Spain

**Study participating centre**

**Ninewells Hospital and Medical School**

NHS Tayside

Ninewells Avenue

Dundee

United Kingdom

DD1 9SY

**Study participating centre**

**Queen Elizabeth Hospital**

Mindelsohn Way

Edgbaston

Birmingham

United Kingdom

B15 2GW

**Study participating centre**

**University Hospital Llandough**

Penlan Road

Llandough

Penarth

United Kingdom

CF64 2XX

**Study participating centre**

**Royal Brompton Hospital**

Sydney Street

London

United Kingdom

SW3 6NP

**Study participating centre**

**Royal Papworth Hospital**  
Papworth Road  
Cambridge Biomedical Campus  
Cambridge  
United Kingdom  
CB2 0AY

**Study participating centre**  
**Hammersmith Hospital**  
Du Cane Road  
Hammersmith  
London  
United Kingdom  
W12 0HS

**Study participating centre**  
**Royal Devon and Exeter Hospital**  
Royal Devon & Exeter Hospital  
Barrack Road  
Exeter  
United Kingdom  
EX2 5DW

**Study participating centre**  
**The Princess Alexandra Hospital**  
Hamstel Road  
Harlow  
United Kingdom  
CM20 1QX

**Study participating centre**  
**Belfast City Hospital**  
51 Lisburn Rd  
Belfast  
United Kingdom  
BT9 7AB

**Study participating centre**  
**Wythenshawe Hospital**  
Southmoor Road  
Wythenshawe



Manchester  
United Kingdom  
M23 9LT

**Study participating centre**  
**Royal Infirmary of Edinburgh at Little France**  
51 Little France Crescent  
Old Dalkeith Road  
Edinburgh  
Lothian  
United Kingdom  
EH16 4SA

**Study participating centre**  
**Northwick Park Hospital**  
Watford Road  
Harrow  
United Kingdom  
HA1 3UJ

## **Sponsor information**

**Organisation**  
University of Dundee

**ROR**  
<https://ror.org/03h2bxq36>

## **Funder(s)**

**Funder type**  
Research organisation

**Funder Name**  
European Respiratory Society

**Alternative Name(s)**  
ERS

**Funding Body Type**

Private sector organisation

Funding Body Subtype

Associations and societies (private and public)

Location

Switzerland

Results and Publications

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are not expected to be made available due to commercial sensitivity.

IPD sharing plan summary

Not expected to be made available

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Basic results</a>		24/06/2025	04/07/2025	No	No
<a href="#">HRA research summary</a>			20/09/2023	No	No
<a href="#">Other files</a>	version 3	29/09/2023	26/01/2024	No	Yes
<a href="#">Participant information sheet</a>	version 1	10/11/2022	17/11/2022	No	Yes
<a href="#">Participant information sheet</a>	version 2	05/05/2023	26/05/2023	No	Yes
<a href="#">Participant information sheet</a>	version 3	29/09/2023	26/01/2024	No	Yes
<a href="#">Participant information sheet</a>	version 2	05/05/2023	26/01/2024	No	Yes
<a href="#">Participant information sheet</a>	Participant information sheet	11/11/2025	11/11/2025	No	Yes
<a href="#">Protocol file</a>	version 4	19/10/2023	06/12/2023	No	No
<a href="#">Protocol file</a>	version 5	13/12/2023	26/01/2024	No	No
<a href="#">Study website</a>	Study website	11/11/2025	11/11/2025	No	Yes