

Alternatives to prophylactic antibiotics for the treatment of recurrent urinary tract infection in women

Submission date 18/05/2016	Recruitment status No longer recruiting	<input checked="" type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 31/05/2016	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 11/05/2022	Condition category Urological and Genital Diseases	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Most women experience infective cystitis (otherwise known as urinary tract infection or UTI) at least once in their life and some get repeated episodes which are uncomfortable and stressful; this is known as recurrent urinary tract infection (rUTI) and affects around 300,000 women each year in the UK. Giving long term low dose antibiotics is the most frequently used prevention for rUTI and although reasonably effective at suppressing the infecting bacteria, it has side effects and sometimes causes bacteria to become resistant to antibiotics. There are some alternative preventative options for rUTI that don't involve antibiotics but doctors are unsure how well they work and so tend mostly to advise antibiotics. The ALTAR trial aims to compare one such alternative non-antibiotic prevention for rUTI, a drug called methenamine hippurate, against the current standard of daily low dose antibiotic to see if the methenamine is at least as good at preventing UTI and has fewer side effects, particularly a lower chance of resistant bacteria developing.

Who can participate?

Women aged at least 18 with rUTI.

What does the study involve?

The participants are randomly allocated to one of 2 groups. Those in group 1 are given a low dose antibiotic once a day for 12 months. Those in group 2 are given methenamine hippurate, a urinary antiseptic given as a tablet, twice a day for 12-months. Each participant is followed during the 12 months that they take the UTI prevention and then for 6 months afterwards to record the benefits, side effects and costs of each treatment. They are all asked to provide urine samples at the start of their time on the study and then every 3 months until the end of their participation in the study (18 months later). If a participant suffers from a UTI during the study, they are asked to provide an additional urine sample to determine the type of bug causing the infection. Blood samples are taken at the start of the study, and then at 3 months, 6 months, 9 months and 12 months. Participants may be asked provide an additional sample at 18 months depending on the results of the previous tests. Skin swabs taken from the area near the anus are also take from each participant at the start of the study and then every 6 months until the end

of their participation in the study at 18 months. Participants are also asked to complete questionnaires on the number of infections the participants have had and what effect this has had on their general health. The questionnaires also measure the impact the infection and treatment is having on the participant's life in terms of cost and benefit. They are completed at the start of the study and at 3, 6, 9, 12, 15 and 18 months during the study. Participants are given a diary to use to keep a record of any urinary infections as they happen to make completing the questionnaires easier.

What are the possible benefits and risks of participating?
Not provided at time of registration

Where is the study run from?
Seven NHS hospitals in the UK

When is the study starting and how long is it expected to run for?
February 2016 to November 2019

Who is funding the study?
National Institute for Health Research (UK)

Who is the main contact?
Gillian Watson, gillian.watson@newcastle.ac.uk

Contact information

Type(s)
Public

Contact name
Ms Gillian Watson

Contact details
Newcastle Clinical Trials Unit
Newcastle University
1-4 Claremont Terrace
Newcastle upon Tyne
United Kingdom
NE2 4AE
+44 191 208 2523
Altar.Trial@ncl.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)
201500348736

Protocol serial number
20802

Study information

Scientific Title

A multicentre, pragmatic patientrandomised noninferiority trial comparing two drugs for the prevention of recurrent urinary tract infection in women both during a 12month period of use and in the subsequent 6months following completion of the prophylactic medication.

Acronym

ALTAR

Study objectives

1. To determine the relative clinical effectiveness and cost effectiveness for the NHS of two licensed preventative treatments for women with recurrent uncomplicated urinary tract infection (UTI).
2. The null hypothesis being tested is that the nonantibiotic treatment (methenamine hippurate) is inferior to the standard treatment of extended course prophylactic antibiotic for prevention of rUTI in women.

Ethics approval required

Old ethics approval format

Ethics approval(s)

North East – Tyne & Wear South Regional Ethics Committee, 23/12/2015, ref: 15/NE/0381

Study design

Randomised; Both; Design type: Treatment, Prevention, Drug, Qualitative

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Specialty: Renal disorders, Primary sub-specialty: Renal disorders; UKCRC code/ Disease: Renal/ Noninflammatory disorders of female genital tract

Interventions

A multicentre, pragmatic patientrandomised noninferiority trial comparing two treatments for the prevention of rUTI in women during a 12month period of treatment and in the 6months following treatment completion. The standard is once daily prophylactic antibiotic, using either trimethoprim 100 mg, nitrofurantoin 50 or 100 mg depending on body weight or cefalexin 250 mg once daily for 12 months which are the recommended drugs licensed for this purpose. The choice of antibiotic will be decided by considering previous bacterial sensitivities, safety, and patient or clinician preference. The alternative (experimental) treatment is a 1 g twice daily oral urinary antiseptic methenamine hippurate for 12 months. Participants in both arms would continue to receive treatment courses of antibiotic for UTI as needed. Apart from random allocation to either option, all participants will receive usual care including use of on demand discrete treatment antibiotic courses for UTI.

Intervention Type

Other

Primary outcome(s)

1. The incidence of symptomatic antibiotic treated UTI selfreported by participants over the 12 month treatment period
2. Incremental Cost per QualityAdjusted Life Year (QALY) gained (based on responses to the Euroqol 5 dimension, 5 level (EQ5D 5L) health status questionnaire completed at baseline and 3, 6, 9 and 12 months

Key secondary outcome(s)

1. The occurrence of symptomatic UTI in the 6 months follow up period after stopping the allocated preventative therapy. (patient reported or clinically recorded)
2. Recorded participant antibiotic use
3. Antimicrobial resistance: Ecological change in terms of type of bacteria and their resistance patterns in isolates from i) midstream urine samples and ii) faecal reservoir (via optional rectal or perineal swabs) during the 12 month treatment period and in the 6 months following completion of treatment.
4. Number of microbiologicalproven UTIs during treatment and follow up
5. Occurrence of asymptomatic bacteriuria (ABU) during treatment and follow up
6. Hospitalisation due to UTI during treatment and follow up
7. HRQoL: Measured using the EQ5D 5L questionnaire
8. Incremental costs to the NHS, personal social services, and the patient at the end of the 12 month treatment and 18month followup phases measured using a cost-utility and a cost-benefit analysis
9. Model based estimates of costs, QALYs and net benefits over the longer terms, potentially over the patients estimated lifetime

Completion date

31/01/2020

Eligibility

Key inclusion criteria

1. Women aged 18 years and over
2. Women with rUTI who, in consultation with a clinician, have decided that prophylaxis is an appropriate option (to include women who have suffered at least three episodes of symptomatic UTI within the preceding 12 months or two episodes in the last 6 months or a single severe infection requiring hospitalisation)
3. Able to take a once daily oral dose of at least one of nitrofurantoin, or trimethoprim, or cephalexin
4. Able to take methenamine hippurate
5. Women who agree to take part in the trial but who are already taking Methenamine or antibiotic prophylaxis will be consented for participation and will stop their preventative therapy for a 3-month washout period. They will then be reassessed and if still eligible undergo baseline assessment and randomisation
6. Able to give informed consent for participation in trial
7. Able and willing to adhere to an 18-month study period

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Female

Total final enrolment

240

Key exclusion criteria

1. Women unable to take methenamine hippurate e.g. known allergy to methenamine hippurate, severe hepatic impairment (Child–Pugh class C, score of 10 or more), gout, eGFR < 10 ml/min, Proteus sp. as consistent proven causative organism for rUTIs
2. Women who are unable to take nitrofurantoin and trimethoprim and cephalexin
3. Women with correctable urinary tract abnormalities that are considered to be contributory to the occurrence of rUTI
4. Presence of symptomatic UTI – this will be treated and symptoms resolved prior to randomisation.
5. Pregnancy or intended pregnancy in next 12 months
6. Women who are breast feeding
7. Women already taking methenamine or antibiotic prophylaxis and declining a 3-month washout period

Date of first enrolment

07/07/2016

Date of final enrolment

31/10/2017

Locations

Countries of recruitment

United Kingdom

England

Scotland

Study participating centre

Freeman Hospital

Freeman Road

High Heaton

Newcastle upon Tyne
United Kingdom
NE7 7DN

Study participating centre

Cambridge University Hospital (Addenbrooks Hospital)

Hills Road
Cambridge
Cambridgeshire
United Kingdom
CB2 0QQ

Study participating centre

Central Manchester University Hospital, Manchester Royal Infirmary

Manchester Royal Infirmary
Oxford Road
Manchester
United Kingdom
M13 9WL

Study participating centre

St James University Hospital

Beckett Street
Leeds
West Yorkshire
United Kingdom
LS9 7TF

Study participating centre

NHS Greater Glasgow and Clyde

Queen Elizabeth University Hospital
1345 Govan Road
Glasgow
United Kingdom
G51 4TF

Study participating centre

Royal Liverpool & Broadgreen University Hospitals NHS Trust

Prescot Street

Liverpool
United Kingdom
L7 8XP

Study participating centre
Mid Yorkshire Hospitals NHS Trust
Rowan House
Aberford Road
Wakefield
West Yorkshire
United Kingdom
WF1 4EE

Study participating centre
Pennine Acute Hospitals NHS Trust
North Manchester General Hospital
Delaunays Road
Crumpsall
Manchester
United Kingdom
M8 5RB

Sponsor information

Organisation
The Newcastle upon Tyne Hospitals NHS Foundation Trust

ROR
<https://ror.org/05p40t847>

Funder(s)

Funder type
Government

Funder Name
National Institute for Health Research

Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

Funding Body Type

Government organisation

Funding Body Subtype

National government

Location

United Kingdom

Results and Publications

Individual participant data (IPD) sharing plan

Not provided at time of registration

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	protocol	09/03/2022	11/03/2022	Yes	No
Protocol article		09/11/2018		Yes	No
Funder report results	Participant information sheet	01/05/2022	11/05/2022	Yes	No
HRA research summary			26/07/2023	No	No
Participant information sheet		11/11/2025	11/11/2025	No	Yes
Study website	Study website	11/11/2025	11/11/2025	No	Yes