

An online study to change perceptions of asthma and asthma medication in adults

Submission date 05/11/2018	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 30/11/2018	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 08/04/2020	Condition category Respiratory	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

The United Kingdom has one of the highest number of people with asthma in Europe and although asthma cannot be cured, there are effective medicines available to control asthma. The most important treatment for controlling asthma is inhaled corticosteroids, also known as 'the preventer inhaler'. However, more than half of people with asthma report not taking their preventer inhaler regularly, which leads to poor asthma control with more frequent and severe asthma symptoms and an increased risk of hospital admission. It is therefore very important that people with asthma are supported to take their preventer inhaler more consistently as prescribed (called adherence). Research shows that patients' beliefs about treatment are fundamental in the decision process to take medication as prescribed, in particular doubts about personal need for the treatment and concerns about potential harm. Moreover the need for the preventer inhaler is often influenced by the experiences a patient has had with their asthma, e.g. only experiencing intermittent symptoms does not reinforce the need for regular treatment. Providing patients with a coherent 'story' that links their experience of having asthma with their asthma medication can prevent patients from developing misconceptions and misplaced concerns, and consequently improve adherence. This study will test whether a new intervention will increase the perceived need for treatment and reduce the concerns about potential harm of the treatment in patients. The intervention uses a simple, medically accurate explanation of asthma and asthma medication, in line with health psychology theory, that helps people to think of asthma and the preventer in a way that makes sense to them.

Who can participate?

People aged 18 and over with or without asthma

What does the study involve?

Participants are randomly allocated to receive either information developed based on health psychology theory (the intervention) or to standard NHS asthma information (the control). The intervention is either presented in a written format (matched to the NHS information) or as a video. This study is run in parallel in a group of people with asthma and a group without asthma. People without asthma are instructed to imagine they had just been diagnosed with asthma, which may allow the researchers to gain a better understanding of how the intervention might affect someone who has just been diagnosed with asthma. Both groups are recruited through

Amazon MechanicalTurk, an online participant panel paying small monetary rewards for study participation. Participants fill in a brief demographics questionnaire and if they report having asthma, a brief questionnaire about their asthma. After viewing the intervention, participants with and without asthma complete questionnaires measuring medication beliefs, side-effect expectations, adherence/behavioural intentions, effectiveness perceptions regarding the preventer, and general understanding of the information. After a period of two weeks, participants are invited to fill out the follow-up questionnaires (similar to the ones directly after the intervention).

What are the possible benefits and risks of participating?

The possible benefits to participants taking part are that they may increase their knowledge about asthma and asthma medication and they receive a small monetary reward for their participation (£2). Further, their input in this study will help to improve the way information is provided for people with asthma. There are no known risks to participants taking part in this study.

Where is the study run from?

UCL School of Pharmacy (UK)

When is the study starting and how long is it expected to run for?

March 2017 to March 2018

Who is funding the study?

National Institute for Health Research (NIHR) (UK)

Who is the main contact?

Caroline Katzer

Contact information

Type(s)

Scientific

Contact name

Miss Caroline Katzer

ORCID ID

<https://orcid.org/0000-0001-6077-7717>

Contact details

UCL School of Pharmacy
Department of Practice and Policy
Centre for Behavioural Medicine
BMA House, Tavistock Square
London
United Kingdom
WC1H 9JP

Additional identifiers

Protocol serial number

1

Study information

Scientific Title

Reframing asthma and inhaled corticosteroids to modify treatment beliefs: an online randomised controlled trial

Study objectives

The aim of the present study is to test a theory-based intervention which reframes asthma and inhaled corticosteroids, to evaluate whether it has the potential to modify medication and illness beliefs in a cohort of people with asthma and a cohort of people who are asthma-naïve (a proxy-group for newly diagnosed patients).

Main hypothesis: In both cohorts, lower concerns about the treatment and higher necessity for the treatment are expected within the intervention groups after receiving the intervention, compared to before the intervention and compared to the control group.

Ethics approval required

Old ethics approval format

Ethics approval(s)

University College London Research Ethics Committee, 31/07/2017, ref: 9293/003

Study design

Parallel-cohort repeated measures randomised controlled trial intervention study conducted online

Primary study design

Intentional

Study type(s)

Other

Health condition(s) or problem(s) studied

Asthma (including people without asthma as a proxy group for newly diagnosed patients)

Interventions

Participants within each cohort were randomised using the computerised block randomisation function in Qualtrics online survey software. All participants received information about asthma and its treatment (in particular inhaled corticosteroids (ICS)). The control group was presented with standard care information about asthma and asthma medication, as presented on the National Health Service (NHS) webpage. Those in the intervention groups were presented with the Balance Model of asthma (BM-intervention). The Balance Model is a medically accurate description of asthma, but by using reframing it moves asthma away from the classical, medical description to a simpler and more positive explanation that is easy to understand for patients and creates a common-sense fit between medication and illness beliefs. As the channel through which an intervention is delivered may also impact its effectiveness, participants were randomised to either receive the BM-intervention in written format (matched NHS format) or as

a four-minute video (same content as written format). Outcome data was collected immediately after the intervention and at two-week follow-up.

Intervention Type

Behavioural

Primary outcome(s)

Participants' beliefs about inhaled corticosteroids are assessed using the Beliefs about Medicines Questionnaire Specific Asthma version (BMQ-S, Horne & Weinman 2002), a 11-item questionnaire used to measure beliefs about the personal need for ICS and the concerns about the potential adverse consequences of ICS. Measurements taken at baseline (asthma cohort only), immediately post-intervention and two-week follow-up.

Key secondary outcome(s)

1. Medication adherence/adherence intention: people in the asthma cohort fill out the 10-item version of the Medication Adherence Report Scale (MARS; Horne, Weinman, & Hankins, 1999) at baseline and 2-week follow-up, and an adherence intention measure immediately post-intervention. The cohort without asthma completes the adherence intention measure immediately post-intervention and at 2-week follow-up.
2. Whether the information provided by the intervention had been understood and found to be coherent, whether perceptions about potential side-effects were reduced, whether efficacy beliefs about ICS were increased and perceptions of self were positively altered. Measured using visual analogue scales immediately post-intervention and at 2-week follow-up.
3. Beliefs about asthma, measured using the Illness Perceptions Questionnaire brief version (B-IPQ; Moss-Morris et al., 2002) at baseline, immediately post-intervention and 2-week follow-up.

Completion date

31/03/2018

Eligibility

Key inclusion criteria

1. At least 18 years old
2. Having been diagnosed with asthma for the asthma patient arm of the study

Participant type(s)

Mixed

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

891

Key exclusion criteria

Does not meet inclusion criteria

Date of first enrolment

13/11/2017

Date of final enrolment

13/03/2018

Locations**Countries of recruitment**

United Kingdom

Australia

Canada

Guernsey

Ireland

Malta

New Zealand

United States of America

Study participating centre

Online study, recruitment of participants through Amazon MechanicalTurk

United Kingdom

N/A

Sponsor information**Organisation**

University College London

ROR

<https://ror.org/02jx3x895>

Funder(s)

Funder type

Government

Funder Name

National Institute for Health Research Collaboration for Leadership in Applied Health Research and Care North Thames

Results and Publications

Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date.

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Abstract results	results presented at the ERS International Congress	19/11/2018	08/04/2020	No	No