

# Long-term comparison of BiPAP® autoSV™ versus continuous positive airway pressure in patients with mixed sleep apnoea syndrome and underlying cardiovascular diseases (Vergleich der Behandlung mit kontinuierlicher konstanter positivdruckatmung [CPAP] und der beatmung mit adaptiver servoventilation mit dem gerat BiPAP® autoSV™ bei patienten mit cheyne-stokes-atmung)

<b>Submission date</b> 16/04/2007	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 16/05/2007	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 11/12/2012	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Individual participant data

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Prof Winfried Randerath

**Contact details**  
Wissenschaftliches Institut Bethanien e.V.  
Aufderhoherstrasse 169  
Solingen  
Germany  
42699

# Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

Rand001

## Study information

Scientific Title

### Study objectives

BiPAP® autoSV™ is superior to Continuous Positive Airway Pressure (CPAP) in reducing breathing disturbances during sleep and improving cardiac function.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

Approved by the local ethics committee (Universitat Witten/Herdecke Ethik-Kommission) on the 4th September 2006.

### Study design

Randomised controlled double blind trial of BiPAP® autoSV™ therapy versus CPAP

### Primary study design

Interventional

### Secondary study design

Randomised controlled trial

### Study setting(s)

Hospital

### Study type(s)

Treatment

### Participant information sheet

### Health condition(s) or problem(s) studied

Sleep disordered breathing and cardiovascular disease.

### Interventions

Control:

Nasal Continuous Positive Airway Pressure (CPAP) is the current gold standard treatment for

Obstructive Sleep Apnoea Syndrome. CPAP can crudely be described as a small air pump attached to a mask that is worn over the nose and pumps a fixed amount of pressurised air through the nares to prevent upper airway collapse whilst sleeping.

#### **Intervention:**

Adaptive Servo-Ventilation machines (ASV) provide Expiratory Positive Airway Pressure (EPAP) or CPAP support to sustain upper-airway patency as with a CPAP machine but also modulate the Inspiratory Positive Airway Pressure (IPAP) in response to central and cheyne-stokes events.

After a baseline polysomnography (first night) patients are randomly assigned to either CPAP or BiPAP® autoSV™. They will then undergo two titration nights on therapy. After the first titration night the patients are asked for side-effects using a standardised questionnaire. On the fourth night the titrated values are validated. The CPAP level should be set to greater than or equal to 10 mbar.

After 3 months and 12 months the patients are re-evaluated polysomnographically and have outcomes measured. The devices are also read out for treatment compliance. After six months and nine months the patients are contacted by telephone to find out any problems with mask, device or compliance.

#### **Intervention Type**

Other

#### **Phase**

Not Specified

#### **Primary outcome measure**

Central AHI, measured at baseline, 3 and 12 months.

#### **Secondary outcome measures**

1. Total AHI, measured at baseline, 3 and 12 months
2. Maximal oxygen uptake (VO2 max), measured at baseline, 3 and 12 months
3. Left ventricular ejection fraction (echocardiography), measured at baseline, 3 and 12 months
4. Minimal and mean oxygen saturation (polysomnography), measured at baseline, 3 and 12 months
5. Compliance, measured at 3 and 12 months only
6. Quality of life (Minnesota questionnaire), measured at baseline, 3 and 12 months
7. Brain Natriuretic Peptide (BNP) level, measured at baseline, 3 and 12 months
8. Mortality, measured at 3 and 12 months only
9. Rate of cardiovascular complications, for example hospital admissions are examined for twelve months, measured at 3 and 12 months only

#### **Overall study start date**

01/01/2007

#### **Completion date**

01/07/2008

## **Eligibility**

#### **Key inclusion criteria**

1. Men and women 18 years of age and older
2. Arterial hypertension, ischaemic heart disease, idiopathic dilated cardiomyopathy with heart failure functional class New York Heart Association (NYHA) II and III and left ventricular ejection fraction greater than 20% (echocardiography)
3. Mixed sleep apnoea syndrome with an Apnoea Hypopnoea Index (AHI) greater than or equal to 15/hour with a proportion of central/periodic disturbances of less than 80% and a proportion of obstructive disturbances of between 20 and 50%
4. Medical treatment of their underlying heart disease has been optimised

**Participant type(s)**

Patient

**Age group**

Adult

**Lower age limit**

18 Years

**Sex**

Both

**Target number of participants**

70

**Key exclusion criteria**

1. Heart failure functional class NYHA IV
2. Myocardial infarction, unstable angina or cardiac surgery within the previous three months, obstructive disturbances of 50% or greater
3. Pregnancy
4. Patients with pure Cheyne-Stokes Respiration (CSR)/Central Sleep Apnoea (CSA) (greater than 80% of disturbances), patients with greater than 50% obstructive disturbances

**Date of first enrolment**

01/01/2007

**Date of final enrolment**

01/07/2008

**Locations****Countries of recruitment**

Germany

**Study participating centre**

Wissenschaftliches Institut Bethanien e.V.

Solingen

Germany

42699

# Sponsor information

## Organisation

Respironics International, Inc (France)

## Sponsor details

Immeuble Hermès  
20, rue Jacques Daguerre  
Rueil Malmaison  
Paris  
France  
92565

## Sponsor type

Industry

## Website

<http://www.respironics.com>

## ROR

<https://ror.org/05jz46060>

# Funder(s)

## Funder type

Industry

## Funder Name

Respironics International, Inc (France)

# Results and Publications

## Publication and dissemination plan

Not provided at time of registration

## Intention to publish date

## Individual participant data (IPD) sharing plan

## IPD sharing plan summary

Not provided at time of registration

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/08/2012		Yes	No