

A feasibility study comparing the safety and acceptability of vacuum-assisted biopsy and conventional 14-gauge core biopsy in the diagnosis of ultrasonically indeterminate and abnormal axillary lymph nodes

Submission date 12/06/2014	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 12/06/2014	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 26/10/2022	Condition category Cancer	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

<https://www.cancerresearchuk.org/about-cancer/find-a-clinical-trial/a-study-comparing-two-ways-taking-samples-tissue-lymph-nodes-under-arm>

Contact information

Type(s)

Scientific

Contact name

Miss Tania Cutts

Contact details

University of Manchester
Nightingale Centre
Southmoor Road
Manchester
United Kingdom
M23 9LT

-
tania.cutts@uhsm.nhs.uk

Additional identifiers

Protocol serial number

14676

Study information

Scientific Title

A feasibility study comparing the safety and acceptability of vacuum-assisted biopsy and conventional 14-gauge core biopsy in the diagnosis of ultrasonically indeterminate and abnormal axillary lymph nodes

Study objectives

Women with invasive breast cancer undergo ultrasound of the axilla (armpit) before surgery to detect spread of disease (metastases) to the lymph nodes. If this is normal the woman undergoes operative sentinel lymph node biopsy (SLNB), usually at the same time as surgery to the breast. If the SLNB is positive the woman undergoes axillary node clearance at a subsequent operation.

Abnormal axillary lymph nodes on ultrasound undergo needle sampling. If metastases are confirmed the woman undergoes node clearance at the same operation as surgery to the breast. The number of women who need to undergo more than one operation can be minimised by maximising the number of women with axillary metastatic disease who are diagnosed before surgery.

Ultrasound has a sensitivity of ~60% for the detection of metastatic lymph nodes. Ultrasound-guided biopsy of nodes that contain metastases has a sensitivity of ~80%. This is less than 100% because the needle may miss the part of the lymph node containing the tumour deposit. Increasing the volume of tissue removed may increase accuracy.

Vacuum-assisted biopsy (VAB) is a needle technique performed under local anaesthetic which allows more tissue to be removed than with a standard needle. Repeated samples can be taken with a single needle insertion, allowing large numbers of samples to be quickly taken. VAB of breast abnormalities is well tolerated by patients.

This study will determine whether the use of VAB in the axilla is safe and acceptable to patients and whether a larger randomised study comparing the two techniques is feasible. The results will inform the design of a larger study which will determine whether VAB significantly increases the preoperative diagnosis rate of axillary metastatic disease. If it does, its use will result in a reduction in the number of second operations in women with breast cancer.

Ethics approval required

Old ethics approval format

Ethics approval(s)

13/NW/0326

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Topic: Cancer; Subtopic: Breast Cancer; Disease: Breast

Interventions

Participants randomised to have axillary lymph node biopsy with either 14-gauge core needle biopsy device or a 13-gauge or 10-gauge vacuum biopsy device.

Study Entry : Registration only

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Proportion of women who would be willing to undergo the biopsy again.; Timepoint(s): 4 to 14 days post-biopsy

Key secondary outcome(s)

1. Procedure pain scores; Timepoint(s): Immediate and at 4-14 days post-biopsy
2. Proportion of women approached who are willing to enter the study.; Timepoint(s): AT recruitment
3. The sensitivity of vacuum biopsy relative to conventional core biopsy (preliminary data only); Timepoint(s): Following surgery

Completion date

17/06/2014

Eligibility

Key inclusion criteria

1. Women in screening and symptomatic clinics aged >35 years with breast masses scored as:
 - 1.1. M4 or M5 (mammographically suspicious or highly suspicious of malignancy) and U5 (ultrasonically highly suspicious of malignancy)
 - 1.2. and/or MRI5 (highly suspicious of malignancy on MRI) and U5
 - 1.3. and/or have histologically proven breast cancer
 - 1.4. and who have ultrasonically indeterminate or abnormal ipsilateral axillary lymph nodes (i.e. suspicious of metastatic disease)
2. Indeterminate / abnormal lymph nodes are defined as those with >2.3mm lymphoid thickness and/or focal cortical bulging and/or non-hilar blood flow and/or loss or reduction in the normal hilar fat

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

Female

Total final enrolment

80

Key exclusion criteria

1. Women who are on anticoagulants or have known clotting disorders
2. Previous ipsilateral axillary surgery
3. Target lymph node not suitable for vacuum biopsy due to its proximity to critical structures such as major blood vessels

Date of first enrolment

20/06/2013

Date of final enrolment

17/06/2014

Locations

Countries of recruitment

United Kingdom

England

Study participating centre

University of Manchester

Manchester

United Kingdom

M23 9LT

Sponsor information

Organisation

University Hospital of South Manchester NHS Foundation Trust (UK)

ROR

<https://ror.org/00he80998>

Funder(s)

Funder type

Charity

Funder Name

British Society of Breast Radiology (BSBR) (UK)

Results and Publications

Individual participant data (IPD) sharing plan

Not provided at time of registration

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/06/2016	11/07/2019	Yes	No
HRA research summary			28/06/2023	No	No
Plain English results			26/10/2022	No	Yes