

Randomised controlled trial of Narrow Band imaging with magnification (NBI) versus white light endoscopy for dysplasia detection in ulcerative colitis surveillance

Submission date 29/09/2006	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 29/09/2006	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 14/09/2017	Condition category Digestive System	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Brian P Saunders

Contact details

The Wolfson Unit 2nd Floor
North West London Hospitals NHS Trust
St Mark's Hospital
Watford Road
Harrow
United Kingdom
HA1 3UJ
+44 020 8235 4225
b.saunders@imperial.ac.uk

Additional identifiers

Protocol serial number

N0515176147

Study information

Scientific Title

Randomised controlled trial of Narrow Band imaging with magnification (NBI) versus white light endoscopy for dysplasia detection in ulcerative colitis surveillance

Study objectives

Does a new colonoscopic viewing technique called narrow band imaging (NBI) help doctors detect more patients with at least pre-cancerous area (dysplasia associated lesion or mass, DALMs) than conventional colonoscopy using white light alone?

Ethics approval required

Old ethics approval format

Ethics approval(s)

Added 17 July 2008: approval granted by Harrow Research Ethics Committee REC 06/Q0408/10

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Diagnostic

Health condition(s) or problem(s) studied

Digestive System: Ulcerative colitis

Interventions

Categorical data will be compared with chi-squared test, t-testing on Mann-Whitney U test will be used for continuous data depending on normality.

Intervention Type

Other

Phase

Not Specified

Primary outcome(s)

The study is to determine if the use of a narrow band of light can help detect more pre-cancerous or cancerous areas than the normal white light used for detection. The outcome measure is to determine the numbers of patients with at least one cancer in either group.

Key secondary outcome(s))

Added 17 July 2008:

1. Are more precancerous lesions found in total with NBI?
2. Are more advanced precancerous lesions found with NBI?
3. Are more patients found with more than one precancerous lesion with NBI?
4. How many lesions are successfully completely removed with endoscope?

Completion date

31/12/2008

Eligibility

Key inclusion criteria

1. Patients with longstanding extensive colitis, with evidence of microscopic inflammation at a previous colonoscopy
2. Patients over 18

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Not Specified

Key exclusion criteria

Pre-intubation:

1. Pregnant patients
2. Severe active colitis
3. Those unwilling or unable to give consent

Pre-randomisation:

1. Those with poor bowel preparations
2. Unable to reach caecum due to stricture

Date of first enrolment

01/03/2006

Date of final enrolment

31/12/2008

Locations

Countries of recruitment

United Kingdom

England

Study participating centre
The Wolfson Unit 2nd Floor
Harrow
United Kingdom
HA1 3UJ

Sponsor information

Organisation

Record Provided by the NHSTCT Register - 2006 Update - Department of Health

Funder(s)

Funder type

Government

Funder Name

North West London Hospitals NHS Trust (UK)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/06/2012		Yes	No
Other publications	case report	01/10/2006		Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes