

A double-blind randomised placebo-controlled trial in patients with diabetes mellitus type 2 and hypertriglyceridemia

Submission date 20/12/2005	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 20/12/2005	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 17/12/2015	Condition category Nutritional, Metabolic, Endocrine	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr M.V. Huisman

Contact details
Leiden University Medical Center
Department of General Internal Medicine
Albinusdreef 2, C2-R
P.O. Box 9600
Leiden
Netherlands
2300 RC
+31 (0)71 625 9111
m.v.huisman@lumc.nl

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

N/A

Study information

Scientific Title

The DALI study: a double-blind randomised placebo-controlled trial in patients with diabetes mellitus type 2 and hypertriglyceridemia

Acronym

DALI

Study objectives

Higher doses of statins will result in additional improvement of the diabetic lipid profile.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics approval received from the local medical ethics committee

Study design

Multicentre randomised double-blind placebo-controlled parallel-group trial

Primary study design

Interventional

Secondary study design

Randomised parallel trial

Study setting(s)

Not specified

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Diabetes Mellitus type II (DM type II)

Interventions

Patients who met the in- and exclusion criteria started with a placebo run-in period. If the lipid levels were still within the inclusion range after two weeks, patients were randomised to treatment with atorvastatin 10 mg, 80 mg, or placebo, administered once daily in the morning. Patients randomised to atorvastatin 80 mg started with 40 mg for four weeks after which the dose was increased to 80 mg. The total treatment period was 30 weeks.

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Atorvastatin

Primary outcome measure

The effect of atorvastatin 10 mg and 80 mg on the reduction of triglyceride levels in patients with diabetes mellitus type 2 and hypertriglyceridemia.

Secondary outcome measures

The effects on other aspects of diabetic dyslipidemia.

Overall study start date

17/06/1998

Completion date

31/01/2000

Eligibility**Key inclusion criteria**

1. Diabetes mellitus type 2 for greater than 1 year
2. Male or female
3. HbA1c 10% or lower
4. Fasting total cholesterol level between 4.0 and 8.0 mmol/L
5. Fasting triglycerides level between 1.5 and 6.0 mmol/L.

Participant type(s)

Patient

Age group

Not Specified

Sex

Both

Target number of participants

217

Key exclusion criteria

1. History of myocardial infarction, Percutaneous Transluminal Coronary Angioplasty (PTCA), Coronary Artery Bypass Graft (CABG), clinical symptoms of manifest coronary artery disease (greater than grade II of the Canadian Cardiovascular Society), severe or unstable angina pectoris (greater than grade II), clinically manifest heart failure (greater than grade II New York Heart Association [NYHA]) and severe cardiac arrhythmias
2. Premenopausal women, patients with acute liver disease or hepatic dysfunction, impaired

renal function (plasma creatinine greater than 150 mmol/l), a history of partial ileal bypass surgery, any surgical procedure or any systemic inflammatory disease within the last three months before randomisation, malignancies, vasculitis, rheumatic arthritis, idiopathic lung fibrosis, ulcerative colitis or Crohns disease

3. Patients who consumed more than 4 alcoholic drinks per day or who used systemic steroids, androgens, cyclosporin, other immunosuppressive drugs, erythromycin or mibefradil

Date of first enrolment

17/06/1998

Date of final enrolment

31/01/2000

Locations

Countries of recruitment

Netherlands

Study participating centre

Leiden University Medical Center

Leiden

Netherlands

2300 RC

Sponsor information

Organisation

Erasmus Medical Centre (Netherlands)

Sponsor details

Dr Molewaterplein 40/50

Rotterdam

Netherlands

3000 CA

Sponsor type

University/education

Website

<http://www.erasmusmc.nl/>

ROR

<https://ror.org/018906e22>

Funder(s)

Funder type

Industry

Funder Name

Parke Davis B.V. (The Netherlands)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/08/2001		Yes	No
Results article	results	01/04/2015		Yes	No