Stapled side-to-side anastomosis versus sutured end-to-end anastomosis following resection for primary or recurrent Crohn's disease

Submission date	Recruitment status No longer recruiting	Prospectively registered		
09/09/2005		☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
09/09/2005	Completed	[X] Results		
Last Edited 04/08/2009	Condition category Digestive System	[] Individual participant data		

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Robin Susan McLeod

Contact details

Mount Sinai Hospital 449-600 University Avenue Toronto Canada M5G 1X5 +1 416-586-8534 rmcleod@mtsinai.on.ca

Additional identifiers

Protocol serial number

MCT-50012

Study information

Scientific Title

Multicentre randomised controlled trial to compare stapled side-to-side anastomosis with sutured end-to-end anastomosis following resection for primary or recurrent Crohn's disease

Acronym

CAST (The Canadian and American Surgical Crohn's Disease Trial)

Study objectives

Patients having an ileocolonic resection for Crohns Disease will have a lower risk of recurrent disease if they have a wide side-to-side anastomosis compared to a end-to-end anastomosis.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Mount Sinai Hospital Research Ethics Board approved on the 28th November 2000

Study design

Randomised controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Crohns disease

Interventions

Group 1: Hand sewn anastomosis Group 2: Stapled anastomosis

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Severe endoscopic recurrence at one year follow-up.

Key secondary outcome(s))

- 1. Symptomatic recurrence (up to 24 months after)
- 2. Post-operative complication rates
- 3. Operative times

Completion date

31/07/2005

Eligibility

Key inclusion criteria

- 1. Women and men aged greater than 16 years
- 2. Elective ileocolonic resection
- 3. Colonoscopy and Small Bowel Endoscopy (SBE) completed with the last 3 years
- 4. Crohns Disease involving the terminal ileum plus or minus right colonic disease
- 5. No other sites of involvement with Crohns disease in the Gastrointestinal (GI) tract current or past
- 6. Willing to return for colonoscopy at 12 months

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

- 1. Require emergency surgery
- 2. Have sites of Crohns involvement elsewhere, except for minimal perianal disease
- 3. Have had strictureplasties or other resections for Crohns disease performed either at this operation or previously, the exception being resections for fistula disease in bowel not affected by Crohns disease
- 4. Require a bypass procedure or defunctioning ileostomy during the index procedure
- 5. Have compromised renal function (serum creatinine level greater than 130 mmol/l or 1.5 mg/l)
- 6. Prednisone, budesonide, 5-Aminosalicylic Acid (5-ASA) medications, ciprofloxacin, metronidazole, cyclosporin and Anti-Tumour Necrotising Factor (Anti-TNF) and any other medications used to treat Crohns cannot be discontinued postoperatively
- 7. Are unable or willing to give informed consent and return for a colonoscopy at 12 months

Date of first enrolment

01/01/2002

Date of final enrolment

31/07/2005

Locations

Countries of recruitment

Canada

Study participating centre

Mount Sinai Hospital

Toronto Canada M5G 1X5

Sponsor information

Organisation

Mount Sinai Hospital (Toronto) (Canada)

ROR

https://ror.org/05deks119

Funder(s)

Funder type

Charity

Funder Name

Canadian Institutes of Health Research (CIHR) (Canada) - http://www.cihr-irsc.gc.ca (ref: MCT-50012)

Funder Name

Crohns & Colitis Foundation of America (USA)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/05/2009		Yes	No