

Comparison of two ready-to-use systems specially designed for physiological intracytoplasmic sperm injection (ICSI): PICSI® versus Sperm Slow™, a prospective randomised trial

Submission date 24/08/2010	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered
Registration date 11/10/2010	Overall study status Completed	<input type="checkbox"/> Protocol
Last Edited 18/12/2020	Condition category Pregnancy and Childbirth	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
		<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Dr Lodovico Parmegiani

Contact details
GynePro Medical Centers
Via T. Cremona, 8
Bologna
Italy
40137
-
l.parmegiani@gynepro.it

Additional identifiers

Protocol serial number
N/A

Study information

Scientific Title

Comparison of two ready-to-use systems specially designed for sperm-hyaluronic acid (HA) binding selection before intracytoplasmic sperm injection (physiological ICSI): PICSI® versus Sperm Slow™, a prospective randomised trial

Study objectives

The objective of this study is to evaluate the role of hyaluronic acid (HA) for sperm selection prior to intracytoplasmic sperm injection (ICSI) comparing two ready-to-use systems specially designed for sperm-HA binding selection: a plastic culture dish with microdots of HA hydrogel attached to the bottom interior of the dish (PICSI® Sperm Selection Device, MidAtlantic Diagnostic) or a viscous medium containing HA (Sperm Slow™, MediCult). These systems, in which the spermatozoa are selected for their capacity to bind to HA (HA-ICSI), allow the execution of a more "physiological" ICSI than conventional PVP-ICSI.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Institutional Medical Ethics Committee of GynePro Medical Center approved in December 2007

Study design

Single centre prospective randomised study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Intracytoplasmic sperm injection (ICSI)

Interventions

100 ICSI treatments randomly carried out with PICSI® Sperm Selection Device or with Sperm Slow™ for sperm selection. Randomisation process conducted with sealed envelopes, assigning 50 subjects to each treatment group.

Controlled ovarian stimulation with gonadotropin-releasing hormone analogs in combination with a graded gonadotropin administration. Oocyte retrieval performed 35 hours after ovulation induction with either 5,000 or 10,000 IU of human chorionic gonadotropin (hCG). Oocytes culture at 37°C in an atmosphere of 6% CO₂. Complete removal of cumulus mass and corona cells by enzymatic digestion of recombinant hyaluronidase (SynVibro® Cumulase® MediCult) and by gentle mechanical aspiration with plastic pipettes (Stripper Tips® MidAtlantic Diagnostic).

Assessment of the nuclear maturation stage of the denuded oocytes. Strict selection of the metaphase II oocytes (MII) by morphological features (zona pellucida thickness, perivitelline space size, oocyte shape, cytoplasm colour and granularity, presence of vacuoles and first polar body morphology) under an inverted microscope with Hoffman modulation contrast.

Classification of "high quality oocytes": colourless and of regular shape, with regular zona pellucida and small perivitelline space without debris, homogeneous cytoplasm and no vacuoles or granulations.

Insemination by ICSI of the best available MII oocytes, in accordance with the Italian law regulating Assisted Reproductive Technology (cryopreservation of the supernumerary MII oocytes reaching the "high quality" standards).

PICSI® Sperm Selection Device procedure as described in www.midatlanticdiagnostics.com.

Sperm Slow™ ICSI procedure:

Spermatozoa must first be treated with a two-layer density gradient system or via Swim-Up. Thus, on a Petri dish, a 2 µL droplet with suspension of treated spermatozoa is connected with a pipette tip to a 5 µL droplet fresh culture medium (Sperm Preparation Medium, MediCult). Simultaneously, A 5 µL droplet of HA-containing medium (Sperm Slow™, MediCult) is connected with a pipette tip to the 5 µL droplet fresh culture medium. The spermatozoa on this Petri dish are incubated for 15 min at 37°C under oil (Liquid Paraffin, MediCult). Spermatozoa bound to HA in the junction zone of the 2 droplets can be selected and easily detached by injecting pipette (ICSI Micropipets, Humagen Fertility Diagnostics) and subsequently injected into oocytes.

Controlled ovarian stimulation duration: from 12 to 20 days

Oocyte recovery, ICSI, embryo culture and embryo transfer duration: from 3 to 5 days

The follow up will be completed at confirmation of clinical pregnancy by ultrasound check of gestational chamber, 30 days after the embryo transfer.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

1. Oocyte fertilisation rate, measured 15 - 22 hours after oocyte insemination
2. Embryo cleavage rate, measured at day of embryo transfer (2, 3 or 5 days after oocyte insemination)
3. Top-quality embryo rate, measured at day of embryo transfer (2, 3 or 5 days after oocyte insemination)
4. Pregnancy rate, measured 30 days after the embryo transfer
5. Implantation rate, measured 30 days after the embryo transfer

Key secondary outcome(s)

Duration of ICSI procedure, measured at time of oocyte insemination

Completion date

01/03/2011

Eligibility

Key inclusion criteria

1. Infertile women not older than 41 years undergoing ICSI treatment at GynePro Medical Center
2. Semen parameters of the male partner:
 - 2.1. Presence of ejaculate motile spermatozoa with total sperm number greater than or equal to 1,000,000
 - 2.2. Sperm motility greater than or equal to 5%

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Total final enrolment

100

Key exclusion criteria

1. Women older than 41 years old
2. Patients with male partners with testicular spermatozoa or severe oligoasthenoteratozoospermia (total sperm number less than 1,000,000 and sperm motility less than 5%)

Date of first enrolment

01/09/2010

Date of final enrolment

01/03/2011

Locations**Countries of recruitment**

Italy

Study participating centre

GynePro Medical Centers

Bologna

Italy

40137

Sponsor information

Organisation

GynePro Medical Centers (Italy)

ROR

<https://ror.org/03segdh23>

Funder(s)

Funder type

Industry

Funder Name

GynePro Medical Centers (Italy)

Funder Name

MidAtlantic Diagnostic (USA)

Results and Publications

Individual participant data (IPD) sharing plan**IPD sharing plan summary**

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/09/2012	18/12/2020	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes