To evaluate efficacy and tolerability of deferred androgen deprivation therapy +/- upfront CRyOtherapy in men with localised radiation recurrent Prostate cancer (RRPC)

| Submission date | Recruitment status No longer recruiting | Prospectively registered | | |
|-------------------|--|---|--|--|
| 26/07/2011 | | <pre>Protocol</pre> | | |
| Registration date | Overall study status Stopped Condition category Cancer | Statistical analysis plan | | |
| 26/07/2011 | | [X] Results | | |
| Last Edited | | Individual participant data | | |
| 26/10/2022 | | Record updated in last year | | |

Plain English summary of protocol

https://www.cancerresearchuk.org/about-cancer/find-a-clinical-trial/a-trial-cryotherapy-and-hormone-therapy-for-prostate-cancer-come-back-after-radiotherapy-crop

Contact information

Type(s)

Scientific

Contact name

Ms Judith Dixon

Contact details

1053 Great Western Road Glasgow United Kingdom G12 0YN +44 141 301 7945 Judith.Dixon@glasgow.ac.uk

Additional identifiers

Protocol serial number 10051

Study information

Scientific Title

randomised controlled trial of deferred androgen deprivation therapy +/- upfront CRyOtherapy in men with localised radiation recurrent Prostate cancer (RRPC) to evaluate efficacy and tolerability

Acronym

CROP

Study objectives

This is an open, multi-centre, phase III randomised controlled trial (RCT) to investigate the role of salvage prostate cryotherapy in patients with localised recurrent prostate cancer following radiotherapy. Recruited patients will be randomised into one of two arms: an intervention arm with salvage prostate cryotherapy followed by non-surgical management at failure (deferred ADT) or a control arm with non-surgical management (deferred ADT) only.

The proposed RCT will evaluate the outcome among patients managed with deferred androgen ablation with or without upfront salvage prostate cryotherapy for their RRPC. The endpoint will be distant metastasis free survival (DMFS). The toxicity profile, assessment of patient quality of life and health economic analysis also form integral parts of the study.

Primary Objective

To determine the efficacy of salvage prostate cryotherapy in RRPC in combination with deferred androgen deprived therapy (ADT) in patients with radiation recurrent prostate cancer

- 1. To evaluate the safety and tolerability of prostate cryotherapy in RRPC
- 2. To determine the side effect profile of salvage cryotherapy and its impact on the quality of life for patients
- 3. To determine within this study the optimal health economic model for cost-effectiveness analysis to assess cryotherapy against current management plans
- 4. To determine the likely costs to the NHS if prostate cryotherapy is adopted widely for RRPC

Ethics approval required

Old ethics approval format

Ethics approval(s)

West of Scotland REC1 approved on 07 April 2011, ref: 11/S0703/2

Study design

Randomised; Interventional; Design type: Treatment

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Prostate cancer

Interventions

Patients will be randomised equally across the two study treatment arms. 270 patients will receive upfront salvage prostate cryotherapy followed by deferred androgen deprivation therapy. 270 patients will receive deferred androgen deprivation therapy alone

Deferred ADT, The option of intermittent or continuous ADT is permitted at the discretion of the Investigator, but the intention is to be declared at randomisation.

Salvage prostate cryotherapy, Salvage prostate cryotherapy will be given upfront to those patients randomised to Arm B (cryotherapy + deferred ADT);

Follow Up Length: 90 month(s); Study Entry: Single Randomisation only

Intervention Type

Other

Phase

Phase III

Primary outcome(s)

Distant Metastasis Free Survival (DMFS); Timepoint(s): DMFS assessed annually by MRI and bone scan

Key secondary outcome(s))

- 1. Acute and late side effects of cryotherapy; Timepoint(s): Meausred using NCI CTCAE; Efficacy; Timepoint(s): PSA 3 monthly in year 1, annually thereafter
- 2. Prostate biopsy at 6 months post cryotherapy; Functional Status; Timepoint(s): validated questionnaires at 3, 6, 12 months and annually thereafter

Completion date

01/06/2020

Reason abandoned (if study stopped)

Lack of efficacy

Eligibility

Key inclusion criteria

- 1. Histologically confirmed relapsed prostate cancer following previous treatment with radiation therapy (either external beam or brachytherapy) for either organ confined or non-metastatic locally advanced prostate cancer, namely T1-3aN0M0 disease
- 2. Life expectancy of at least 5 years
- 3. Clinical/radiological T1c-T3a
- 4. Prostrate specific antigen (PSA) level = 20 ng/ml
- 5.. Aged 18 years or over
- 6. Eastern Cooperative Oncology Group Performance Status Scale (ECOG PS) = 0 or 1 or 2
- 7. Ability to provide informed consent
- 8. Adequate haematological function as defined by haemoglobin (Hb) = 100g/L; platelets = $100 \times 109/L$; neutrophils = $1.5 \times 109/L$
- 9. Adequate biochemical function as defined by bilirubin = 1.5 upper limit of normal (ULN); alanine aminotransferase (ALT), aspartate aminotransferase (AST) = $2.5 \times 100 \times 10^{-5} = 1.5 \times 1$

ULN OR calculated/measured creatinine clearance = 60mls/min (as defined by Cockcroft and Gault formula); Target Gender: Male; Lower Age Limit 18 years

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Male

Key exclusion criteria

- 1. Lymph node >10mm short axis on magnetic resonance imaging (MRI) abdomen and pelvis
- 2. Previous transurethral resection of prostate gland with evidence of a significant defect (>10mm in width) on transrectal ultrasound scan
- 3. Significant lower urinary tract symptoms, including bladder outflow obstructive symptoms
- 4. History of abdomino-perineal resection of rectum.
- 5. Known coagulation disorder
- 6. Complex perianal fistula
- 7. Previous combined external beam radiotherapy and brachytherapy to the prostate
- 8. Significant symptoms/toxicity related to the rectum following radiotherapy National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE) grade 3 or above
- 9. Failed androgen deprivation therapy as second line therapy for radiation recurrent prostate cancer
- 10. Prior cryotherapy to the prostate
- 11. Any evidence of severe or uncontrolled systemic conditions (e.g. severe hepatic impairment, interstitial lung disease [bilateral, diffuse, parenchymal lung disease]) or current unstable or uncompensated respiratory or cardiac conditions which make it undesirable for the patient to participate in the study or which could jeopardise compliance with the protocol 12. Other prior malignancy with estimated =30% chance of relapse within 5 years

Date of first enrolment

01/06/2011

Date of final enrolment

01/06/2020

Locations

Countries of recruitment

United Kingdom

Scotland

Study participating centre 1053 Great Western Road Glasgow

United Kingdom G12 0YN

Sponsor information

Organisation

NHS Greater Glasgow & Clyde (UK)

ROR

https://ror.org/05kdz4d87

Funder(s)

Funder type

Charity

Funder Name

Clinical Trials Awards and Advisory Committee (CTAAC) (UK)

Results and Publications

Individual participant data (IPD) sharing plan

Not provided at time of registration

IPD sharing plan summary

Not provided at time of registration

Study outputs

| Output type | Details | Date created | Date added | Peer reviewed? | Patient-facing? |
|-------------------------------|-------------------------------|--------------|------------|----------------|-----------------|
| Results article | results | 19/10/2015 | | Yes | No |
| Participant information sheet | Participant information sheet | 11/11/2025 | 11/11/2025 | No | Yes |
| Plain English results | | | 26/10/2022 | No | Yes |