

# Comprehensive studies of a new improved oral rehydration solution for diarrhoeal disease: physiological and molecular studies, clinical trials, and acceptability and efficacy studies in a rural south Indian community

<b>Submission date</b> 22/07/2005	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 22/07/2005	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 06/02/2015	<b>Condition category</b> Signs and Symptoms	<input type="checkbox"/> Individual participant data

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Mr Balakrishnan S Ramakrishnan

**Contact details**  
Christian Medical College  
Williams Research Building  
Department of Gastrointestinal Science  
Ida Scudder Road  
Vellore  
India  
632004  
+91 (0)416 2282052  
rama@cmcvellore.ac.in

## Additional identifiers

**EudraCT/CTIS number**

**IRAS number**

**ClinicalTrials.gov number**

**Secondary identifying numbers**

063150

## **Study information**

### **Scientific Title**

Comprehensive studies of a new improved oral rehydration solution for diarrhoeal disease: physiological and molecular studies, clinical trials, and acceptability and efficacy studies in a rural south Indian community

### **Study objectives**

The hypothesis is that addition of amylase resistant starch to a low osmolality Oral Rehydration Solution (ORS) will reduce the duration and severity of diarrhoea in a variety of diarrhoeal diseases including cholera and non-cholera diarrhoea. The studies consist of two randomised trials in adults and children respectively comparing the newly recommended low osmolar ORS to low osmolar ORS with added amylase resistant starch. The objective is to determine whether subjects in the trial arm will have reduced stool output and duration of diarrhoea.

The overall trial end date was changed from 15/03/2006 to 31/01/2007.

### **Ethics approval required**

Old ethics approval format

### **Ethics approval(s)**

Not provided at time of registration

### **Study design**

Randomised single-blinded controlled clinical trial

### **Primary study design**

Interventional

### **Secondary study design**

Randomised controlled trial

### **Study setting(s)**

Not specified

### **Study type(s)**

Treatment

### **Participant information sheet**

### **Health condition(s) or problem(s) studied**

Diarrhoea

## **Interventions**

Eligible subjects will be randomised to one of two treatment arms to receive either reduced osmolarity ORS or reduced osmolarity ORS along with amylase resistant starch. All other treatment measures will be identical in the two groups. ORS will be administered according to World Health Organisation (WHO) treatment guidelines for management of acute diarrhoea. Stool consistency and weight will be determined by personnel unaware of the treatment received by the patient.

## **Intervention Type**

Drug

## **Phase**

Not Applicable

## **Drug/device/biological/vaccine name(s)**

Amylase resistant starch, low osmolarity Oral Rehydration Solution (ORS)

## **Primary outcome measure**

1. Stool weight in successive 12 hour periods
2. Stool weight in the second 24 hours
3. Time to last unformed stool
4. Requirement for unscheduled intravenous fluids

## **Secondary outcome measures**

No secondary outcome measures

## **Overall study start date**

01/01/2005

## **Completion date**

31/01/2007

# **Eligibility**

## **Key inclusion criteria**

In the adult study, all adult male patients presenting to the Emergency Service with acute severe watery diarrhoea of less than three days' duration, with moderate to severe dehydration, will be enrolled after informed written consent.

In a separate trial, male children aged six months to five years with diarrhoea (defined as more than three watery stools/24 hours) of three days or less, with mild or moderate dehydration, or those with severe dehydration who have received intravenous rehydration, will be enrolled after informed written consent from the parent.

## **Participant type(s)**

Patient

## **Age group**

Mixed

**Sex**

Male

**Target number of participants**

To be added

**Key exclusion criteria**

In the adult trial, subjects without clinical dehydration, those with presence of blood in stool, and presence of significant systemic illness unrelated to the diarrhoea, will be excluded.

In the children's trial, bloody diarrhoea, concurrent severe illness (such as pneumonia, meningitis or infections requiring antibiotics), and severe malnutrition Grade III and IV will be treated as exclusion criteria.

**Date of first enrolment**

01/01/2005

**Date of final enrolment**

31/12/2006

**Locations****Countries of recruitment**

India

**Study participating centre**

Christian Medical College

Vellore

India

632004

**Sponsor information****Organisation**

Christian Medical College (India)

**Sponsor details**

c/o George M Chandy

Ida Scudder Road

Vellore

India

632004

+91 (0)416 2282052

director@cmcvellore.ac.in

**Sponsor type**

Hospital/treatment centre

**Website**

<http://cmcvellore.ac.in>

**ROR**

<https://ror.org/00c7kvd80>

## Funder(s)

**Funder type**

Charity

**Funder Name**

Wellcome Trust

**Alternative Name(s)****Funding Body Type**

Private sector organisation

**Funding Body Subtype**

International organizations

**Location**

United Kingdom

## Results and Publications

**Publication and dissemination plan**

Not provided at time of registration

**Intention to publish date****Individual participant data (IPD) sharing plan****IPD sharing plan summary**

Not provided at time of registration

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	13/02/2008		Yes	No