

Oral challenge test with gluten versus placebo to identify patients with nonceliac gluten sensitivity (NCGS)

Submission date 28/06/2014	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 31/07/2014	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 24/02/2015	Condition category Digestive System	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Non-celiac gluten sensitivity (NCGS) is a condition with symptoms similar to that of celiac disease. These include depression, abdominal pain, bloating, diarrhoea, constipation, fatigue, joint pain and a foggy mind. However, despite the number of symptoms common to both conditions, people with NCGS do not test positive for celiac disease or a wheat allergy. Indeed, there is a debate on whether the disease exists at all. The aim of this study is to test the effect of low amounts (doses) of gluten in patients suspected of having NCGS.

Who can participate?

Adult patients with persistent NCGS symptoms and having eaten a diet containing gluten for at least the last two months.

What does the study involve?

Participants are randomly assigned to one of two groups. Those in group 1 are given a pill containing 4.375 grams of gluten for a week. Those in group 2 are given a dummy (placebo) pill made out of rice starch. Then, after a one week wash-out period (where no treatment is given to either group), the groups cross over with Group 1 taking the placebo and group 2 taking the gluten containing pill for a week. The effects of taking the gluten compared with the placebo are then analysed.

What are the possible benefits and risks of participating?

There are no risks associated with taking part in the trial as participants are not taking drugs but common components of a normal diet. Benefits associated with taking part in our trial might be the diagnosis of true NCGS.

Where is the study run from?

St. Matteo Hospital (Italy)

When is the study starting and how long is it expected to run for?

October 2012 to January 2014

Who is funding the study?
St. Matteo Hospital (Italy)

Who is the main contact?
Dr Antonio Sabatino
a.disabatino@smatteo.pv.it

Contact information

Type(s)
Scientific

Contact name
Prof Gino Roberto Corazza

Contact details
First Department of Medicine
St Matteo Hospital Foundation
Piazzale Golgi 19
Pavia
Italy
27100

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
20120011203

Study information

Scientific Title
Effect of low gluten doses in patients suspected of having nonceliac gluten sensitivity (NCGS): a randomized, double-blind, placebo-controlled cross-over gluten challenge trial

Acronym
NCGS

Study objectives
We hypothesized that even low gluten doses could worsen overall (intestinal and extraintestinal) symptoms in patients strongly suspected of having NCGS

Ethics approval required
Old ethics approval format

Ethics approval(s)

Ethics Committee of the St. Matteo Hospital Foundation - Pavia (Italy), 19/10/2012, ref. 4468 /2012

Study design

Randomized double-blind placebo-controlled cross-over trial

Primary study design

Interventional

Secondary study design

Randomised cross over trial

Study setting(s)

Hospital

Study type(s)

Diagnostic

Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

Health condition(s) or problem(s) studied

Nonceliac gluten sensitivity

Interventions

Patients will be asked to fill in a daily questionnaire in order to assess a rating scale of both intestinal and extraintestinal symptoms over a 5-week period. Participants will be asked to follow a strict gluten-free diet (GFD) starting one week before the randomization and continuing until the end of the study period. After the first week of GFD, patients will be given either gastrosoluble capsules containing purified wheat gluten (10 capsules ingested in no more than two times over the day, corresponding to a daily gluten intake of 4.375 grams, equivalent to about 2 slices of white bread) or gastrosoluble capsules containing rice starch (10 capsules ingested in no more than two times over the day, corresponding to a daily rice starch intake of 4.375 grams) as placebo for one week. At the end of the first treatment week, patients from both arms will continue only their wash-out from gluten, without taking a capsule. Subsequently, individuals belonging to the first arm will be given placebo capsules, while individuals belonging to the second arm will be given gluten capsules. After the second treatment week, all patients will continue with their wash-out from dietary gluten. During the trial, patients will undergo five outpatient weekly appointments, during which patients will provide their filled questionnaires and investigators will count any unused capsules remaining in capsule dispensers.

Intervention Type

Other

Phase

Not Applicable

Primary outcome measure

The primary outcome is the change in the weekly overall symptom score, as assessed by the sum of intestinal and extraintestinal scores, between the 1-week treatment with gluten and the 1-week treatment with placebo. The 15 intestinal symptoms, which patients will be asked to grade daily from 0 to 3 (0=absent; 1=mild; 2=relevant, 3=severe and interfering with daily activities) are:

1. Abdominal pain
2. Abdominal bloating
3. Wind
4. Diarrhea
5. Borborygmus
6. Reduced consistency of stools
7. Increased consistency of stools
8. Constipation
9. Urgency
10. Incomplete evacuation
11. Nausea
12. Heartburn
13. Belching
14. Acid regurgitation
15. Epigastric pain.

The 13 extraintestinal symptoms, which patients will be asked to grade daily from 0 (absent) to 1 (present), are:

1. Tiredness
2. Malaise
3. Headache
4. Depression
5. Anxiety
6. Foggy mind
7. Aphthous stomatitis
8. Paresthesia
9. Arthralgia
10. Myalgia
11. Asthma
12. Rhinitis
13. Skin rash

Secondary outcome measures

Secondary outcomes are:

1. The change in individual symptom scores between the 1-week treatment with gluten and the 1-week treatment with placebo
2. The identification of patients with true NCGS
3. To verify whether laboratory parameters at baseline are predictive of true NCGS.

True NCGS patients are defined as having at the end of the trial a delta overall score - calculated by subtracting the weekly overall score under placebo from the weekly overall score under gluten- higher than the mean delta overall score plus two standard deviations. Laboratory predictors include serum IgG AGA, fecal calprotectin, HLA genotyping and intraepithelial lymphocyte density.

Overall study start date

20/10/2012

Completion date

31/01/2014

Eligibility

Key inclusion criteria

1. Adult patients referred to our outpatients' clinics because of persistence of relevant intestinal and extraintestinal symptoms believed by them to be caused by the ingestion of food containing even low doses of gluten.
2. Under gluten-containing diet at the time of screening since at least two months
3. Happy to undergo ad hoc screening, including:
 - 3.a. Serum determination of IgA anti-transglutaminase and anti-endomysial antibodies
 - 3.b. IgG anti-gliadin antibodies (AGA)
 - 3.c. Total IgA and IgE
 - 3.d. Wheat-specific IgE
 - 3.e. Upper endoscopy with collection of multiple duodenal biopsies
 - 3.f. HLA genotyping
 - 3.g. Fecal calprotectin
 - 3.h. Lactose breath test

Participant type(s)

Patient

Age group

Adult

Sex

Both

Target number of participants

At least 58 randomized participants

Key exclusion criteria

1. Other significant gastrointestinal diseases, such as inflammatory bowel disease or cirrhosis,
2. Other clinically significant comorbidities such as diabetes, use of nonsteroidal anti-inflammatory agents or systemic immunomodulators
3. Pregnancy
4. Inability to give written informed consent
5. Reported psychiatric disorder
6. Excessive alcohol intake

Only patients, who at the end of the screening are found not to be affected by celiac disease, wheat allergy, lactose or FODMAP intolerance, *Helicobacter pylori* infection, giardiasis, and who complain of relevant gluten-dependent symptoms that is affecting the overall quality of life, will be randomized according to a computer-generated list of random numbers held by an independent observer to either the gluten or the placebo treatment group.

Date of first enrolment

20/10/2012

Date of final enrolment

31/01/2014

Locations

Countries of recruitment

Italy

Study participating centre

St Matteo Hospital Foundation

Pavia

Italy

27100

Sponsor information

Organisation

St. Matteo Hospital Foundation (Italy)

Sponsor details

Piazzale Golgi 5

Pavia

Italy

27100

Sponsor type

Hospital/treatment centre

ROR

<https://ror.org/05w1q1c88>

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

St Matteo Hospital Foundation, Pavia (Italy)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/09/2015		Yes	No