# Study of the beneficial effects of a combination of food ingredients on cardiovascular risk factors in overweight subjects

Submission date	Recruitment status	[X] Prospectively registered
14/09/2017	No longer recruiting	[_] Protocol
Registration date	Overall study status	[] Statistical analysis plan
15/09/2017	Completed	[_] Results
Last Edited	Condition category	Individual participant data
15/09/2017	Nutritional, Metabolic, Endocrine	[] Record updated in last year

### Plain English summary of protocol

Background and study aims

The aim of this study is to test whether a combination of food ingredients can decrease cardiovascular (heart disease) risk factors by reducing visceral adiposity (abdominal fat), hypercholesterolemia (high blood cholesterol) and endothelial (blood vessel) dysfunction. These ingredients are based on extracts rich in polyphenols and non-digestible fibers already known individually for their health benefits.

Who can participate? Overweight people aged 18 and over

What does the study involve?

Participants are randomly allocated to take either the study medication or a placebo (dummy) capsule twice a day, 2 capsules each time (4 capsules per day), for 3 months. Clinical data, blood samples and faeces samples are collected and the brachial artery is examined by ultrasound at the start of the study, at the end of the first, second and third months of treatment, and one month after the treatment has finished.

What are the possible benefits and risks of participating? Treatment does not cause any adverse health effects. This study does not involve any risks apart from those related to a conventional blood collection procedure or possible gastrointestinal discomfort.

Where is the study run from? CHU UCL Namur Godinne (Belgium)

When is the study starting and how long is it expected to run for? May 2017 to May 2019

Who is funding the study? Competitivity pole WAGRALIM (Public service of Wallonia region) (Belgium) Who is the main contact? Prof. Laurence Galanti laurence.galanti@uclouvain.be

## **Contact information**

**Type(s)** Scientific

**Contact name** Prof Laurence Galanti

**Contact details** Avenue du Dr Gaston Thérasse 1 Yvoir Belgium 5530 +32 (0)81 42 32 40 laurence.galanti@uclouvain.be

## Additional identifiers

EudraCT/CTIS number

**IRAS number** 

ClinicalTrials.gov number

Secondary identifying numbers B039201733170

## Study information

## Scientific Title

Study of the beneficial effects of a combination of food ingredients on cardiovascular risk factors in overweight subjects

Acronym ADIPOSTOP

### Study objectives

Decrease of cardiovascular risk factors (waist circumference, BMI, inflammatory parameters, lipid balance and oxidative status, endothelial function, intestinal microbiota) by taking a new combination of food ingredients.

**Ethics approval required** Old ethics approval format

Ethics approval(s)

Comité d'éthique médicale de CHU UCL Namur, site Godinne (Ethics committee of CHU UCL Namur, Godinne), 11/07/2017, ref: CE Mont-Godinne: 79/2017

### Study design

Interventional single-center double-blind randomized placebo-controlled study

**Primary study design** Interventional

#### Secondary study design

Randomised controlled trial

### Study setting(s)

Hospital

#### Study type(s)

Prevention

#### Participant information sheet

Not available in web format, please use the contact details to request a patient information sheet

#### Health condition(s) or problem(s) studied

Overweight patients

#### Interventions

Participants are randomized by sealed envelope block method:

1. Study medication (VERUM) capsule composition: pomegranate extract 163 mg, chitin-glucan

375 mg, siliciumdioxide 18 mg, magnesium stearate 18 mg

2. Placebo capsule composition: cellulose 500mg, silicium dioxide 18 mg, magnesium stearate 18 mg.

Medication is administered per oral twice a day, 2 capsules each time (4 capsules per day). Duration of treatment: 3 months.

Duration of the study - 4 months with 5 visits performed: V0 - at the beginning of the study, V1 - at the end of the first month of treatment, V2 - at the end of the second month of treatment, V3 - at the end of third (last) month of treatment, V4 - one month after the treatment has finished.

#### Intervention Type

Supplement

#### Primary outcome measure

- 1. Blood collection with blood analysis at V0, V1, V3, V4:
- 1.1. Inflammatory parameters at V0, V1, V3, V4:
- 1.1.1. C-reactive protein high-sensitivity (hsCRP), µg/ml, measured using turbidimetric method
- 1.1.2. Interleukin 6, pg/ml, measured using ELISA
- 1.1.3. Interleukin 8, pg/ml , measured using ELISA
- 1.1.4. Interleukin 10, µmol/l, measured using ELISA
- 1.1.5. Tumor necrosis factor α, pg/ml, measured using ELISA
- 1.1.6. Monocyte chemotactic protein 1, MCP1, pg/ml, measured using ELISA

- 1.2 Parameters of oxidative and endothelial function at V0, V1, V3, V4:
- 1.2.1. Vitamin A, mg/L, measured using UPLC
- 1.2.2. Vitamin E, mg/L, measured using UPLC
- 1.2.3. Vitamin C, mg/l, measured using HPLC
- 1.2.4. Coenzyme Q10, μg/l, measured using HPLC
- 1.2.5  $\beta$ -carotene, ng/ml, measured using HPLC
- 1.2.6. Superoxide dismutase (SOD), U/ml, measured using enzyme assay
- 1.2.7. Glutathione peroxidase (GPX), U/ml, measured using spectrophotometry
- 1.2.8. Soluble vascular adhesion molecule 1 (sVCAM 1), ng/ml, measured using ELISA
- 1.2.9. Soluble intercellular adhesion molecule 1 (sICAM 1), ng/ml, measured using ELISA
- 1.2.10. Asymmetric dimethylarginine (ADMA), µmol/l, measured using ELISA
- 1.2.11. Cluster of differentiation 40 ligand (CD40-L), pg/ml, measured using ELISA
- 1.3. Lipid parameters at V0, V1, V3, V4:
- 1.3.1. Cholesterol, mg/dl, measured using reflectometry
- 1.3.2. High density lipoprotein (HDL), mg/dl, measured using reflectometry
- 1.3.3. Low density lipoprotein (LDL), mg/dl, measured using spectrophotometry
- 1.3.4. Triglycerides, mg/dl, measured using reflectometry
- 1.3.5. Lipoprotein a (LPa), g/l, measured using nephelometry
- 1.3.6. Apolipoprotein A1 (ApoA1), g/l, measured using nephelometry
- 1.3.7. Apolipoprotein B (ApoB), g/l, measured using nephelometry
- 1.3.8. Small density low density lipoprotein (Sd-LDL), nmol/ml, measured using ELISA
- 1.3.9. Anti-oxidized low density lipoprotein antibodies (anti ox-LDL ab), measured using ELISA

2. Endothelial function assessed by ultrasound examination of the brachial artery at V0, V1, V2, V3, V4:

- 2.1. Diameter baseline (D baseline), mm
- 2.2. Diameter maximum (D max), mm
- 2.3. Flow mediated dilatation (FMD), %

### Secondary outcome measures

- 1. Blood collection with confounding parameters analysis at V0, V1, V3, V4:
- 1.1. Glycaemia, mg/dl, measured using reflectometry
- 1.2. Glycated hemoglobin A1c (HbA1c), %, measured using chromatography
- 1.3. Glutamic oxaloacetic transaminase (GOT), UI/L, measured using enzyme assay
- 1.4. Glutamic-pyruvic transaminase (GPT), UI/L, measured using enzyme assay
- 1.5. Gamma-glutamic transaminase (GGT), UI/L, measured using enzyme assay
- 1.6. Iron, μg/dl, measured using reflectometry
- 1.7. Albumin (ALB), g/l, measured using reflectometry
- 1.8. Ferritin, µg/l, measured using immunoassay
- 1.9. Magnesium, mmol/l, measured using reflectometry
- 1.10. Uric acid, mg/dl, measured using reflectometry
- 1.11 Creatinine (CRS), mg/dl, measured using enzyme assay

2. Collection of feces for intestinal microbiota analysis (V0, V3) during randomization and at the end of treatment

3. Assessment of nutritional status (via questionnaire) and bioelectrical impedance analysis at V0, V1, V2, V3, V4:

- 3.1. Height, m
- 3.2. Weight, kg
- 3.3. Waist circumference, cm
- 3.4. Abdominal diameter, cm
- 3.5. Fat, %

3.6. Fat, kg 3.7. Body mass index (BMI) 3.8. Waist/hip ratio

Overall study start date 01/05/2017

**Completion date** 

30/05/2019

## Eligibility

### Key inclusion criteria

- 1. Male and female
- 2. Aged at least 18 years
- 3. Caucasian type
- 4. 25<BMI <35 ± 5%

5. Stable patient's weight  $\pm$  5% for at least 3 last months and patient's agreement not to change food habits during the study

- 6. Menopausal women without or with HRT with a stable rate for at least 3 months
- 7. Able to understand and to give consent

8. Compliant with treatment

Participant type(s)

Mixed

## Age group

Adult

### Lower age limit

18 Years

Sex

Both

**Target number of participants** 100

### Key exclusion criteria

- 1. Current cancer treatment
- 2. Diabetes
- 3. Severe intestinal problems
- 4. Autoimmune or chronic inflammatory diseases
- 5. Unintentional weight loss (> 10% of body weight) during the previous 6 months
- 6. Particular nutritional practice (vegetarians, vegans, high consumption of fiber)
- 7. Intake of food supplements
- 8. History of eating disorders (anorexia, bulimia)
- 9. Dieting at the time of inclusion
- 10. Underwent bariatric surgery
- 11. Consumption in the preceding 6 weeks of antibiotics or probiotics

12. Pregnancy, breastfeeding

13. Active smoker or ex-smoker for less than 2 years

14. Food allergy to one of the components of the supplement

15. Participation in another clinical study at the time of inclusion

16. Usually taking medication: anti-inflammatory drugs, corticosteroids, antiepileptics, neuroleptics, laxatives, antibiotics

### Date of first enrolment

01/10/2017

## Date of final enrolment

31/12/2018

## Locations

**Countries of recruitment** Belgium

## Study participating centre

CHU UCL Namur Godinne

Avenue du Dr Gaston Therasse 1 Yvoir Belgium 5530

## Sponsor information

## Organisation

KitoZyme

### Sponsor details

Rue de Milmort, 680 Herstal Belgium 4040 +32 (0)42 59 85 00 info@kitozyme.com

#### Sponsor type Industry

Website http://kitozyme.com/

## ROR

## https://ror.org/008a8ch32

## Funder(s)

**Funder type** Other

**Funder Name** Competitivity pole WAGRALIM (Public service of Wallonia region)

## **Results and Publications**

#### Publication and dissemination plan

Planned publication in a high-impact peer reviewed journal. Additional documents (study protocol, statistical analysis plan, informed consent form, clinical study report, analytic code) could be available for sharing with researchers or investigators beginning 3 months and ending 36 months following article publication (also after obtaining all participant's permission).

#### Intention to publish date

30/12/2019

#### Individual participant data (IPD) sharing plan

The ADIPOSTOP study represents a part of the multidisciplinary project containing several participants. Patient level data can be available only after obtaining data sharing permission from all the participants.

#### IPD sharing plan summary

Other