Pressure Support Ventilation (PSV) or Synchronised Intermittent Mandatory Ventilation (SIMV) for weaning preterm infants on mechanical ventilation

Submission date 23/02/2011	Recruitment status No longer recruiting	 Prospectively registered Protocol
Registration date 08/03/2011	Overall study status Completed	 Statistical analysis plan Results
Last Edited 20/11/2017	Condition category Respiratory	Individual participant dataRecord updated in last year

Plain English summary of protocol

Background and study aims

Most premature infants need breathing support through a tube in their wind pipe. We know providing artificial breathing support can lead to long-term problems with the growth of the lungs. Although most clinicians would agree when to start breathing support for a preterm infant, there is as yet no consensus on when to reduce this support, also called 'weaning on mechanical ventilation'. This study is investigating two different methods of 'weaning' breathing support in premature infants. Until recently, most units in the UK used synchronised intermittent mandatory ventilation (SIMV). SIMV matches a fixed number of breaths on the breathing support machine and over a period of time the fixed number of breaths is reduced before the breathing tube is taken out (extubation). However, advances in technology have led to newer modes of ventilation. One such mode is pressure support ventilation (PSV). In PSV the infant is in charge of all aspects of breathing. The medical team would only set the permissible pressure level to support the spontaneous breaths, so weaning on mechanical ventilation would only be to decrease the pressure support before the tube is taken out. In adults PSV is preferred as this is more comfortable.

Who can participate?

Preterm infants born between 23+0 weeks and 32 weeks gestation who were ventilated for at least 6 hours for Respiratory Distress Syndrome (RDS).

What does the study involve?

Participating infants are randomly allocated to one of the two modes of 'weaning' breathing support. It is usual for some infants to not tolerate the reduction in breathing support and 'fail'. If at any stage the infant does not tolerate any of the weaning methods then the medical team assess them as per usual policy. All other aspects of their care are provided as per unit policy and current practice.

What are the possible benefits and risks of participating? At present there is no perceived benefit or risk of one method over the other. However, at the end of the study both the groups would be compared to find out if there is a difference.

Where is the study run from? University Hospital of North Tees and James Cook University Hospital (UK)

When is the study starting and how long is it expected to run for? January 2010 to May 2013

Who is funding the study? University Hospital of North Tees (UK)

Who is the main contact? Prof. Samir Gupta samir.gupta@nth.nhs.uk

Contact information

Type(s) Scientific

Contact name Dr Samir Gupta

Contact details

Neonatal Unit University Hospital of North Tees Hardwick Road Stockton United Kingdom TS19 8PE +44 (0)16 4262 4250 samir.gupta@nth.nhs.uk

Type(s)

Scientific

Contact name Dr Prashant Mallya

Contact details

Neonatal Unit University Hospital of North Tees Hardwick Road Stockton United Kingdom TS19 8PE + 44 (0)16 4262 4248 mallyaprashant@yahoo.com

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers Version 13, 29/12/2010

Study information

Scientific Title

Pressure Support Ventilation (PSV) or Synchronised Intermittent Mandatory Ventilation (SIMV) for weaning preterm infants on mechanical ventilation: a multi-centre randomised controlled trial

Acronym POST-UK Study

Study objectives

The trial is investigating the practicability of PSV as opposed to SIMV for weaning on mechanical ventilation in preterm infants needing mechanical ventilatory support for Respiratory Distress Syndrome (RDS).

The primary endpoint of the trial is defined as time from entry into the trial (predefined priori of mean airway pressure of less than 10cm, oxygen requirement of less than 40% and spontaneous breath rates of at least 50%) to the time when the infant is ready for extubation demonstrated by passing then minute ventilation test (MVT).

The null hypothesis for the trial would find no difference between the two modes. However any difference between the two modes would be calculated using appropriate statistical calculations.

The study is planned across two tertiary care units in the North East of England.

Ethics approval required Old ethics approval format

Ethics approval(s) Newcastle and North Tyneside Research Ethics Committee, 10/01/2011

Study design Multi-centre randomised controlled trial

Primary study design Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Not available in web format, please use the contact details to request a patient information sheet

Health condition(s) or problem(s) studied

Respiratory distress syndrome in preterm babies

Interventions

The intervention arm would receive pressure supported mechanical breaths for all the spontaneous breaths generated by the infant and the control arm would receive SIMV mode of respiratory support as is currently practiced.

Intervention Type

Procedure/Surgery

Primary outcome measure

The primary endpoint is duration of weaning on mechanical ventilation - defined as time from commencing weaning in the assigned mode of respiratory support to the time when the infant is ready for extuabtion demonstrated by passing the MVT

Secondary outcome measures

- 1. Total duration of mechanical support through the endotracheal (ET) tube
- 2. Total duration of respiratory support including non invasive respiratory support
- 3. Discharge on home oxygen
- 4. Complications of prematurity

Overall study start date 26/01/2011

Completion date 26/01/2013

Eligibility

Key inclusion criteria

Preterm infants with RDS, stratified into three groups (23+0 to 25+6, 26+0 to 28+6 and 29+0 to 31+6 weeks) based on their gestation at birth would be enrolled if they are needing mechanical ventilation through an endotracheal tube for at least 6 hours
 Signed parental consent is obtained before randomisation

Participant type(s) Patient

Age group

Neonate

Sex Both

Target number of participants 110

Key exclusion criteria

1. Severe congenital malformation

2. Neuromuscular disorder demonstrated clinically with decreased muscular tone

3. Upper airway anomaly

4. Infants transferred from other unit who have met the study entry criteria on admission to the neonatal unit

Date of first enrolment

26/01/2011

Date of final enrolment 26/01/2013

Locations

Countries of recruitment England

United Kingdom

Study participating centre University Hospital of North Tees Stockton United Kingdom TS19 8PE

Sponsor information

Organisation North Tees and Hartlepool NHS Foundation Trust (UK)

Sponsor details Univeristy Hospital of North Tees Hardwick Road Stockton England United Kingdom TS19 8PE +44 (0)16 4238 3251 jane.greenaway@nth.nhs.uk

Sponsor type Hospital/treatment centre

ROR https://ror.org/04zzrht05

Funder(s)

Funder type Hospital/treatment centre

Funder Name University Hospital of North Tees (UK)

Results and Publications

Publication and dissemination plan

Data analysis is ongoing. All aspects of the study would be sent for publication including the study protocol. The researchers aim to disseminate the research findings in 2018.

Intention to publish date

20/05/2018

Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study will be stored in a repository.

IPD sharing plan summary

Stored in repository