

The effect of hyaluronidase on dispersal of local anaesthetic solution after sub-Tenon's injection

Submission date 30/09/2005	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 30/09/2005	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 11/12/2008	Condition category Surgery	<input type="checkbox"/> Individual participant data

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Ms M Khandwala

Contact details
Leeds Metropolitan University
48 Riding House Street
Leeds
United Kingdom
LS9 7TF
+44 (0)113 283 2600

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers
N0436146624

Study information

Scientific Title

Study objectives

To study the difference in distribution of anaesthetic fluid with and without hyaluronidase after sub-Tenon's block using B scan ultrasonography. Patients will be randomised to a group receiving local anaesthetic plus hyaluronidase. An ultrasound scan of the orbit will be performed prior to injection and 1, 3 and 5 minutes after injection to assess dispersal of local anaesthetic solution. The maximum depth of the local anaesthetic solution will be used to compare groups. The quality of the block will also be assessed in terms of akinesia, patient pre-operative pain scores and surgical assessment of the block.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Not provided at time of registration

Study design

Randomised controlled trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Not specified

Study type(s)

Not Specified

Participant information sheet

Health condition(s) or problem(s) studied

Surgery: Anaesthesia

Interventions

1. Anaesthetic fluid with hyaluronidase
2. Anaesthetic fluid without hyaluronidase

Intervention Type

Procedure/Surgery

Phase

Not Specified

Primary outcome measure

Ultrasound measured depth of local anaesthetic solution behind the eye at 1, 3 and 5 minutes after the injection.

Secondary outcome measures

Akinesia, pain scores and surgical satisfaction.

Overall study start date

24/03/2004

Completion date

15/06/2004

Eligibility

Key inclusion criteria

Consecutive patients presenting for routine cataract extraction surgery will be approached to enrol in the study. They are mostly elderly patients (from previous studies in Leeds age range 65-80 years)

Participant type(s)

Patient

Age group

Senior

Sex

Not Specified

Target number of participants

Added December 2008: 19

Key exclusion criteria

1. Allergy to local anaesthetic
2. Inability/refusal to give informed consent
3. Any patient with pre existing extra ocular muscle palsy, patients with single eye who could be unable to follow a target so assessment of akinesia would be impossible.

Date of first enrolment

24/03/2004

Date of final enrolment

15/06/2004

Locations

Countries of recruitment

England

United Kingdom

Study participating centre
Leeds Metropolitan University
Leeds
United Kingdom
LS9 7TF

Sponsor information

Organisation
Department of Health

Sponsor details
Richmond House
79 Whitehall
London
United Kingdom
SW1A 2NL
+44 (0)20 7307 2622
dhmail@doh.gsi.org.uk

Sponsor type
Government

Website
<http://www.dh.gov.uk/Home/fs/en>

Funder(s)

Funder type
Government

Funder Name
Leeds Teaching Hospitals NHS Trust (UK), own account

Results and Publications

Publication and dissemination plan
Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/08/2008		Yes	No