

The improvement effect of probiotics using tablet containing *Lactobacillus salivarius* WB21 on bad breath

Submission date 27/12/2012	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 10/01/2013	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 17/12/2020	Condition category Signs and Symptoms	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

Oral malodor (bad breath) is primarily the result of the breakdown of amino acids in the local debris of the oral cavity by the micro organisms. Because probiotics possess potential capacity for improving microbiota, it may improve oral malodor. *Lactobacillus salivarius* WB21 has been recognized as a probiotic for oral health, and we previously reported that oral administration of tablets containing *L. salivarius* WB21 improved oral malodor in a study. The aim in this study was to confirm the effect of probiotics using *L. salivarius* WB21 on bad breath.

Who can participate?

Eighty-two patients complaining of halitosis (bad breath) were assessed for study eligibility at the Oral Malodor Clinic of Fukuoka Dental College Medical and Dental Hospital, Japan between June 2010 and September 2011. The eligible participants were 26 patients (22 females and four males, age range, 22-67 years). They were patients with oral malodor above a questionable level, not currently visiting a dentist for treatment, no acute symptoms requiring immediate oral cavity treatment, no use of probiotic supplements, no antibiotic use within the last month, no daily smoking habit, no systemic illness and no adverse reactions to lactose or fermented milk products.

What does the study involve?

The intervention (treatment) compared test tablets with placebo. All participants received the same treatment that ingested both tablets containing *L. salivarius* WB21 and placebo. Placebo samples contained only xylitol.

What are the possible benefits and risks of participating?

The participants have a chance of improving oral malodor by oral consumption of test tablets. There is not any side effect.

Where is the study run from?

The study took place at the Oral Malodor Clinic of Fukuoka Dental College Medical and Dental Hospital, Japan.

When is the study starting and how long is it expected to run for?
The participants were recruited between June 2010 and September 2011.

Who is funding the study?
This study was supported in part by a Grant-in-Aid for Young Scientists (no. 23792532), Grant-in-Aid for Scientific Research (no. 23593078) and a Grant-in-Aid for Advanced Science Research from the Ministry of Education, Culture, Sports, Science, and Technology, Japan.

Who is the main contact?
Nao Suzuki
naojasz@college.fdcnet.ac.jp

Contact information

Type(s)
Scientific

Contact name
Dr Nao Suzuki

Contact details
2-15-1, Tamura Sawara-ku
Fukuoka
Japan
814-0193
-
naojasz@college.fdcnet.ac.jp

Additional identifiers

Protocol serial number
N/A

Study information

Scientific Title
A double-blind, randomized, placebo-controlled crossover trial of a Lactobacillus salivarius WB21-containing tablet for treatment of oral malodor

Study objectives
Oral malodor is a common problem in humans. Most oral malodor originates directly from the oral cavity owing to conditions such as periodontitis, tongue debris, poor oral hygiene, deep caries, inadequately fitted restorations, and endodontic lesions. Oral malodor is primarily the result of microbial metabolism of amino acids in local debris in the oral cavity. The most common compounds associated with oral malodor are volatile sulfur compounds (VSCs), such as hydrogen sulfide (H₂S) and methyl mercaptan (CH₃SH).

Lactobacillus salivarius WB21 has been recognized as a probiotic for oral health, and we previously reported that oral administration of tablets containing L. salivarius WB21 improved oral malodor in an open trial. In this study, we conducted a 14-day, double-blind, placebo-

controlled, randomized crossover trial of tablets containing *L. salivarius* WB21 or placebo taken orally by patients having oral malodor to confirm the authenticity of the effect of this organism on oral malodor.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Ethics Committee for Clinical Research of Fukuoka Dental College and Fukuoka College of Health Sciences, July 2, 2008, approval number 125

Study design

Randomized double-blind crossover placebo-controlled clinical trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Oral malodor

Interventions

The tablets (MINNA NO ZENDAMAKIN WB21 TABLET®; Wakamoto Pharmaceutical Co., Tokyo, Japan) contained 6.7×10^8 colony-forming units (CFU) of *L. salivarius* WB21 and 280-mg xylitol per tablet.

Placebo samples contained only xylitol (280 mg/tablet).

The dose throughout the test period was maintained at three tablets per day, taken orally after eating. Subjects were directed to place a tablet on the tongue for a few minutes and allow it dissolve.

The period that the patients had tablets was four weeks, 2 weeks for test tablets and 2 weeks for placebo tablets with a wash out period of 2 weeks.

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Lactobacillus salivarius WB21

Primary outcome(s)

1. Organoleptic (OLT) score: The OLT scores were estimated by two of the three evaluators (with training and experience in calibration tests) using a scale of 0 to 5 (0, absence of oral malodor; 1, questionable odor; 2, slight malodor; 3, moderate malodor; 4, strong malodor; 5, severe malodor), and the mean of the scores given by the evaluators was used.

2. Total volatile sulfur compounds concentration: The total VSCs condition was measured by gas chromatography
3. Periodontal health: Periodontal health was assessed using the average probing pocket depth (PPD) and the number of bleeding on probing (BOP) sites. PPD and BOP were measured at six points around each tooth in all subjects.
4. Plaque control was evaluated using the Silness L  e Plaque Index
5. Degree of tongue coating was determined by tongue coating score (TCS): 0, no tongue coating; 1, thin tongue coating covering less than one-third of the tongue dorsum; 2, thick tongue coating covering approximately one-third of the tongue dorsum or thin tongue coating covering one-third to two-thirds of the tongue dorsum; 3, thick tongue coating covering one-third to two-thirds of the tongue dorsum or thin tongue coating covering more than two-thirds of the tongue dorsum; and 4, thick tongue coating more than two-thirds of the tongue dorsum.
6. Volume of stimulated salivary flow was measured using the chewing gum test. The patient was asked to pool saliva in the oral cavity and spit into a vessel each minute throughout a 5-min collection period.

Key secondary outcome(s)

1. Volatile sulfur compound concentration
2. The number of each probing depth
3. The number of oral bacteria in the saliva

Completion date

30/09/2011

Eligibility

Key inclusion criteria

1. 26 eligible patients (22 females and four males; mean age, 43.5 \pm 11.5 years; age range, 22-67 years)
2. Oral malodor above a questionable level (OLT score \geq 1.5)
3. Not currently visiting a dentist for treatment
4. No acute symptoms requiring immediate oral cavity treatment
5. No use of probiotic supplements
6. No antibiotic use within the last month
7. No daily smoking habit
8. No systemic illness
9. No adverse reactions to lactose or fermented milk products

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Sex

All

Key exclusion criteria

Does not meet inclusion criteria

56 were excluded: 48 did not meet the inclusion criteria, three refused to participate, and five had personality traits that were considered difficult for adherence to the protocol.

Date of first enrolment

01/06/2010

Date of final enrolment

30/09/2011

Locations

Countries of recruitment

Japan

Study participating centre

2-15-1, Tamura Sawara-ku

Fukuoka

Japan

814-0193

Sponsor information

Organisation

Fukuoka Dental College (Japan)

ROR

<https://ror.org/04zkc6t29>

Funder(s)

Funder type

Government

Funder Name

Ministry of Education, Culture, Sports, Science, and Technology (Japan) - Grant-in-Aid for Young Scientists (no. 23792532), Grant-in-Aid for Scientific Research (no. 23593078) and a Grant-in-Aid for Advanced Science

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/04/2014	17/12/2020	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes