Neoadjuvant short-term radiotherapy followed by surgery versus surgery alone for patients over 70 years of age with local advanced rectal cancer and who are medically unfit for standard neoadjuvant radiochemotherapy treatment

Recruitment status	[X] Prospectively registered
06/06/2012 No longer recruiting	☐ Protocol
Overall study status	Statistical analysis plan
Completed	Results
Condition category	Individual participant data
	Record updated in last year
	No longer recruiting Overall study status

Plain English summary of protocol

Background and study aims

Rectal cancer is cancer that starts in the rectum, the lower part of the large bowel. Locally advanced rectal cancer means that the cancer has spread into the tissues around the rectum but hasn't spread to other organs. The aim of this study is to find out whether it is better to treat patients aged over 70 with radiotherapy before they undergo surgery to remove the cancer.

Who can participate?

Patients aged over 70 with locally advanced rectal cancer who are medically unfit for standard radiochemotherapy treatment

What does the study involve?

Participants are randomly allocated to one of two groups. One group are treated with radiotherapy followed by surgery followed by chemotherapy if tolerated. The other group are treated with surgery followed by chemotherapy. Both groups are followed up to see which treatment has better results.

What are the possible benefits and risks of participating? Not provided at time of registration

Where is the study run from?
South City Hospital Rostock (Klinikum Suedstadt Rostock) (Germany)

When is the study starting and how long is it expected to run for? January 2013 to January 2021

Who is funding the study?
South City Hospital Rostock (Klinikum Suedstadt Rostock) (Germany)

Who is the main contact? Prof. Dr med. habil. Kaja Ludwig kfch@kliniksued-rostock.de

Contact information

Type(s)

Scientific

Contact name

Prof Kaja Ludwig

Contact details

Department of Surgery Klinikum Suedstadt Rostock Suedring 81 Rostock Germany 18059

_

kfch@kliniksued-rostock.de

Additional identifiers

Clinical Trials Information System (CTIS)

2007-006758-24

Protocol serial number

N/A

Study information

Scientific Title

NEROP-RC (NEoadjuvant Radiotherapy for Older Patients Rectal Cancer): a randomised controlled phase III study

Acronym

NEROP-RC

Study objectives

The primary aim of the study is to demonstrate the superiority of preoperative radiotherapy (5x5Gy) followed by total mesorectal excision (TME) or abdomino-sacral rectum extirpation in patients over 70 years of age with locally advanced rectal cancer, up to a max. of uT1-3N1-2M0 versus surgery alone with TME or abdomino-sacral rectum extirpation with regard to locoregional control.

Ethics approval required

Old ethics approval format

Ethics approval(s)

University of Rostock Ethics Commission, 31/05/2010, ref: FK-2010-0008

Study design

Prospective randomised controlled multicentre phase III study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Optimal therapy for patients > 70 years with rectal cancer

Interventions

All patients over 70 years of age and having preoperatively diagnosed locally advanced rectal cancer (uT1-3 and/or uN1-2), for whom the neoadjuvant standard RCT appears to present too high a risk, are randomised to two study arms:

Arm A: neoadjuvant short-term RT (5x5Gy) followed by TME or abdominosacral rectum extirpation

Arm B (control arm): TME alone or abdomino-sacral rectum extirpation.

Four weeks following surgery re-evaluation takes place in both arms if possible, then administration of 5-FU- based Chemotherapy (CT) for 6 months. If no CT is possible, then control arm is followed up.

Intervention Type

Mixed

Primary outcome(s)

Superiority of preoperative radiotherapy (5x5Gy) followed by TME and abdomino-sacral rectum extirpation in patients over 70 years of age with locally advanced rectal cancer up to a maximum of uT1-4N1-2M0 versus surgery alone with TME or abdomino-sacralen rektum extirpation with respect to locoregional control.

Key secondary outcome(s))

- 1. Disease-free survival (DFS) (first case of tumor illness locoregional or distant metastases)
- 2. Occurrence of distant metastases
- 3. Frequency of other, non-tumor-related deaths
- 4. Determination for comorbidity based on the Charlson Index and the ADL Index
- 5. Determination of how often chemotherapy can be performed postoperatively
- 7. Acute and long-term toxicity of preoperative radiotherapy
- 8. Overall survival (OS)

Completion date

01/01/2021

Eligibility

Key inclusion criteria

- 1. Patients with primary diagnosis of histologically bioptically secured diagnosis of rectal cancer
- 2. Rectal cancer at any height (up to 16cm lower edge of tumor measured from the anal line using inflexible rectoscopy)
- 3. Decision made by the treating tumor team that intensified RCT is not possible
- 4. Preoperative tumor staging max. uT1-4uN1-2M0
- 5. Age >/= 70 years
- 6. Karnofsky index >/= 70%
- 7. ASA-classification < IV
- 8. Oral and written consent according to
- 9. Good Clinical Practice and corresponding local, national and international regulations

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Senior

Sex

Αll

Key exclusion criteria

- 1. Other previous cancer illnesses with the exception of appropriately treated in situ carcinoma of the cervix or skin tumors with no indication of melanoma or in cases of tumor-free state ≥ 10 years subsequent to appearance of cancer which has undergone appropriate treatment
- 2. Participation in studies elsewhere which include chemotherapy or radiotherapy
- 3. Inability to participate in regular check-ups

Date of first enrolment

01/01/2013

Date of final enrolment

01/01/2021

Locations

Countries of recruitment

Germany

Study participating centre

Klinikum Suedstadt Rostock

Rostock Germany 18059

Sponsor information

Organisation

South City Hospital Rostock (Klinikum Suedstadt Rostock) (Germany)

ROR

https://ror.org/02m0p4y77

Funder(s)

Funder type

Hospital/treatment centre

Funder Name

South City Hospital Rostock (Klinikum Suedstadt Rostock), Department of Surgery (Germany)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type Details Date created Date added Peer reviewed? Patient-facing?

Participant information sheet 11/11/2025 No Yes