

# Can Teriparatide prevent the onset of hypocalcemia after surgical thyroid removal?

<b>Submission date</b> 09/04/2016	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered
		<input type="checkbox"/> Protocol
<b>Registration date</b> 12/04/2016	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan
		<input checked="" type="checkbox"/> Results
<b>Last Edited</b> 08/05/2017	<b>Condition category</b> Nutritional, Metabolic, Endocrine	<input type="checkbox"/> Individual participant data

## Plain English summary of protocol

### Background and study aims

Hypoparathyroidism is a hormonal disorder in which the parathyroid glands in the neck do not produce enough parathyroid hormone. Parathyroid hormone (PTH) is responsible for regulating the amount of calcium in the blood, by acting on the bones, kidneys and intestines. If the body does not produce enough PTH then blood calcium levels fall (hypocalcaemia), leading to unusual muscle movements (such as jerking or twitching), a tingling sensation in the fingers and toes, muscle cramps, mental problems such as confusion and irritability, and even heart problems. The most common cause of hypoparathyroidism is unintentional or unavoidable damage to or removal of the parathyroid glands during surgery. Predicting the risk of a surgical patient developing hypoparathyroidism is an important part of care. It has been found that measuring intact parathyroid hormone (iPTH) during or shortly after surgery is a good method of predicting risk. Teriparatide (PTH 1-34) is drug containing a man-made form of parathyroid hormone. A recent study has shown that it able to correct blood calcium levels in patients with hypocalcemia following surgery. The aim of this study is to find out whether teriparatide is able to prevent post-surgical hypocalcemia in patients who have a high risk of developing hypocalcemia after thyroid surgery.

### Who can participate?

Adults who have low iPTH levels after having part or all of the thyroid gland removed (thyroidectomy).

### What does the study involve?

Participants are randomly allocated to one of two groups. Participants in the first group receive 20 micrograms of teriparatide from a trained nurse using an injection pen after surgery. This is then repeated every 12 hours until they are discharged from hospital. Participants in the second group do not receive any treatment and are observed throughout their hospital stay. Every day they are in hospital after their surgery (at least two days), participants in both groups have a sample of blood taken so that their iPTH levels can be measured. The length of time participants from each group spend in hospital is also compared.

### What are the possible benefits and risks of participating?

Participants who receive treatment may benefit from improved PTH levels, avoiding

hypocalcemia. There are no notable risks of participating, although blood testing may cause pain and bruising to some participants.

Where is the study run from?

Università Campus Bio-Medico di Roma, Surgical Ward (Italy)

When is the study starting and how long is it expected to run for?

January 2016 to May 2016

Who is funding the study?

Università Campus Bio-Medico di Roma (Italy)

Who is the main contact?

Dr Andrea Palermo

a.palermo@unicampus.it

## Contact information

**Type(s)**

Scientific

**Contact name**

Dr Andrea Palermo

**Contact details**

Università Campus Bio-Medico di Roma

via Alvaro del Portillo 5

Rome

Italy

00128

## Additional identifiers

**EudraCT/CTIS number**

2016-000481-32

**IRAS number**

**ClinicalTrials.gov number**

**Secondary identifying numbers**

N/A

## Study information

**Scientific Title**

PTH(1-34) for the primary prevention of post-surgical hypocalcemia: the THYPOS trial

**Acronym**

THYPOS

**Study objectives**

The aim of this study is to evaluate whether teriparatide is able to prevent post-surgical hypocalcemia in subjects with high risk of hypocalcemia after thyroid surgery.

**Ethics approval required**

Old ethics approval format

**Ethics approval(s)**

Ethic Committee Università Campus-Biomedico (Rome), 01/02/2016, ref: 16.16

**Study design**

Prospective phase II single-centre open label randomized controlled trial

**Primary study design**

Interventional

**Secondary study design**

Randomised controlled trial

**Study setting(s)**

Hospital

**Study type(s)**

Prevention

**Participant information sheet**

Please see additional files

**Health condition(s) or problem(s) studied**

Hypocalcemia due to Post-surgical hypoparathyroidism

**Interventions**

Subjects with iPTH levels at 4 hours after thyroidectomy  $< 10$  pg/ml will be enrolled. After surgery, they will be randomized (1:1) to receive treatment with teriparatide or following the standard clinical care.

Treatment group: A nurse administers 20 mcg of teriparatide using an injection pen (Forsteo® Eli Lilly Nederland B.V.) to patients immediately after randomisation. Following this, administrations will be done every 12 hours until discharge.

Control group: Participants will receive no treatment and they will be accurately observed until the discharge as suggested by national and international guidelines.

At 8.00 AM on the first and second postoperative, in fasting state, a blood sample will be drawn and calcium, phosphate, albumin, magnesium, will be measured. A clinical evaluation and ECG will be performed to exclude signs or symptoms of hypocalcemia.

**Intervention Type**

Drug

**Phase**

Phase II

**Drug/device/biological/vaccine name(s)**

Teriparatide (Forsteo®)

**Primary outcome measure**

1. Serum calcium is measured using hematological testing at baseline (4 hours after thyroidectomy), in the morning (from 8:00 to 8:30 AM) of postoperative day 1 and 2, and daily until the discharge (if hospitalisation lasts more than two days)
2. Signs and symptoms of hypocalcemia are determined through physical examinations at baseline (4 hours after thyroidectomy) and daily until discharge.

**Secondary outcome measures**

Duration of hospitalization is measured counting the days from the thyroidectomy until the discharge.

**Overall study start date**

04/01/2016

**Completion date**

10/05/2016

**Eligibility****Key inclusion criteria**

1. Aged 18 and over
2. Patients with a formal surgical indication for thyroidectomy.

**Participant type(s)**

Patient

**Age group**

Adult

**Lower age limit**

18 Years

**Sex**

Both

**Target number of participants**

11 patients for each group ( total number: 22). Estimating a dropout rate of 20%: total number: 26)

**Key exclusion criteria**

1. Aged under 18 years
2. Pregnancy
3. Renal failure (glomerular filtration rate < 30 mL/min)
4. Hypersensitivity to the active substance or excipients

5. Any prior parathyroid pathology
6. Preexisting hypercalcemia
7. Metabolic bone disease other than osteoporosis
8. Ongoing therapy for osteoporosis
9. In the last 6 months administration of calcitonin
10. Systemic corticosteroids, estrogens, raloxifene, fluoride, lithium, loop or thiazide diuretics, aromatase inhibitors or other drugs that could interfere with calcium metabolism
11. History of skeletal malignancies (primary or metastatic)
12. Active or recent urolithiasis
13. Unexplained elevation of serum alkaline phosphatase levels
14. Prior radiation therapy involving the skeleton, serum magnesium levels below the lower limits or above the upper limits of normal

**Date of first enrolment**

01/02/2016

**Date of final enrolment**

30/04/2016

## **Locations**

**Countries of recruitment**

Italy

**Study participating centre**

**University Campus Biomedico of Rome**

Surgical ward

via Alvaro del Portillo 20

Rome

Italy

00128

## **Sponsor information**

**Organisation**

Università Campus Bio-Medico di Roma

**Sponsor details**

Department of Endocrinology and Diabetes

Via Alvaro del Portillo 20

Rome

Italy

00128

**Sponsor type**

University/education

**ROR**

<https://ror.org/04gqx4x78>

## Funder(s)

**Funder type**

University/education

**Funder Name**

Università Campus Bio-Medico di Roma

**Alternative Name(s)**

Campus Bio-Medico University

**Funding Body Type**

Private sector organisation

**Funding Body Subtype**

Universities (academic only)

**Location**

Italy

## Results and Publications

**Publication and dissemination plan**

Planned publication in the journals JCEM, JBMR or Bone.

**Intention to publish date**

31/12/2016

**Individual participant data (IPD) sharing plan**

**IPD sharing plan summary**

Available on request

**Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Participant information sheet</a>			13/04/2016	No	Yes
<a href="#">Results article</a>	results	01/11/2016		Yes	No