# The effect of breakfast consumption on physical activity in girls

Submission date	Recruitment status	<ul><li>Prospectively registered</li></ul>
17/05/2017	No longer recruiting	Protocol
Registration date	Overall study status	Statistical analysis plan
23/05/2017	Completed	[X] Results
Last Edited	Condition category	Individual participant data
07/03/2018	Nutritional, Metabolic, Endocrine	

### Plain English summary of protocol

Background and study aims

Many children and adolescents often skip breakfast and do not engage in the recommended amounts of physical activity. Research has shown that breakfast consumption frequency (i.e., days per week that breakfast is consumed) is linked with increased physical activity in young people. However, these results have not been consistent when using an accelerometer (activity monitor) to measure physical activity. It is also not known whether increased breakfast consumption frequency causes an increase in physical activity, or whether the link is simply because those who eat breakfast generally engage in a range of healthy lifestyle habits. Although some studies in adults have indicated that breakfast consumption can increase physical activity, others have reported no effect, and no study has examined the effects of breakfast consumption frequency on physical activity in children or adolescents. This is particularly important for adolescent girls, who frequently skip breakfast and have low physical activity levels. Therefore, the aim of this study is to examine the effect of daily compared with intermittent breakfast consumption on physical activity in adolescent girls.

Who can participate? Healthy girls aged 11 to 13

### What does the study involve?

The study consists of two periods that last 7 days each: daily breakfast consumption and intermittent breakfast consumption. Daily breakfast consumption involves eating a standard breakfast every day for 7 days. Intermittent breakfast consumption involves eating breakfast on three intermittent days (i.e., days 2, 4 and 6) with four days without breakfast (i.e., days 1, 3, 5 and 7). There is a 7-10 day break in between the two periods. The girls are fitted with a chestworn combined heart-rate accelerometer device to measure their physical activity throughout each 7-day period.

What are the possible benefits and risks of participating?

The benefits of participating include education on health-related scientific research and use of research equipment. In addition, the participants receive a certificate stating that they have completed the study and receive their individual results in an anonymous format. The research also has indirect benefits for the participants, as the findings will help to inform interventions to

increase physical activity in adolescent girls. In the long term, this could help with the prevention of obesity and diseases such as type 2 diabetes and heart disease. Possible risks included feeling light-headed during the exercise tests, but the cool down helps to reduce this and it rarely lasts more than a few minutes. There is also a risk of skin irritation caused by wearing the activity monitor. Therefore, all participants are shown how to remove and fit the monitor and wash the skin appropriately. Those who are allergic to any of the ingredients in the breakfast meals or had certain health conditions (e.g. diabetes, epilepsy) are not able to participate for their own safety.

Where is the study run from?

- 1. University of Bedfordshire (UK)
- 2. Loughborough University (UK)

When is study starting and how long is it expected to run for? July 2014 to December 2016

Who is funding the study?

- 1. British Academy (UK)
- 2. Leverhulme Trust (UK)
- 2. University of Bedfordshire (UK)
- 3. Loughborough University (UK)

Who is the main contact? Dr Julia Zakrzewski-Fruer Julia.Fruer@beds.ac.uk

### Contact information

### Type(s)

Scientific

#### Contact name

Dr Julia Zakrzewski-Fruer

#### **ORCID ID**

https://orcid.org/0000-0003-4167-4100

### Contact details

University of Bedfordshire Bedford United Kingdom MK41 9EA +44 (0)123 479 3410 Julia.Fruer@beds.ac.uk

### Additional identifiers

**Protocol serial number** SG142106

### Study information

#### Scientific Title

Comparison of daily and intermittent breakfast consumption on physical activity energy expenditure in girls

### **Study objectives**

Seven days of daily breakfast consumption will result in higher physical activity energy expenditure when compared with intermittent breakfast consumption.

### Ethics approval required

Old ethics approval format

### Ethics approval(s)

- 1. University of Bedfordshire Institute for Sport and Physical Activity Research Ethics Panel, 22 /09/2015, ref: 2015ISPAR013
- 2. Loughborough University Ethics Approvals (Human Participants) Sub-Committee, 30/03/2016, ref: R16-PO39

### Study design

Interventional within-participants cross-over study

### Primary study design

Interventional

### Study type(s)

Prevention

### Health condition(s) or problem(s) studied

Physical inactivity

#### **Interventions**

The participants completed two, 7-day conditions. The order of the conditions for each participant was produced using a computer-based random number generator by the principal investigator ensuring that the order across the total sample was counterbalanced. The conditions were not masked to the participant.

- 1. Daily breakfast consumption (DBC): the consumption of a standardized 400 kcal breakfast before 0900 h for 7 consecutive days
- 2. Intermittent breakfast consumption (IBC): the abstinence from all energy-providing nutrients until at least 1030 h on days 1, 3, 5 and 7 and the consumption of the 400 kcal standardized breakfast on days 2, 4 and 6

All breakfasts were weighed, pre-packaged and provided to the participants prior to each condition. The participants were instructed to consume the breakfasts provided at home. Participants completed both of the conditions and there was a 7-10 day washout between the conditions.

### Intervention Type

Other

### Primary outcome(s)

- 1. Physical activity energy expenditure from sedentary activities (<1.5 metabolic equivalent (METs)) during wake time to <1030 h (kJ/day)
- 2. Physical activity energy expenditure from sedentary activities (<1.5 METs) during 1030 h to <1530 h (kJ/day)
- 3. Physical activity energy expenditure from sedentary activities (<1.5 METs) during 1530 to bed time (kJ/day)
- 4. Physical activity energy expenditure from light activities (1.5-2.9 METs) during wake time to <1030 h (kJ/day)
- 5. Physical activity energy expenditure from light activities (1.5-2.9 METs) during 1030 h to <1530 h (kJ/day)
- 6. Physical activity energy expenditure from light activities (1.5-2.9 METs) during 1530 to bed time (kJ/day)
- 7. Physical activity energy expenditure from moderate activities (3.0-5.9 METs) during wake time to <1030 h (kJ/day)
- 8. Physical activity energy expenditure from moderate activities (3.0-5.9 METs) during 1030 h to <1530 h (kJ/day)
- 9. Physical activity energy expenditure from moderate activities (3.0-5.9 METs) during 1530 to bed time (kJ/day)
- 10. Physical activity energy expenditure from vigorous activities (>5.9 METs) during wake time to <1030 h (kJ/day)
- 11. Physical activity energy expenditure from vigorous activities (>5.9 METs) during 1030 h to <1530 h (kJ/day)
- 12. Physical activity energy expenditure from vigorous activities (>5.9 METs) during 1530 to bed time (kJ/day)

All primary outcomes are measured using combined heart rate/accelerometry continuously throughout each 7-day condition.

### Key secondary outcome(s))

- 1. Time spent in sedentary activities (<1.5 metabolic equivalent (METs)) during wake time to <1030 h (min/day)
- 2. Time spent in sedentary activities (<1.5 METs) during 1030 h to <1530 h (min/day)
- 3. Time spent in sedentary activities (<1.5 METs) during 1530 to bed time (min/day)
- 4. Time spent in light activities (1.5-2.9 METs) during wake time to <1030 h (min/day)
- 5. Time spent in light activities (1.5-2.9 METs) during 1030 h to <1530 h (min/day)
- 6. Time spent in light activities (1.5-2.9 METs) during 1530 to bed time (min/day)
- 7. Time spent in moderate activities (3.0-5.9 METs) during wake time to <1030 h (min/day)
- 8. Time spent in moderate activities (3.0-5.9 METs) during 1030 h to <1530 h (min/day)
- 9. Time spent in moderate activities (3.0-5.9 METs) during 1530 to bed time (min/day)
- 10. Time spent in vigorous activities (>5.9 METs) during wake time to <1030 h (min/day)
- 11. Time spent in vigorous activities (>5.9 METs) during 1030 h to <1530 h (min/day)
- 12. Time spent in vigorous activities (>5.9 METs) during 1530 to bed time (min/day)

All secondary outcomes are measured using combined heart rate/accelerometry continuously throughout each 7-day condition.

### Completion date

16/12/2016

### **Eligibility**

Key inclusion criteria

- 1. Aged 11 to 13 years old
- 2. Female

### Participant type(s)

Healthy volunteer

### Healthy volunteers allowed

No

### Age group

Child

### Lower age limit

11 years

### Upper age limit

13 years

### Sex

Female

### Key exclusion criteria

- 1. Allergies to the breakfast meals
- 2. Fitted with a pacemaker
- 3. Unable to walk
- 4. Unable to wear a chest-worn combined heart rate/accelerometer
- 5. Health related issues that could be affected by participation in the study (e.g., uncontrolled exercise-induced asthma, diabetes, epilepsy)

### Date of first enrolment

01/10/2015

### Date of final enrolment

12/09/2016

### **Locations**

### Countries of recruitment

United Kingdom

England

## Study participating centre University of Bedfordshire

Polhill Avenue Bedford United Kingdom MK41 9EA

### Study participating centre Loughborough University

Epinal Way Loughborough United Kingdom LE11 3TU

### Sponsor information

### Organisation

University of Bedfordshire

### Organisation

Loughborough University

### Organisation

University of Bedfordshire

### **ROR**

https://ror.org/0400avk24

### Funder(s)

### Funder type

University/education

### **Funder Name**

**British Academy** 

### Alternative Name(s)

BA British Academy, The British Academy, BA

### **Funding Body Type**

Private sector organisation

### Funding Body Subtype

Universities (academic only)

### Location

**United Kingdom** 

#### **Funder Name**

University of Bedfordshire

### Alternative Name(s)

### **Funding Body Type**

Private sector organisation

### Funding Body Subtype

Universities (academic only)

### Location

**United Kingdom** 

### **Funder Name**

Loughborough University

### Alternative Name(s)

Lbого

### **Funding Body Type**

Private sector organisation

### **Funding Body Subtype**

Universities (academic only)

#### Location

United Kingdom

### **Funder Name**

Leverhulme Trust

### Alternative Name(s)

The Leverhulme Trust

### **Funding Body Type**

Private sector organisation

### **Funding Body Subtype**

Other non-profit organizations

### Location

**United Kingdom** 

### **Results and Publications**

### Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are/will be available upon request from Dr Julia Zakrzewski-Fruer (Julia.Fruer@beds.ac.uk).

### IPD sharing plan summary

Available on request

### **Study outputs**

Output type	Details	Date created Date added	Peer reviewed?	Patient-facing?
Results article	results	01/02/2018	Yes	No
Participant information sheet	Participant information sheet	11/11/2025 11/11/2025	No	Yes