Does Acamprosate Decrease Cue-induced Alcohol Craving?

Submission date	Recruitment status	Prospectively registered
13/11/2007	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
18/01/2008	Completed	☐ Results
Last Edited	Condition category	Individual participant data
21/01/2008	Mental and Behavioural Disorders	☐ Record updated in last year

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Johan Franck

Contact details

Department of Clinical Neuroscience Section for Alcohol and Drug dependence Research Karolinska University Hospital, M4:02 Stockholm Sweden 17176 johan.franck@ki.se

Additional identifiers

EudraCT/CTIS number

2004-004514-17

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

EudraCT number: 2004-004514-17

Study information

Scientific Title

A randomised placebo-controlled trial of acamprosate effects on alcohol cue reactivity and alcohol priming in dependent patients

Acronym

DADCAC

Study objectives

- 1. Acamprosate attenuates cue-induced subjectively experienced and physiolological correlates of craving
- 2. Acamprosate attenuates alcohol-induced subjectively experienced and physiolological correlates of craving

Ethics approval required

Old ethics approval format

Ethics approval(s)

The trial was approved by:

- 1. The Regional Ethical Review Board in Stockholm on the 23rd March 2005 (ref: 2005/30-31/3)
- 2. The Swedish Medical Products Agency on the 5th April 2005 (EudraCT number: 2004-004514-17)

Study design

The study used a randomised, double blind, single-site, placebo-controlled design comparing cue- and alcohol-induced craving for acamprosate and placebo treated patients

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Not specified

Study type(s)

Treatment

Participant information sheet

Health condition(s) or problem(s) studied

Alcohol dependence

Interventions

Patients were assigned to 22 days of either acamprosate (1998 mg/day) or placebo treatment according to a randomisation process conducted by the Karolinska University Hospital pharmacy.

Medication (150 tablets containing 333 mg acamprosate or placebo) was dispensed once per patient, at the start of the trial, with instructions to intake 6 tablets per day (2 in AM, 2 midday, 2 in PM).

Intervention Type

Drug

Phase

Not Specified

Drug/device/biological/vaccine name(s)

Acamprosate

Primary outcome measure

- 1. The difference in subjectively experienced craving post exposure to non-alcohol related stimuli
- 2. The difference in subjectively experienced craving between pre- and post-consumption of an alcoholic drink

For both measures comparisons are made between acamprosate versus placebo treated patients. Measures are collected post 22 days of treatment. Patients then come to the clinic and go through a series of sessions where they are exposed to alcohol versus non-alcohol related stimuli, and also take part in a alcohol priming paradigm. Present alcohol craving is measured in connection to each exposure.

Secondary outcome measures

Physiological measures of craving:

- 1. Pulse
- 2. Blood-pressure
- 3. Cortisol in blood
- 4. Galvanic skin response
- 5. Skin temperature

Measures are collected post 22 days of treatment. Patients then come to the clinic and go through a series of sessions where they are exposed to alcohol versus non-alcohol related stimuli, and also take part in an alcohol priming paradigm. Present alcohol craving is measured in connection to each exposure.

Overall study start date

01/09/2005

Completion date

05/02/2007

Eligibility

Key inclusion criteria

- 1. A male or a non-pregnant/non-nursing female between 18 and 65 years of age
- 2. A goal of controlled drinking
- 3. An intact sense of smell

- 4. Fulfilling the criteria for alcohol dependence according to Diagnostic and Statistical Manual of Mental Disorders, 4th edition (DSM-IV)
- 5. Willingness to give informed consent and comply with study procedures
- 6. Alcohol consumption in 15 of the last 90 days

Participant type(s)

Patient

Age group

Adult

Lower age limit

18 Years

Sex

Both

Target number of participants

56

Key exclusion criteria

- 1. Seeking complete alcohol abstinence
- 2. Current use of any medication that interferes with salivation
- 3. A diagnosis of an Axis I psychiatric disorder according to DSM-IV criteria (including all forms of substance dependence other than nicotine and alcohol)
- 4. A current use of psychoactive medications to manage schizophrenia, bipolar disorder, or major depression
- 5. Inpatient alcohol detoxification within the last 4 days
- 6. Acamprosate medication during the last year
- 7. Use of illegal drugs during the course of the study

Date of first enrolment

01/09/2005

Date of final enrolment

05/02/2007

Locations

Countries of recruitment

Sweden

Study participating centre
Department of Clinical Neuroscience
Stockholm

Sweden 17176

Sponsor information

Organisation

Addiction Centre Stockholm (Beroendecentrum Stockholm) (Sweden)

Sponsor details

Box 17914 Stockholm Sweden 118 95

Sponsor type

Hospital/treatment centre

Website

http://www.beroendecentrum.com

ROR

https://ror.org/04g380834

Funder(s)

Funder type

Research organisation

Funder Name

AFA insurances (AFA försäkringar) (Sweden) - http://www.afaforsakring.se

Funder Name

Systembolagets Council for Alcohol Research (Systembolagets råd för alkoholforskning) (Sweden) - http://www.can.se/sa/node.asp?node=1663

Funder Name

Milan Valverius Foundation (Sweden) - http://www.salusansvar.se/info/default.aspx? FolderID=3063a24d-26b2-416d-b00a-1d4d06810b2e

Funder Name

Foundation for Research on Psychiatric Diseases (Psykiatrifonden) (Sweden) - http://www.psykiatrifonden.se

Results and Publications

Publication and dissemination planNot provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summaryNot provided at time of registration