

Traditional laparoscopic cholecystectomy versus SILS™ port laparoscopic cholecystectomy

Submission date	Recruitment status	<input type="checkbox"/> Prospectively registered
13/01/2009	No longer recruiting	<input type="checkbox"/> Protocol
Registration date	Overall study status	<input type="checkbox"/> Statistical analysis plan
27/02/2009	Completed	<input checked="" type="checkbox"/> Results
Last Edited	Condition category	<input type="checkbox"/> Individual participant data
21/05/2013	Surgery	

Plain English summary of protocol

Not provided at time of registration

Contact information

Type(s)

Scientific

Contact name

Dr Jeffrey Marks

Contact details

11100 Euclid Avenue
Mail Stop 5047
Cleveland
United States of America
44106

Additional identifiers

ClinicalTrials.gov (NCT)

NCT00832767

Protocol serial number

AS08012

Study information

Scientific Title

Prospective randomised controlled trial of traditional laparoscopic cholecystectomy versus SILS™ port laparoscopic cholecystectomy

Acronym

SILS™ Port Laparoscopic Cholecystectomy Study

Study objectives

The objectives of this trial are: 1) to assess the feasibility and safety of performing SILS™ port cholecystectomy, and 2) to monitor and compare the outcomes of SILS™ port cholecystectomy versus 4PLC to objectively document the scientific merit, and the perceived advantages of SILS™ port cholecystectomy.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Italy: Ethics Committee, Universita Cattolica del Sacro Cuore, approved on 02/12/2008 (ref: Prot. cm.P822 [A.1505]/C.E./2008)

UK: To be submitted to Ethics Committee, Imperial College London, St Mary's Hospital, in January 2009.

USA: University Hospital Case Medical Center, Institutional Review Board for Human Investigations. Approval pending as of 13/01/2009.

Study design

Randomised controlled single-blind multi-centre study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Acute calculus, acalculous cholecystitis

Interventions

Traditional laparoscopic cholecystectomy versus SILS™ port laparoscopic cholecystectomy

Intervention Type

Procedure/Surgery

Phase

Phase IV

Primary outcome(s)

1. Feasibility and safety of SILS™ port cholecystectomy versus four-port traditional laparoscopic cholecystectomy as indicated by intraoperative and postoperative adverse events up to one year
2. Operative time
3. Blood loss

Key secondary outcome(s)

1. Pain, assessed by the Pain Intensity numerical rating scale at 8 different timepoints within the first month post surgery

2. Cosmesis, assessed at 5 different timepoints between 1 week and 1 year post surgery by the following:
 - 2.1. Modified Hollander scale
 - 2.2. Body image questionnaire
 - 2.3. Photo series questionnaire
3. Quality of life, assessed by the SF-8® and SF-12® Health Survey questionnaires at 7 different timepoints, once prior to surgery and 6 times up to 1 month post surgery
4. Time to return to normal activity
5. Time required for insertion of SILS™ Port compared to 4 standard ports

Completion date

01/02/2011

Eligibility

Key inclusion criteria

1. Both males and females, between 18 and 65 years old
2. The patient has a diagnosis of biliary colic with documented gallstones or polyps by imaging or biliary dyskinesia with documented ejection fraction (EF) <30%
3. Body mass index (BMI) <35 kg/m²
4. The patient or patient's legal representative has been informed of the nature of the study, agrees to its provisions, and has provided written informed consent as approved by the Institutional Review Board/Ethics Committee (IRB/EC) of the respective clinical site
5. The patient agrees to return to the same research facility for all study-required post-procedure follow-up visits

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Key exclusion criteria

1. Any female patient who is pregnant, suspected pregnant, or nursing
2. Any patient with acute calculus or acalculous cholecystitis
3. Any patient who has had an upper midline or right sub costal incision
4. Any patient with pre-operative indication for a cholangiogram
5. Any patient with American Society of Anesthesiologists (ASA) >3 with normal liver function
6. Any patient who is undergoing peritoneal dialysis (PD)
7. Any patient who has an unrepaired umbilical hernia or has had prior umbilical hernia repair

Date of first enrolment

03/02/2009

Date of final enrolment

01/02/2011

Locations

Countries of recruitment

United Kingdom

Italy

United States of America

Study participating centre

11100 Euclid Avenue

Cleveland

United States of America

44106

Sponsor information

Organisation

Covidien Surgical Devices (USA)

ROR

<https://ror.org/00grd1h17>

Funder(s)

Funder type

Industry

Funder Name

Covidien Surgical Devices (USA)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	01/06/2013		Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes