# To evaluate the impact of a Carer Support Needs Assessment Tool (CSNAT) intervention in hospice home care

Submission date	Recruitment status	[X] Prospectively registered
22/06/2012	No longer recruiting	Protocol
Registration date	Overall study status	Statistical analysis plan
22/06/2012	Completed	Results
Last Edited	Condition category	Individual participant data
06/09/2016	Other	Record updated in last year

#### Plain English summary of protocol

Background and study aims

Most people prefer to be cared for and to die in the familiar surroundings of their own home. Being able to make this choice very much depends on the efforts of family carers. While taking on a caring role is normally done willingly, it can result in considerable burden and negative effects on carers health. Therefore it is important to ensure that carers are well supported to enable them to look after patients at home while also maintaining their own health and wellbeing. So far health professionals have had no clear guidelines to help them ensure that they assess and address carer support needs regularly and in detail. So we developed a simple assessment form, the Carer Support Needs Assessment Tool (CSNAT), based on what carers told us their main support needs were in end of life home care. The procedures for CSNAT assessment, prioritisation and follow-up together form the CSNAT intervention. The aim of the study is to find out whether the using the CSNAT intervention with carers during care giving makes a difference to carer outcomes in bereavement.

### Who can participate?

All adult carers of patients looked after by the hospice home care (HHC) services can take part in the study.

#### What does the study involve?

We will ask six HHC services to use the CSNAT intervention with carers of patients under their care. We will then compare bereavement outcomes for carers who had usual care with those who had the intervention. The HHC services will start using the CSNAT at three month intervals so we can make a comparison between services who have started the intervention with those who have not, and also make a comparison before and after the start of the intervention within each service. We will measure outcomes using a postal survey of carers 4-5 months after bereavement. To reduce any potential risks, HHC will exclude carers that are believed to suffer particularly complicated grief in bereavement, and contact details to HHC bereavement support will be provided. The study will also ask the views of the HHC nurses and some of the carers who

had the intervention on what made the intervention successful or not so successful. Finally, we will look at HHC CSNAT records to see whether some factors, such as how often assessments were done, seem to make a difference.

What are the possible benefits and risks of participating? Participants in the CSNAT intervention group may receive better support during care giving and experience better health, well being and lower levels of grief in bereavement.

Where is the study run from? University of Manchester (UK).

When is the study starting and how long is it expected to run for? The study started in May 2012 and will run for 2 years.

Who is funding the study? National Institute for Health Research (NIHR) (UK).

Who is the main contact? Gunn Grande gunn.grande@manchester.ac.uk

## **Contact information**

### Type(s)

Scientific

#### Contact name

Prof Gunn Grande

#### Contact details

School of Nursing, Midwifery and Social Work Oxford Road Manchester United Kingdom M13 9PL +44 161 306 7695 gunn.grande@manchester.ac.uk

## Additional identifiers

**EudraCT/CTIS** number

**IRAS** number

ClinicalTrials.gov number

Secondary identifying numbers 12422

## Study information

#### Scientific Title

A randomised trial to evaluate the impact of a Carer Support Needs Assessment Tool (CSNAT) intervention in hospice home care

#### **Acronym**

**CSNAT** 

#### **Study objectives**

To test whether a formalised, comprehensive procedure for carer support needs assessment, prioritisation and follow up (CSNAT intervention) improves quality of care and carer outcomes in end of life home care, compared to usual care.

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

NRES Committee North West - Greater Manchester East, 23/03/2012, ref: 12/NW/0206

#### Study design

Stepped wedge cluster randomised trial

#### Primary study design

Interventional

#### Secondary study design

Randomised controlled trial

### Study setting(s)

Hospital

## Study type(s)

Quality of life

### Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

## Health condition(s) or problem(s) studied

Topic: Primary Care Research Network for England, Generic Health Relevance and Cross Cutting Themes; Subtopic: Not Assigned, Generic Health Relevance (all Subtopics); Disease: All Diseases, Health Services Research

#### **Interventions**

CSNAT, Formalised, comprehensive procedure for carer support needs assessment, prioritisation and follow up using the Carer Support Needs Assessment Tool (CSNAT)

Control, informal carer assessment as usual

Follow Up Length: 5 month(s)

#### Intervention Type

Other

#### **Phase**

Phase II/III

#### Primary outcome measure

SF-12 4-5 months post bereavement

#### Secondary outcome measures

- 1. Quality of care; Timepoint(s): 4-5 months post bereavement
- 2. Patient's place of death; Timepoint(s): Date of death
- 3. Texas Revised Inventory of Grief; Timepoint(s): 4-5 months post bereavement

#### Overall study start date

21/09/2012

#### Completion date

31/12/2013

## **Eligibility**

#### Key inclusion criteria

- 1. The main bereaved carer of patients who have been supported by participating hospice home care services (HHCs)
- 2. Target Gender: Male & Female; Lower Age Limit 18 years

#### Participant type(s)

**Patient** 

#### Age group

Adult

#### Lower age limit

18 Years

#### Sex

Both

#### Target number of participants

Planned Sample Size: 1003; UK Sample Size: 1003

#### Key exclusion criteria

- 1. Age < 18 years
- 2. Identified by HHCs as experiencing particularly complicated grief in bereavement

#### Date of first enrolment

21/09/2012

#### Date of final enrolment

## Locations

#### Countries of recruitment

England

**United Kingdom** 

Study participating centre School of Nursing, Midwifery and Social Work Manchester United Kingdom M13 9PL

## Sponsor information

## Organisation

University of Manchester (UK)

## Sponsor details

c/o Samantha Wade Oxford Road Manchester England United Kingdom M13 9PL +44 208 843 8054 Samantha.Wade@nihr-ccf.org.uk

### Sponsor type

University/education

#### **ROR**

https://ror.org/027m9bs27

## Funder(s)

### Funder type

Government

#### **Funder Name**

National Institute for Health Research (NIHR) (UK) - Central commissioning facility (CCF); Grant Codes: PB-PG-0610-22291

## **Results and Publications**

**Publication and dissemination plan**Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

**IPD sharing plan summary**Not provided at time of registration