Evaluation of the clinical effects of supplementation with probiotic supplement Proxian® in frail elderly patients treated with enteral nutrition

Submission date	Recruitment status No longer recruiting	Prospectively registered		
07/07/2020		[X] Protocol		
Registration date 09/07/2020	Overall study status Completed	Statistical analysis plan		
		[X] Results		
Last Edited	Condition category	[] Individual participant data		

Plain English summary of protocol

Background and study aims

Enteral feeding refers to intake of food via the gastrointestinal (GI) tract. The GI tract is composed of the mouth, esophagus, stomach, and intestines. Enteral feeding may mean nutrition taken through the mouth or through a tube that goes directly to the stomach or small intestine.

The main aim of the study is to evaluate the effects of probiotic supplement Proxian® on infections and inflammation in frail elderly patients treated with enteral nutrition. The secondary objectives of the study are to evaluate the effects of the administration of Proxian® food supplement on the nutritional status and intestinal motility and to assess the compliance of caregivers of HEN patients and their perception of the product in different stages of administration (before and after).

Who can participate?

Patients older than 65 years treated with HEN for at least 1 month at INRCA service, who didn't assume the antibiotic therapy for at least 1 month before the enrollment, without diagnosis of pancreatitis and/or hypersensitivity to any ingredient contained in Proxian®, who didn't use any probiotic supplement in the period of 30 days previous to the enrollment and didn't participate in any clinical trial in 3 months previous to the invitation to participate in the study, whose CRP hs value is lower than 10 mg / dL, who will give their written informed consent (for patients with a serious cognitive impairment the informed consent signed by legal administrator) and whose caregiver is willing to participate.

What does the study involve?

The study will run for 90 days after the enrolment (signing the free consent). The Proxian® probiotic supplement will be administered to the intervention group for 60 days. The placebo (maltodextrin) will be administered to the control group for the same duration. The follow up will be performed for both groups 30 days after the end of administration. CRP values will be measured at enrolment, 45 days and 90 days. Information on clinical manifestations of infections

and antibiotic therapy will be gathered from caregivers on regular basis. Patients are free to decide about their participation and to withdraw from the study at any time.

What are the possible benefits and risks of participating? There are no risks of participating in the study.

Where is the study run from? IRCCS Istituto Nazionale Ricovero e Cura per Anziani (Italy)

When is the study starting and how long is it expected to run for? April 2018 to September 2019

Who is funding the study? Investigator initiated and funded

Who is the main contact? Dr Paolo Orlandoni p.orlandoni@inrca.it

Contact information

Type(s)

Scientific

Contact name

Dr Paolo Orlandoni

ORCID ID

https://orcid.org/0000-0003-1786-6743

Contact details

IRCCS Istituto Nazionale Ricovero e Cura per Anziani Via Della Montagnola 81 Ancona Italy 60127 +39 (0)71 8006353 p.orlandoni@inrca.it

Type(s)

Public

Contact name

Mrs Nikolina Jukic Peladic

ORCID ID

https://orcid.org/0000-0002-6023-2917

Contact details

IRCCS Istituto Nazionale Ricovero e Cura per Anziani Via Della Montagnola 81 Ancona Italy 60127 +39 (0)71 8003327 t.nutrizionale@inrca.it

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

CE INRCA 18015

Study information

Scientific Title

May the use of probiotic supplements in older patients treated with enteral nutrition reduce infections and inflammation?

Acronym

INTEGPro

Study objectives

Supplementation with Proxian® probiotic supplement in frail elderly patients treated with Enteral Nutrition (EN) reduces clinical infections and inflammation

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 20/09/2018, Ethics board of IRCCS INRCA (Via della Montagnola 81, 60129 Ancona, Italy; +39 (0)718003500; comitatoetico@inrca.it), ref: CE INRCA 18015

Study design

Randomized double-blind two-arm single-centre pilot study

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Clinical infections and inflammation in older subjects treated with home enteral nutrition

Interventions

A total of 32 subjects will be randomly assigned to the intervention group (16) or to controls (16). The Proxian® probiotic supplement will be administered to the intervention group for 60 days. Proxian® is a food supplement (Errekappa spa patent) containing encapsulated and gastro-protected probiotic micro-organisms. A Proxian® stick contains the following doses of probiotic microorganisms: Bifidobacterium lactis (DSM 23032) ≥ 1 billion cells / dose, Lactobacillus plantarum LP01 (LMG P-21021) ≥ 1 billion cells / dose and Lactobacillus buchneri Lb26 (DSM 16341) tindalized, naturally rich in selenium (20 mg). Both bifido bacteria and lactobacilli (in particular Lactobacillus plantarum) have proven to be very effective in reducing intestinal permeability (22). Being Lactobacillus buchneri Lb26 naturally rich in Selenium, Proxian® also guarantees a better bio-availability of this mineral which, together with Zinc, also contained in the Proxian® supplement, contributes to the protection of cells from oxidative stress and to the normal functioning of the immune system.

The placebo (maltodextrin) will be administered to the control group for the same duration.

The packaging of the two products - Proxian and placebo - will be identical and will be identifiable through a special code that neither the patients or caregivers responsible for the administration or researcher will know. The intervention group will be administered, through the enteral probe, 1 sachet / day of Proxian® dissolved in water. The administration will be carried out by formal and / or informal caregivers (carers and family members) who deal with the management of the patient and will be carried out 1 hour after the administration of the enteral formula. To ensure correct administration of the product, all caregivers will be specifically trained and they will be given the written instructions concerning the method of administration of the product. The caregivers will also be instructed on how to recognize and note the progress of the alvo and eventual adverse effects due to the administration of the product. In order to ensure the adherence of the participants to the project, telephone contacts on daily bases will be made with the caregivers.

Intervention Type

Supplement

Primary outcome(s)

- 1. The onset of infections observed by detection of clinical manifestations of infections (prevalence) and antibiotic therapy (prevalence) during the period of administration and 90 days after the enrollment (30 days after the last administration)
- 2. The inflammation is assessed by C Reactive Protein (CRP) values in the intervention group and in controls following the administration of products (Proxian® and placebo) and lab analyses will be performed at baseline, 45 and 90 days after the enrollment

Key secondary outcome(s))

- 1. Nutritional status is assessed by Body Mass Index values assessed at baseline, 45 and 90 days after the enrollment
- 2. Motility is assessed by the evidence on bowel function and its variations as reported by caregivers on the occasion of each phone contact with caregivers
- 3. Compliance of caregivers and their perception of the product in different stages of administration (before and after) are assessed by specific questionnaires administered at the enrollment and after 60 days of administration

Completion date

30/09/2019

Eligibility

Key inclusion criteria

- 1. Age over 65 years
- 2. Patient treated with HEN for at least 1 month
- 3. No antibiotic therapy for at least 1 month before the enrollment
- 4. Written informed consent; for patients with a serious cognitive impairment the informed consent signed by legal administrator
- 5. Availability of a caregiver (formal or informal)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Senior

Sex

All

Total final enrolment

43

Key exclusion criteria

- 1. Diagnosis of pancreatitis
- 2. Known hypersensitivity to any ingredient contained in Proxian®
- 3. Participation in other clinical trials in 3 months previous to the invitation to participate in the study
- 4. Use of probiotic supplements in the period of 30 days previous to the enrollment
- 5. CRP hs> 10 mg / dL

Date of first enrolment

02/10/2018

Date of final enrolment

15/06/2019

Locations

Countries of recruitment

Italy

Study participating centre IRCCS Istituto Nazionale Ricovero e Cura per Anziani

Via della Montagnola 81 Ancona

Italy

60127

Sponsor information

Organisation

IRCCS Istituto Nazionale Ricovero e Cura per Anziani

ROR

https://ror.org/04tfzc498

Funder(s)

Funder type

Other

Funder Name

Investigator initiated and funded. ERREKAPPA Spa. provided free samples of Proxian® and placebo.

Results and Publications

Individual participant data (IPD) sharing plan

Dataset generated and analyzed during the current study will be available upon request which is to be addressed to MD Paolo Orlandoni, Clinical Nutrition, IRCCS INRCA Ancona, Via della Montagnola 81, 60127 Ancona, Italy (p.orlandoni@inrca.it; 0039/(0)71/8003653). Data will be available only after the publication of study results in a high-impact peer- reviewed journal. Data may be requested for the involvement of participants in studies on larger samples relatively to the efficacy of probiotics in similar populations. The request must be formulated by scientific contact/responsible of the study to MD Paolo Orlandoni. Data will be shared only after the acceptance of participants or legal representatives.

IPD sharing plan summary

Available on request

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	27/01/2021	27/01/2021	Yes	No
Participant information sheet			07/08/2020		Yes
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Protocol file	version V2	03/09/2018	07/08/2020	No	No