A pre-pregnancy study examining the effects of an intensive lifestyle package supported with Liraglutide treatment, a medication equivalent to a natural hormone produced in the stomach, in obese women with previous history of pregnancy diabetes

Submission date 21/12/2016	Recruitment status No longer recruiting	[X] Prospectively registeredProtocol
Registration date 22/12/2016	Overall study status Completed	Statistical analysis planResults
Last Edited 30/05/2017	Condition category Nutritional, Metabolic, Endocrine	Individual participant dataRecord updated in last year

Plain English summary of protocol

Background and study aims

Up to 15% of pregnant women develop a temporary type of diabetes during pregnancy called gestational diabetes mellitus (GDM). In GDM, women develop high blood sugar levels as the body is unable to produce enough insulin (a hormone that helps convert sugar into stored energy). GDM is associated with increased risks for mother and baby including increased risk of later life diabetes for the mother and increased risk of childhood obesity for the baby. If a woman overweight or obese or has previously had GDM, then their risk of developing it is higher. Treatments during pregnancy to reduce the risk of GDM to date have been unsuccessful, and so it is important to look at the months before pregnancy as a time to intervene and reduce the chance of developing GDM in a subsequent pregnancy. Optimising mother's health prior to a pregnancy holds considerable potential to improve mother and baby's health in pregnancy. This study is looking at one such intervention in women at risk of GDM before pregnancy. The aim of this study is to find out whether this programme, which includes intensive lifestyle changes and treatment with a medication called Liraglutide, is feasible and acceptable to women.

Who can participate?

Women who are obese (without being diabetic) and/or have previously had GDM.

What does the study involve?

Participants are randomly allocated to one of two groups. Those in the first group take part in the GDM prevention intervention. This involves taking part in an intensive lifestyle change programme involving a healthy diet and physical activity, as well as receiving daily injections of liraglutide (a drug which mimics the action of a protein called GLP-1 which triggers the release of insulin). Those in the second group receive intensive lifestyle advice also. Participants in both

groups come to a clinic service in a monthly basis over a 6-month period, where various measurements and blood collection are carried out.

What are the possible benefits and risks of participating?

Participants may benefit from potentially improved the blood sugar levels and a reduction of the incidence of diabetes in subsequent pregnancies, as well as improved control of blood glucose in participants with type 2 diabetes mellitus and a reduction body weight. Liraglutide is generally tolerated well, however the most frequently reported adverse effects of Liraglutide are gastrointestinal (gut), more than 5% of patients will present with nausea, diarrhoea, vomiting, constipation, abdominal pain, and dyspepsia (indigestion). Serious side effects are rare during Liraglutide treatment.

Where is the study run from?
National Maternity Hospital (Ireland)

When is the study starting and how long is it expected to run for? September 2016 to December 2020

Who is funding the study?
University College Dublin (Ireland)

Who is the main contact? Professor Fionnuala McAuliffe fionnuala.mcauliffe@ucd.ie

Contact information

Type(s)

Scientific

Contact name

Prof Fionnuala McAuliffe

ORCID ID

https://orcid.org/0000-0002-3477-6494

Contact details

UCD Perinatal Research Centre
Obstetrics & Gynaecology
School of Medicine
University College Dublin
National Maternity Hospital
Dublin
Ireland
Dublin 2
+353 1 6373216
fionnuala.mcauliffe@ucd.ie

Additional identifiers

Protocol serial number

EC 26.2016

Study information

Scientific Title

An open-label randomized trial of an intensive lifestyle package supported with Liraglutide treatment in obese, non-pregnant women with previous history of Gestational Diabetes Mellitus

Study objectives

Pre-pregnancy treatment with an intensive lifestyle intervention supported by daily Liraglutide injections in obese non-pregnant women with previous gestational diabetes is feasible and acceptable.

Ethics approval required

Old ethics approval format

Ethics approval(s)

National Maternity Hospital Ethics Committee, 14/11/2016, ref: EC 26.2016

Study design

Single-centre open label randomised controlled trial

Primary study design

Interventional

Study type(s)

Prevention

Health condition(s) or problem(s) studied

- 1. Gestational diabetes
- 2. Glucose intolerance
- 3. Obesity

Interventions

Eligible participants are randomised to one of two groups in a ratio of 1:1 for usual care versus intervention using computer-generated allocation.

Intervention arm: The study intervention involves a treatment package of an intensive lifestyle approach (including diet and physical activity advice) supported by a daily treatment of Liraglutide for a period of 6 months. The control group will have usual care.

At baseline and at 6 months an OGTT will be carried out and bloods analysed for metabolic parameters such as HOMA-IR, hsCRP, lipids and HbA1c.

A 3-day food diary and a lifestyle questionnaire will be given to participants, at baseline and at 6 months, to assess behaviour change, readiness and analyse nutritional quality of the diet. At 6 months the intervention arm will complete an acceptability questionnaire to assess patient views of the intervention and a diary card to assess compliance with study medication. At the end of the treatment/intervention period of 6 months the participant will have finished the study trial. Participants will be contacted if conception is achieved, to follow up in their potential development of gestational diabetes or impaired glucose tolerance.

Control arm as of 02/05/2017:

Control arm: Participants will receive intensive lifestyle alone. Participants will be followed up after 6 months.

Original control arm:

Control arm: Participants receive usual care, including general lifestyle advice. Participants will be followed up after 6 months.

Intervention Type

Mixed

Primary outcome(s)

- 1. Proportion of eligible women who would agree to participate in the study. This will be calculated by the % of women who agree to study participation out of all those eligible women who were approached for study participation.
- 2. Acceptability of women of taking daily Liraglutide injections. This will be assessed by an acceptability questionnaire completed after 6 months of study participation
- 3. Proportion of women that complied with the study protocol and completed the study intervention. This will be assessed by adherence to study drug using a drug compliance diary, attendance at study visits, satisfactory collection of study endpoints

Key secondary outcome(s))

- 1. Fasting glucose levels are measured in fasting plasma samples at baseline and 6 months
- 2. Effect on glucose homeostasis after treatment, as measured by OGTT, HbA1c, HOMA-ir at baseline and 6 months
- 3. Effect on inflammatory markers, including C- reactive protein, as measured by blood samples at baseline and 6 months
- 4. Weight loss as assessed by weighing participants at baseline and 6 months
- 5. Gestational diabetes mellitus incidence and/or impaired glucose tolerance in a subsequent pregnancy as measured by a positive result in a future pregnancy for gestational diabetes using standard diagnostic criteria

Completion date

31/12/2020

Eligibility

Key inclusion criteria

- 1. Severe obesity BMI≥ 35 kg/m2 without Type 2 Diabetes Mellitus and with previous gestational diabetes (with or without insulin)
- 2. Participants must be able and willing to give written informed consent and to comply with the requirements of this study protocol
- 3. Participant must be a female, aged 18 years or above at baseline
- 4. Planning a pregnancy within the next one to two years
- 5. Negative pregnancy test
- 6. Contraception during the study period

Removed 02/05/2017:

Post pregnancy Type 2 Diabetes Mellitus with BMI≥ 30 kg/m2 and HbA1C >42 mmol/mol

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Female

Key exclusion criteria

- 1. Allergy/sensitivity to study medication
- 2. Female subjects who are pregnant or breast-feeding or considering becoming pregnant during the study period
- 3. Medical disorder requiring medication other than stable hypertension, hypothyroidism, polycystic ovarian syndrome
- 5. Ongoing abuse of alcohol or narcotics
- 6. Family or personal history of multiple endocrine neoplasia type 2 (MEN2) or familial medullary thyroid carcinoma (FMTC)
- 7. Personal history of non-familial medullary thyroid carcinoma
- 8. History of acute or chronic pancreatitis
- 9. Obesity induced by drug treatment
- 10. Use of approved weight lowering pharmacotherapy
- 11. Previous surgical treatment of obesity
- 12. History of major depressive disorder or suicide attempt
- 13. Uncontrolled hypertension
- 14. Subjects unable to provide written informed consent

Added 30/05/2017:

15. Type 2 diabetes mellitus

Date of first enrolment

01/04/2017

Date of final enrolment

31/01/2020

Locations

Countries of recruitment

Ireland

Study participating centre National Maternity Hospital Holles St, Grand Canal Dock

Sponsor information

Organisation

Clinical Research Centre - University College Dublin

ROR

https://ror.org/05m7pjf47

Funder(s)

Funder type

University/education

Funder Name

University College Dublin

Alternative Name(s)

UCD

Funding Body Type

Private sector organisation

Funding Body Subtype

Universities (academic only)

Location

Ireland

Results and Publications

Individual participant data (IPD) sharing plan

The current data sharing plans for the current study are unknown and will be made available at a later date

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type

Details

Date created Date added Peer reviewed? Patient-facing?

Participant information sheet Participant information sheet 11/11/2025 No

Yes