A clinical trial to see if LPRI-424 is safe and works for Polycystic Ovary Syndrome compared to a placebo for 9 months

Submission date	Recruitment status	Prospectively registered		
19/07/2022	No longer recruiting	∐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
14/09/2022	Completed	Results		
Last Edited	Condition category	Individual participant data		
21/12/2022	Nutritional, Metabolic, Endocrine	Record updated in last year		

Plain English summary of protocol

Background and study aims

The test medicine (LPRI-424) is being given to women with polycystic ovary syndrome (PCOS) to see if there is a change in hair growth when given for 9 months. This is being tested against a placebo and the hair growth is being measured by a tool called the Adapted Modified Ferriman-Gallwey (mFG) score. This score is used by Doctors to see how much hair growth there is for those suffering with polycystic ovary syndrome.

Who can participate?

Women aged 14 to 40 years diagnosed with PCOS.

What does the study involve?

Not provided at time of registration

What are the possible benefits and risks of participating? Benefits:

Improvement of PCOS symptoms in particular of hirsutism. Information learned from the trial may help other patients with hirsutism in the future.
Risks:

The main burden for subjects would be the E-Diary completion each day. The burden is minimised by taking less than 3 minutes to complete and the subjects being compensated for their time/compliance with the E-Diary. The amount of compensation will be given within the patient information sheet. Subjects will not be allowed any permanent hair removal during the trial, they can only use shaving as a method of hair removal. If the subject chooses to shave areas of their bodies, they must do so 15 days prior to the mFG score being tested to ensure the same standardised assessment of hair growth is done. This burden will be minimised by providing compensation to those patients who comply and complete the trial.

Other risks and burdens include blood withdrawals, where the subject may get injection-site pain /discomfort. To mitigate this risk only a trained, experienced and delegated member of the study site team will perform blood withdrawals. The subjects will also have gynaecological examinations and intravaginal ultrasounds, this is an intrusive examination and the experience

will be minimised by a trained and experienced member of the site team performing the assessments.

Where is the study run from? Chemo Research S.L. (Spain)

When is the study starting and how long is it expected to run for? September 2020 to November 2023

Who is funding the study? Chemo Research S.L. (Spain)

Who is the main contact? Dr Thozhukat Sathyapalan, thozhukat.sathyapalan@hyms.ac.uk

Contact information

Type(s)

Scientific

Contact name

Dr Andrew Bradshaw

Contact details

Scope international UK
St George's House
St George's Road
Bolton
United Kingdom
BL1 2DD
+44 1204 322 725
regulatory.uk@scope-international.com

Type(s)

Principal investigator

Contact name

Dr Thozhukat Sathyapalan

Contact details

Hull Royal Infirmary Anlaby Road Hull United Kingdom HU3 2JZ

thozhukat.sathyapalan@hyms.ac.uk

Additional identifiers

Clinical Trials Information System (CTIS)

2021-002178-17

Integrated Research Application System (IRAS)

1006162

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

LPRI-424/304, IRAS 1006162, CPMS 47924

Study information

Scientific Title

A multicentre, phase III, double-blind, randomised clinical trial to assess the efficacy and safety of LPRI-424 (dienogest 2.00 mg / ethinyl estradiol 0.02 mg) in the treatment of polycystic ovary syndrome (PCOS) versus placebo during 9 cycles

Study objectives

Primary objective:

To demonstrate the efficacy of LPRI-424 on hirsutism, measured by an adapted modified Ferriman-Gallwey (mFG) score based on the investigator decision against placebo

Secondary objective:

To assess the safety of LPRI-424, bleeding pattern and health-related quality of life

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 13/09/2022, Wales REC 5 (Health and Care Research Wales Support and Delivery Centre, Castlebridge 4, 15-19 Cowbridge Road East, Cardiff, CF11 9AB, UK; no telephone number provided; Wales.REC5@wales.nhs.uk), ref: 22/WA/0230

Study design

Interventional double blind randomized placebo controlled trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Excess of body hair in women with polycystic ovarian syndrome (PCOS)

Interventions

Duration of treatment: 9 sequential cycles of 28 days (252 days in total)

Dose and mode of administration:

a) Test product: LPRI-424 tablets (dienogest [DNG] 2 mg / ethinyl estradiol [EE] 0.02 mg), film-coated tablets, oral administration. Each blister contains 28 tablets. One treatment cycle consists of 24 active white tablets followed by 4 green placebo tablets.

The tablets will be taken once daily at approximately the same time each day.

b) Reference therapy: LPRI-424 tablets matching placebo tablets, film-coated tablets, oral administration. Each blister contains 28 tablets. One treatment cycle consists of 24 white placebo tablets followed by 4 green placebo tablets.

The tablets will be taken once daily at approximately the same time each day.

Follow-up activity for both arms:

- Document concomitant medication
- Vital signs (blood pressure, heart rate, respiration rate, body temperature), height only in adolescent subjects, body weight, waist circumference
- Gynaecological examination
- Urine pregnancy test
- Check for Adverse events

Randomisation process: Online tool

Intervention Type

Drug

Phase

Phase III

Drug/device/biological/vaccine name(s)

dienogest 2.00 mg, ethinyl estradiol 0.02 mg

Primary outcome(s)

Hirsutism is measured using a modified Ferriman-Gallwey score (mFG) based on the investigator's decision at baseline, treatment cycle 6 (day 20+/- 7 of medication cycle) and at the end of treatment.

Key secondary outcome(s))

Changes in laboratory parameters from baseline to trial termination

- 1. Adverse events (AEs)
- 2. Vital signs
- 3. Clinical laboratory parameters
- 4. Physical examination
- 5. Electrocardiogram (ECG)
- 6. Gynaecological examination
- 7. Cervical cytology
- 8. Changes in weight and waist
- 9. Change in bleeding pattern from baseline
- 10. Number of subjects with prolonged bleeding/spotting >10 days
- 11. IP acceptability based on IP satisfaction and wellbeing
- 12. Changes in quality of life based on the Polycystic Ovarian Syndrome Questionnaire (PCOSQ) from baseline

Completion date

Eligibility

Key inclusion criteria

At Visit 1a (Screening) and at Visit 1b (IP Dispensation), subjects must meet ALL of the following criteria:

1. Two years postmenarcheal women of any ethnic origin between 14 and 40 years (inclusive at the time of enrolment), not seeking pregnancy.

Female subjects at risk of pregnancy aged between 14 and 17 years (inclusive) provided that:

- a. Applicable national, state and local laws allow subjects in this age group to consent/assent to receive contraceptive services,
- b. All applicable laws and regulations regarding the informed consent/assent of the subjects to participate in clinical trials are observed.
- 2. Diagnosed with PCOS according to the following criteria:
- a. Presence of hirsutism measured using an adapted mFG with a cut-off value of ≥7 (at V1a only based on patient's history or interview; to be determined only at Visit 1b) and
- b. Presence of one of the following criteria:
- 1. Oligomenorrhea (≤6 menses per year) while not using hormonal contraceptives
- 2. Polycystic ovaries defined as presence of 12 or more follicles measuring 2 9 mm throughout the entire ovary or an ovarian volume ≥ 10 cm³ determined by ultrasound. This criterion applies only for subjects aged ≥ 16 years due to the high incidence of multi-follicular ovaries in young subjects.

(Ovarian cyst are any fluid filled structure >30 mm in diameter that persisted for more than 2 cycles and enlarged follicles are any structure similar to an ovarian cyst that did not persist).

- 3. Informed consent form (ICF)/assent form signed voluntarily before any study-related procedure is performed; indicating that the subject understands the purpose of and procedures required for the study and is willing to participate in the study.
- 4. Body Mass Index (BMI): 18 kg/m²≤ BMI <35 kg/m²
- 5. At screening, maximum systolic blood pressure (median value of 3 values) \leq 140 mm Hg and diastolic blood pressure (median value of 3 values) \leq 90 mm Hg.
- 6. Haematology and chemistry parameters, heart rate (HR) and electrocardiogram (ECG) within reference range, or showing no clinically relevant deviations, as judged by the investigator.
- 7. Willing to use trial medication for nine 28-day cycles.
- 8. Is consenting to use reliable nonhormonal contraceptive methods (condoms, female or male sterilisation or sexual abstinence) during the study from screening until the final examination visit.
- 9. Must be willing to avoid the use of all hair removal or growth procedures and products apart from shaving during the trial and not to shave the hairs for a period of 14 days before the modified adapted mFG score will be determined during study participation.
- 10. Has a good physical and mental health as determined on the basis of medical history and general physical examination performed at screening.
- 11. Willing to adhere to the prohibitions and restrictions specified in the protocol and being able to comply with the use of the trial medication or the use of the trial diary.
- 12. Agrees to not participate in any other clinical trials during the course of this trial.

Participant type(s)

Patient

Healthy volunteers allowed

Age group

Mixed

Sex

Female

Key exclusion criteria

Subjects are to be excluded from the trial for ANY ONE of the following reasons:

- 1. Pregnancy or wish for pregnancy.
- 2. Breastfeeding women.
- 3. Current smoker with age \geq 36 years and/or with BMI >30 kg/m² (at the time of trial enrolment)
- 4. Abnormal finding on pelvic, breast or intravaginal ultrasound examination (or transabdominal ultrasound examination for adolescents) that in the investigator's opinion contraindicates participation in the trial.
- 5. Women ≥21 years of age with a Papanicolaou (pap) smear reading low-grade squamous intraepithelial lesion (LSIL) or higher at screening (or 6 months prior to screening date). Subjects with atypical squamous cells of undetermined significance (ASC-US) can be included if they are negative for high-risk Human papilloma virus (HPV) strains. Subjects <21 years of age do not require a Pap smear.
- 6. Poorly controlled diabetes (HgbA1c >6.5%). Treatment with metformin is allowed only if it is expected to remain stable during the trial (defined as a stable dose for at least 3 months before V1a).
- 7. Anaemia (haemoglobin <10 g/dL).
- 8. Subjects with disorders other than PCOS, that can result in menstrual irregularity and hyperandrogenism
- 9. Known contraindication or hypersensitivity to ingredients or excipients of the IP, including:
- 9.1. Presence or risk of a venous thromboembolism (VTE)
- 9.2. Presence or risk of an arterial thromboembolism (ATE)
- 9.3. Presence or history of pancreatitis, if it is associated with severe hypertriglyceridemia
- 9.4. Presence or history of liver diseases in which liver function has not returned to normal (also Dubin-Johnson and Rotor syndrome)
- 9.5. Current or previous liver tumours
- 9.6. Known or suspected sex hormone-dependent malignant tumours
- 9.7. Undiagnosed vaginal bleeding
- 9.8. Unexplained amenorrhoea
- 9.9. Concomitant use of medicinal products containing ombitasvir/paritaprevir/ritonavir or dasabuvir
- 10. Uncontrolled chronic concomitant diseases (i.e., not on a stable treatment dose for at least 2 months at the time
- 11. Severe Covid-19 disease or less than 3 months after hospitalisation due to a Covid-19 disease.
- 12. Evidence or history of alcohol, medication or drug abuse (within the last 12 months prior to consent/assent).
- 13. Known or suspected human immunodeficiency virus (HIV) and/or hepatitis infection at screening.
- 14. Previous intake of hormonal contraceptives within the last 3 months prior consent/assent or concomitant intake of hormonal contraceptives
- 15. Previous or concomitant intake of 5a-reductate inhibitors (e.g. flutamide) or similar inhibitors
- 16. Long-term treatment (longer than 7 consecutive days within a month prior to V1b) of any medication that might interfere with the efficacy of hormonal contraceptives

- 17. Prohibited medication include the use of oestrogens, progestogens, strong microsomal enzyme-inducing drugs (intensive and moderate frequency).
- 18. Other prohibited medications include antiandrogens, insulin sensitizers or drugs that might interfere with blood pressure regulation, lipid profile or carbohydrate metabolism within the last 6 months prior to consent/assent.
- 19. Administration of human chorionic gonadotropin (hCG) or intake of co-medication containing hCG within a month prior to V1b.
- 20. Inositol treatment within the last 3 months prior to consent/assent.
- 21. Evidence or history of clinically significant psychiatric illness or suicide risk.
- 22. Planned surgery during the anticipated time of participation in this trial requiring withdrawal of an oral contraceptive.
- 23. Participation in any other trial of an investigational drug or device parallel to the current trial or less than 90 days before consent/assent, or previous participation in a clinical trial with LPRI-421 or LPRI-424, or previous participation in the current trial including dispensed trial medication.
- 24. Subject is a member of the investigator's or sponsor's staff or a relative or family member thereof.
- 25. Any condition that, in the opinion of the investigator, may jeopardize protocol compliance or the scientific integrity of the trial.

Date of first enrolment 01/11/2021

Date of final enrolment 31/01/2023

Locations

Countries of recruitment United Kingdom	
Czech Republic	
Hungary	
Lithuania	
Poland	
Serbia	
Slovakia	
Spain	

Study participating centre

Ukraine

United Kingdom

Sponsor information

Organisation

Chemo Research S.L.

Funder(s)

Funder type

Industry

Funder Name

Chemo Research S.L.

Results and Publications

Individual participant data (IPD) sharing plan

The current data sharing plans for this study are unknown and will be available at a later date

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023	No	No