# Multicultural Information on Depression: Effects of a multilingual information website intervention on the levels of depression literacy and depression related stigma in Greek-born and Italian-born immigrants living in Australia

Submission date	Recruitment status  No longer recruiting	Prospectively registered		
08/01/2010		☐ Protocol		
Registration date	Overall study status	Statistical analysis plan		
04/02/2010	Completed	[X] Results		
Last Edited	Condition category	Individual participant data		
29/12/2020	Mental and Behavioural Disorders			

## Plain English summary of protocol

Not provided at time of registration

## Study website

http://www.MIDonline.com.au

## Contact information

## Type(s)

Scientific

#### Contact name

Dr Litza Kiropoulos

### Contact details

Monash University
School of Psychology and Psychiatry
Monash Medical Centre
Level 3, Block P
246 Clayton Rd, Clayton
Melbourne
Australia
3168
+61 (0)3 9594 1467
litza.kiropoulos@med.monash.edu.au

## Additional identifiers

**EudraCT/CTIS** number

**IRAS** number

ClinicalTrials.gov number

**Secondary identifying numbers** N/A

## Study information

### Scientific Title

Effects of a multilingual information website intervention on the levels of depression literacy and depression related stigma in Greek-born and Italian-born immigrants living in Australia: a cross-sectional randomised controlled trial

### **Acronym**

**MIDonline** 

### **Study objectives**

In this study, we evaluate the impact of a new web-based multilingual intervention (MIDonline) on depression literacy, personal and perceived stigma and level of depression among Greekborn and Italian-born immigrants living in Melbourne, Australia.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Monash University Standing Committee on Ethics in Research involving Humans approved on the 27th June 2006 (ref: 2006/546)

## Study design

Single centre cross-sectional randomised controlled trial

## Primary study design

Interventional

## Secondary study design

Randomised controlled trial

### Study setting(s)

Other

## Study type(s)

Quality of life

## Participant information sheet

Not available in web format, please use the contact details below to request a patient information sheet

### Health condition(s) or problem(s) studied

Depression and depression related stigma

### **Interventions**

Participants were randomly assigned either to the MIDonline intervention or the control/no intervention interview condition. Participants in both conditions spent on average 1 - 1.5 hours to complete the MIDonline condition or the control interview.

### MIDonline Intervention:

The intervention comprised the consumer stream of the MIDonline website (http://www. MIDonline.com.au) which provides online multilingual information about depression designed for middle- to older-aged consumers from a non-English speaking background. The material is available in the Greek, Italian and English languages. The website content incorporates information about symptoms and case studies of depression, how depression is diagnosed, related disorders, causes, treatment options, how to find a bilingual mental health professional and professional psychological care, stigma related to mental illness, and multilingual translated resources. The MIDonline website also includes separate sections for carers and mental health professionals. The information is provided is a culturally relevant way. For example, case studies are representative of middle- to older-aged people of both genders who are Greek-born and Italian-born.

### Control condition:

The control intervention consisted of a semi-structured interview with a bilingual interviewer who asked open-ended questions about the participant's depression related illness schemas. Specifically, questions were asked about the causes, important symptoms, course and development, treatments and outcomes of depression.

The follow-up questionnaire was completed an average of 7.95 days (sd = 2.34) after participants completed the post-intervention questionnaire for both the MIDonline and control/interview arms.

## Intervention Type

Other

### Phase

Not Applicable

### Primary outcome measure

- 1. Depression literacy, assessed using translated and adapted versions of the D-Lit scale which is a 22-item true/false test of knowledge about depression. A higher score on this scale indicated greater literacy.
- 2. Depression related stigma; changes in personal (reflecting the participants' personal attitudes) and perceived (reflecting participants' beliefs about the attitudes of others) depression stigma were assessed using the 18-item Depression Stigma Scale. Half of the items required participants to rate how strongly they personally agreed with a statement about depression (e.g., people with depression are unpredictable). The other half of the items asked the participant to indicate what they thought most other people believed about the same issue (e.g. 'most people believe that people with depression are unpredictable'). Ratings were made

on a five-point Likert scale. Scores on the total scale range from 0 to 36 for the full scale and 0 to 18 for each of the two nine-item sub-scales, with higher scores indicating greater stigma.

3. Level of depression. Depression severity was measured with the Beck Depression Inventory - version 2 (BDI-II) which is a 21-item instrument for measuring severity of depression in adults. Responses to items covered the 'past two weeks, including today'. Responses on the BDI-II items range from 0 to 6 with higher values indicating higher severity.

All participants who took part in the study completed the pre-, post- and 1 week follow-up questionnaires which included the above outcome measures. The pre- questionnaire was completed just prior to taking part in either the MIDonline intervention or the control condition and the post- assessment straight after the completion of the intervention or control condition. The follow up questionnaire was completed an average of 7.95 days (s.d. = 2.34) after participants completed the post-intervention questionnaire.

### Secondary outcome measures

Socio-demographic variables and clinical history such as:

- 1. Age
- 2. Gender
- 3. Birthplace
- 4. Marital status
- 5. Level of education
- 6. Occupation level
- 7. Currently working
- 8. Length of stay in Melbourne
- 9. Greek identification
- 10. Italian identification
- 11. Australian identification
- 12. Current living situation
- 13. English language proficiency
- 14. Current work status
- 15. Alcohol use
- 16. Memory functioning
- 17. Reported physical conditions
- 18. Frequency of physical activity
- 19. Tobacco use
- 20. Currently receiving treatment for an emotional or psychological problem
- 21. What type of treatment being received and from whom
- 22. Indicators of quality of life

Recorded at the pre-assesment time point only.

### Overall study start date

01/11/2006

### Completion date

01/06/2009

## **Eligibility**

Key inclusion criteria

- 1. Middle- to older-aged (45+ years of age), either sex
- 2. Greek-born or Italian-born first generation immigrants
- 3. Living in Melbourne, Australia

### Participant type(s)

**Patient** 

### Age group

Adult

### Sex

Both

### Target number of participants

At least 60 Greek-born immigrants and 60 Italian-born immigrants in each of the intervention and control groups (202 in total)

### Total final enrolment

202

### Key exclusion criteria

Does not meet inclusion criteria

### Date of first enrolment

01/11/2006

### Date of final enrolment

01/06/2009

## Locations

### Countries of recruitment

Australia

## Study participating centre Monash University Melbourne Australia

3168

## Sponsor information

### Organisation

The beyondblue Victorian Centre of Excellence in Depression and Related Disorders (Australia)

### Sponsor details

PO BOX 6100 Hawthorn West Melbourne, Victoria Australia 3122 +61 (0)3 9810 6100 bb@beyondblue.org.au

### Sponsor type

Research organisation

### Website

http://www.beyondblue.org.au/index.aspx?

### **ROR**

https://ror.org/05mwvz623

## Funder(s)

### Funder type

Research organisation

### **Funder Name**

The beyondblue Victorian Centre of Excellence in Depression and Related Disorders (Australia)

## **Results and Publications**

### Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

## IPD sharing plan summary

Not provided at time of registration

## **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	19/04/2011	29/12/2020	Yes	No