

Effect of acupuncture treatment on stiff neck

Submission date 22/02/2011	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input checked="" type="checkbox"/> Protocol
Registration date 10/03/2011	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
Last Edited 27/12/2023	Condition category Musculoskeletal Diseases	<input type="checkbox"/> Individual participant data

Plain English summary of protocol

Background and study aims

There have been no reported randomized placebo-controlled trials investigating the effectiveness of acupuncture under double-blind conditions using practitioner-patient blinded needles. The aim of this study is to verify the effectiveness of acupuncture for functional neck /shoulder stiffness.

Who can participate?

Patients aged over 18 with stiff neck not from organic disease.

What does the study involve?

Before treatment, patients will be asked about the intensity of their stiff neck. Patients will be randomly allocated to receive one of the following four acupuncture treatments:

1. Treatment with penetrating needles
2. Treatment with skin-touch placebo needles (the needle tip presses against the skin)
3. Treatment with no-touch placebo needles (the needle tip does not reach the skin)
4. Treatment without any needle applications (control group)

Immediately after and 24 hours after treatment, the patients are asked about the intensity of their stiff neck. Also, the acupuncturists and patients are asked to record whether the treatment is 'no-touch control', 'skin-touch placebo', 'penetrating', or 'unidentifiable'.

What are the possible benefits and risks of participating?

Participants will contribute to developing acupuncture and moxibustion studies. Although rare, there is the possibility of needle pain, unpleasantness, bleeding and hematoma with acupuncture.

Where is the study run from?

Tokyo Ariake University of Medical and Health Sciences (Japan).

When is the study starting and how long is it expected to run for?

The study started in June 2010 and will run until March 2015.

Who is funding the study?

Grants-in-Aid for Scientific Research (B) (Japan).

Who is the main contact?
Prof Nobuari Takakura

Contact information

Type(s)
Scientific

Contact name
Prof Nobuari Takakura

Contact details
2-9-1 Ariake, Koto-ku
Tokyo
Japan
135-0063

Additional identifiers

Protocol serial number
RCT

Study information

Scientific Title
Effect of acupuncture treatment on stiff neck: a randomised, double-blind, placebo-controlled trial

Study objectives
Differences exist in therapeutic effect on stiff neck between treatments with the penetrating, skin-touch placebo, no-touch control needle and a control group without any needles. The aim of the study is to assess whether verum penetrating needle treatment has specific effect over placebo needle treatment on stiff neck.

On 23/07/2014 the following changes were made to the trial record:
1. The overall trial start date was changed from 24/01/2011 to 09/06/2010.
2. The overall trial end date was changed from 31/03/2013 to 31/03/2015.

Ethics approval required
Old ethics approval format

Ethics approval(s)
Ethics Committee of Tokyo Ariake University of Medical and Health Sciences, 08/07/2010

Study design
Randomised double-blind placebo-controlled single-centre trial

Primary study design
Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Musculoskeletal discomfort

Interventions

Treatments using the double-blind needle:

1. The no-touch needle which the needle tip does not reach the skin
2. Skin-touch placebo needle which the needle tip presses against the skin
3. Verum penetrating needle
4. Treatment without any needle applications (control group)

Before treatment, patients are asked about the intensity of their stiff neck on a visual analogue scale. In this trial, each of five acupuncturists applies acupuncture treatment to 60 patients. For each treatment, the acupuncturist applies four needles to four acupoints on the neck and shoulder in the patients. Immediately after and 24 hours after treatment, the patients are asked about the intensity of their stiff neck on a visual analogue scale. Also, acupuncturists (practitioners) and patients are asked to record whether the treatment is 'no-touch control', 'skin-touch placebo', 'penetrating', or 'unidentifiable'. The same methods are applied to the control group without any needles and relating questions are asked.

Intervention Type

Other

Phase

Not Applicable

Primary outcome(s)

Improvement in stiff neck: before, immediately and 24 hours after needle application, the patients are asked about the intensity of their stiff neck on a Visual Analogue Scale (VAS) ranging from 0 (no stiff neck) to 100 (the most intense stiff neck imaginable)

Key secondary outcome(s)

Effect of patient and practitioner blinding using the double-blind needle: after completion of treatment, the practitioners and patients are asked to record whether they think the treatment is 'penetrating', 'skin-touch', 'no-touch' or 'utterly unidentifiable'

Completion date

31/03/2015

Eligibility**Key inclusion criteria**

1. Patients with stiff neck not from organic disease
2. Age range: over 18 years, both male and female

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

All

Total final enrolment

400

Key exclusion criteria

Patients who have:

1. Cervical spondylosis
2. Cervical disc herniation
3. Cervico-omo-brachial syndrome
4. Thoracic outlet syndrome
5. Liver, gallbladder or heart disease
6. High blood pressure
7. Cerebral vascular disease
8. Neurological disease
9. Under any pharmacological treatment
10. Patients who will receive any cure for stiff neck or acupuncture treatment within 24 hours after treatment in this study

Date of first enrolment

24/01/2011

Date of final enrolment

31/08/2014

Locations**Countries of recruitment**

Japan

Study participating centre

2-9-1 Ariake, Koto-ku

Tokyo

Japan

135-0063

Sponsor information

Organisation

Japan Society for the Promotion of Science (Japan)

ROR

<https://ror.org/00hhkn466>

Funder(s)

Funder type

Government

Funder Name

Grants-in-Aid for Scientific Research (B) (Japan)

Results and Publications

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		09/12/2023	27/12/2023	Yes	No
Protocol article	protocol	16/07/2014		Yes	No