# Evaluation of the efficiency of autologous albumin gel mixed with I-PRF and artificial bone graft in alveolar ridge preservation following tooth extraction

Submission date 25/05/2024	<b>Recruitment status</b> No longer recruiting	<ul> <li>Prospectively registered</li> <li>Protocol</li> </ul>
<b>Registration date</b> 20/06/2024	<b>Overall study status</b> Completed	<ul><li>Statistical analysis plan</li><li>Results</li></ul>
Last Edited 22/07/2025	<b>Condition category</b> Oral Health	<ul><li>Individual participant data</li><li>[X] Record updated in last year</li></ul>

### Plain English summary of protocol

#### Background and study aims

When a tooth is lost, the surrounding bone, known as the alveolar bone, begins to deteriorate. This process, termed alveolar bone resorption, results in a decrease in both the quantity and guality of the bone, leading to a reduction in its external width and height. Ridge preservation is a treatment carried out during or after tooth extraction to mitigate this bone loss. It aims to slow down bone resorption, promote new bone formation, and enhance the healing of both bone and surrounding soft tissues. Various biomaterials have been tested for ridge preservation, including autografts (bone from the same individual), bone substitutes, blood derivatives, and bioactive agents. Platelet-rich fibrin (PRF), a second-generation platelet concentrate, has shown promise in promoting tissue regeneration by enhancing blood vessel formation, stem cell capture, immune response, and tissue closure. However, PRF is rapidly absorbed, typically degrading within 10 to 14 days, which limits its effectiveness as a long-term barrier. Recent advancements have led to the development of an injectable albumin gel combined with PRF, which prolongs the material's absorption time to 4-6 months. This new compound leverages the benefits of PRF while offering a slower degradation rate, potentially providing more sustained support for bone and tissue regeneration. The purpose of this research is to evaluate the efficacy of this innovative injectable albumin gel mixed with PRF and artificial bone graft in preserving the alveolar ridge following tooth extraction. The study will focus on 1. assessing the pain experienced by patients during the treatment; 2. measuring the radiographic density of the newly formed bone; and, 3. analyzing the changes in the dimensions of the alveolar ridge, specifically the vestibular (cheek side) and lingual (tongue side) walls, six months post-surgery. Alveolar ridge resorption is a significant issue after dental extractions, leading to aesthetic and functional problems and complicating future dental implant placement. This study aims to offer a potential solution to enhance the preservation of the alveolar structure, thereby improving patient outcomes.

Who can participate?

Adult patients aged between 22 and 50 years old requiring the preservation of the alveolar structure following tooth extraction

What does the study involve?

This study aims to compare the effectiveness of two different combinations of treatments (one with autologous albumin gel and PRF, and the other with injectable PRF) in maintaining alveolar bone dimensions post-extraction.

What are the possible benefits and risks of participating? The benefits gained from the procedure are improved healing and more bone. There are no risks after the application of bone graft.

Where is the study run from? Damascus University (Syria)

When is the study starting and how long is it expected to run for? August 2022 to April 2024

Who is funding the study? Damascus University (Syria)

Who is the main contact? Dr Zain Moalla, zainmoalla150@gmail.com

# **Contact information**

**Type(s)** Public, Scientific, Principal Investigator

**Contact name** Dr Zain Moalla

ORCID ID https://orcid.org/0009-0001-6184-3699

**Contact details** Damascus university, Mazzeh highway Damascus Syria -+963 94758006

+963 94758006 zainmoalla150@gmail.com

**Type(s)** Principal Investigator

**Contact name** Dr Zain Moalla

**Contact details** 

Damascus university, Mazzeh highway Damascus Syria -+963 94758006 zainmoualla06@damascusuniversity.edu.sy

# Additional identifiers

EudraCT/CTIS number Nil known

### **IRAS number**

**ClinicalTrials.gov number** Nil known

**Secondary identifying numbers** Nil known

# Study information

#### Scientific Title

Evaluation of the efficiency of autologous albumin gel mixed with I-PRF and artificial bone graft in alveolar-ridge-preservation following tooth extraction

#### **Study objectives**

Is the application of autologous albumin gel mixed with injectable platelet-rich fibrin and artificial bone graft able to improve healing, relieve pain, prevent post-extraction alveolar bone resorption, and improve the bone density of new bone?

#### Ethics approval required

Ethics approval required

#### Ethics approval(s)

Approved 29/08/2022, Scientific Research and Postgraduate Studies Council (Baramkeh, Damascus, -, Syria; +963 1133923192; ap.srd@damascusuniversity.edu.sy), ref: 4243

### Study design

Comparative clinical trial with a split mouth technique

#### **Primary study design** Interventional

Secondary study design Non randomised study

**Study setting(s)** University/medical school/dental school

# Study type(s)

Efficacy

Participant information sheet

No participant information sheet available

## Health condition(s) or problem(s) studied

Symmetrical teeth indicated for extraction

## Interventions

The study aims to compare two different treatments for maintaining alveolar bone dimensions after tooth extraction. On one side, autologous albumin gel mixed with platelet-rich fibrin (PRF) and artificial bone graft will be applied, while on the opposite side, injectable platelet-rich fibrin (I-PRF) mixed with artificial bone graft will be used. To ensure unbiased results, the specific treatment for each side will be randomly assigned using Excel software, and blinding will be maintained by pulling papers to determine the treatment for each party. The intervention involves atraumatic extraction of two teeth from a single patient in one session, with each extraction taking 20-40 minutes. Both extraction sites will receive bone grafts, but with different PRF combinations to assess their effectiveness.

## Intervention Type

Procedure/Surgery

### Primary outcome measure

 Radiographic Bone Density: ROI Region of Interest By drawing a rectangle in the extraction socket in the image taken six months after the alveolar ridge preservation procedure and recording the average value given, measured using the Ez3Di viewer program
 Study of changes in alveolar bone dimensions: The two images are uploaded to the OnDemand3D program. We perform matching in the three axes - sagittal, frontal, and vertical and then choose the appropriate section of the image in which the surgical intervention was performed.

3. Measuring vertical changes (alveolar ridge height): A line is drawn that connects the top of the vestibular plate to the top of the lingual or palatal plate, and from the middle of this line a line is drawn to the height of the socket. In this way, the vertical height of the socket is measured immediately after the extraction and six months after the extraction, and the amount of change that occurs is calculated after that.

4. Measuring horizontal bone changes (width of the alveolar ridge): Two horizontal lines are drawn at level 3 and level 5 mm from the edge of the alveolar bone on the image taken immediately after the extraction. These lines are fixed when measuring on the second image (six months later) after merging the two images together, and the amount is then calculated. The change is happening.

# Secondary outcome measures

Pain is measured using a visual analogue scale (VAS) at baseline 24, 48, and 72 h

# Overall study start date

29/08/2022

**Completion date** 01/04/2024

# Eligibility

### Key inclusion criteria

1. There are no general diseases

- 2. There is no periodontitis
- 3. There is no contraindication for local anesthesia or oral surgery
- 4. Extraction socket type 1 according to Elian, 2007

### Participant type(s)

Patient

### Age group

Adult

#### **Lower age limit** 22 Years

Upper age limit

50 Years

**Sex** Both

**Target number of participants** 12

**Total final enrolment** 12

### Key exclusion criteria

- 1. Pregnant and breastfeeding
- 2. The presence of an acute abscess or vestibular fistula in the work area
- 3. Smoker patient
- 4. Alcoholics
- 5. Taking medications that affect bone metabolism

# Date of first enrolment 01/02/2023

# Date of final enrolment 01/01/2024

# Locations

**Countries of recruitment** Syria

Study participating centre

**Damascus University** Mazzeh highway Damascus Syria

# Sponsor information

**Organisation** Damascus University

**Sponsor details** Mazzeh highway Damascus Syria -+963981555778 damascusuniversity.edu.sy@gmail.com

**Sponsor type** University/education

Website http://www.damascusuniversity.edu.sy

ROR https://ror.org/03m098d13

# Funder(s)

**Funder type** University/education

**Funder Name** Damascus University

Alternative Name(s) University of Damascus, , DU

**Funding Body Type** Government organisation

**Funding Body Subtype** Universities (academic only) **Location** Syria

# **Results and Publications**

### Publication and dissemination plan

Planned publication in a peer-reviewed journal

# Intention to publish date 31/12/2024

#### Individual participant data (IPD) sharing plan

The dataset to be generated will be stored in a publicly available repository

#### IPD sharing plan summary

Study outputs

Stored in publicly available repository

Output type	Details	Date created	Date added	Peer reviewed?	Patient- facing?
<u>Other</u> publications	Alveolar Ridge Preservation Using a Mixture of Alb-PRF and Alloplastic Bone Graft: A Case Report	13/03 /2025	22/07 /2025	Yes	No