# Impact of poor nutrition on survival rates in patients with aortic dissection undergoing heart surgery

Submission date	Recruitment status	Prospectively registered
22/10/2024	No longer recruiting	☐ Protocol
Registration date	Overall study status	Statistical analysis plan
25/10/2024	Completed	Results
Last Edited	Condition category	Individual participant data
24/10/2024	Surgery	[] Record updated in last year

## Plain English summary of protocol

Background and study aims

Aortic dissection is a serious and potentially deadly condition where the inner layer of the aorta tears. This study looks at how a person's nutritional status, measured by the prognostic nutritional index (PNI), affects their chances of survival after surgery for type B aortic dissection (TBAD).

#### Who can participate?

The study included patients from the Aortic Dilatation Database of the First Affiliated Hospital of Naval Medical University. Patients with connective tissue diseases, traumatic dissection, secondary interventions, tumors, or incomplete data were not included.

#### What does the study involve?

Participants' medical records were reviewed to gather data. The study used various statistical models to analyze the relationship between PNI and patient outcomes after surgery.

#### What are the possible benefits and risks of participating?

The study aims to identify if PNI can predict survival rates, which could help doctors better assess and manage patients with TBAD. Since this is a retrospective study, there are no direct risks or benefits to the participants.

#### Where is the study run from?

The study is conducted at the First Affiliated Hospital of Naval Medical University in China.

When is the study starting and how long is it expected to run for? The study reviewed data from January 2009 to July 2023.

#### Who is funding the study?

The study is funded by the National Natural Science Foundation of China, the Shanghai Emerging Cross Disciplinary Research Project, and the Research Project of Shanghai Municipal Health Commission.

Who is the main contact?
Prof. Jian Zhou: zhoujian1\_3@163.com
Dr. Shuangshuang Li: lishuangshuangfy@163.com
Ms. Wen Li: daiyaliwen@hotmail.com

# **Contact information**

# Type(s)

Scientific

#### Contact name

Dr Shuangshuang Li

#### **ORCID ID**

https://orcid.org/0000-0001-5887-2280

#### Contact details

168 Changhai Road, Yangpu District Shanghai China 200433 +86 21-31161670/13611826460 lishuangshuangfy@163.com

#### Type(s)

Principal investigator

#### Contact name

Prof Jian Zhou

#### Contact details

168 Changhai Road, Yangpu District Shanghai China 200433 +86 13818896067 zhoujian1 3@163.com

#### Type(s)

Public

#### Contact name

Ms Wen Li

#### Contact details

225 Changhai Road, Yangpu District Shanghai China 200433 +86 17821451628 daiyaliwen@hotmail.com

# Additional identifiers

# Clinical Trials Information System (CTIS)

Nil known

#### ClinicalTrials.gov (NCT)

Nil known

#### Protocol serial number

Nil known

# Study information

#### Scientific Title

Low prognostic nutritional index predicts case fatality of type B aortic dissection undergoing thoracic endovascular aortic repair

#### **Study objectives**

PNI could serve as a valuable marker for predicting 1-year all-cause mortality in patients with Type B aortic dissections.

# Ethics approval required

Ethics approval required

# Ethics approval(s)

approved 24/08/2020, Shanghai Changhai Hospital Ethics Committee (168 Changhai Road, Yangpu District, Shanghai, 200433, China; +86-21-31162338; changhaiec@126.com), ref: CHEC-Y2020-042

## Study design

Single-center observational retrospective cohort study

# Primary study design

Observational

# Study type(s)

Prevention

# Health condition(s) or problem(s) studied

Type B aortic dissection (TBAD)

#### **Interventions**

This retrospective study, conducted from January 2009 to July 2023, included 1,524 patients with aortic dissection who were treated with Thoracic endovascular aortic repair in the aortic dilation database of the First Affiliated Hospital of Naval Medical University. Among them, 563 patients were subsequently excluded from the study. Therefore, only 912 patients were

included in the final analysis.

Demographic variables, laboratory tests, intra-operative details and anatomical characteristics were extracted from medical records. BMI calculated as weight in kilograms divided by height in meters squared. Aortic dissection was classified into acute (1–14 days), sub-acute (15–90 days), or chronic (>90 days) stages according to "2014 ESC guidelines on the diagnosis and treatment of aortic diseases". Blood samples were taken from all patients upon admission and tested by the laboratory. The Prognostic Nutritional Index (PNI) was subsequently calculated using the formula: PNI = albumin (g/L) +  $5 \times \text{lymphocyte}$  (109/L). Based on the derived PNI, patients were stratified into two groups: those with a low PNI (PNI < 46.83) and those with a high PNI (PNI  $\geq 46.83$ ), utilizing the specified cut-off point.

Qualified investigators meticulously monitored all patients at two distinct intervals: 30 days and one-year post-surgery, utilizing either telephonic surveys or medical records for data collection. Furthermore, an exhaustive review was conducted on the comprehensive clinical files of patients who were readmitted or treated as outpatients to identify any adverse events.

#### Intervention Type

Other

#### Primary outcome(s)

All-cause mortality at (30 days and one year post-surgery) using telephonic surveys or medical records for data collection

#### Key secondary outcome(s))

Aortic-related adverse events (ARAEs) and major adverse cardiac and cerebrovascular events (MACCEs) at (30 days and one year post-surgery) using telephonic surveys or medical records for data collection

# Completion date

01/07/2023

# **Eligibility**

## Key inclusion criteria

- 1. Age range: 18-80 years old
- 2. Patients with TBAD undergoing TEVAR surgery
- 3. Complete preoperative albumin, lymphocyte, and other blood test data available
- 4. No history of aortic surgery
- 5. No malignant tumor or connective tissue disease

# Participant type(s)

Patient

#### Healthy volunteers allowed

No

#### Age group

Adult

#### Lower age limit

18 years

#### Upper age limit

80 years

#### Sex

All

# Total final enrolment

912

#### Key exclusion criteria

- 1. Traumatic or iatrogenic aortic dissection
- 2. Connective tissue diseases such as:
- 2.1. Marfan syndrome
- 2.2. Turner's syndrome
- 2.3. Bicuspid aortic valve
- 2.4. Bechet's disease
- 2.5. Ehlers-Danlos syndrome
- 2.6. Giant cell arteritis
- 2.7. Ankylosing spondylitis
- 2.8. Takayasu arteritis
- 3. Previous aortic surgery
- 4. History of malignancy
- 5. Patients with incomplete clinical data

#### Date of first enrolment

01/01/2009

#### Date of final enrolment

01/07/2023

# Locations

#### Countries of recruitment

China

# Study participating centre

Department of Vascular Surgery, the First Affiliated Hospital of the Navy Medical University

Changhai Hospital, 168 Changhai Road, Yangpu District Shanghai

China

200433

# Sponsor information

# Organisation

National Natural Science Foundation of China

#### **ROR**

https://ror.org/01h0zpd94

# Funder(s)

## Funder type

Government

#### **Funder Name**

The National Natural Science Foundation of China [82270513]

#### **Funder Name**

Shanghai Emerging Cross Disciplinary Research Project [2022JC011]

#### **Funder Name**

Research project of Shanghai Municipal Health Commission [20224Y0351]

# **Results and Publications**

# Individual participant data (IPD) sharing plan

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request. zhoujian1\_3@163.com

# IPD sharing plan summary

Available on request

## **Study outputs**

Output type Details Date created Date added Peer reviewed? Patient-facing?

Participant information sheet 11/11/2025 No Yes