

Self-perceived impacts on daily living after periodontal treatment in a Brazilian elderly population.

Submission date 24/07/2015	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 21/08/2015	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 11/08/2015	Condition category Oral Health	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol

Background and study aims

Periodontitis (PD) is a very common chronic gum infection that damages the soft tissue and destroys the bone supporting the teeth. It can lead to tooth loss, difficulties chewing, poor appearance of teeth and gums and it can even increase the risk of a heart attack or stroke. It is caused by the build-up of bacteria in the mouth which, over time, combines with saliva and small food particles to form a sticky film over the teeth, called plaque. The bacteria in the plaque can result in gum disease, leading to swollen, painful gums. If not treated, this gum disease will get worse and will develop into periodontitis. This study investigates how elderly people think their quality of life has been affected by having treatment for their periodontitis, taking into account physical, psychological, social and environmental aspects as well as individual belief (OMS). The patients with chronic periodontal disease may experience emotional changes that can affect their daily life. The aim of this study was to compare these effects on daily living in an elderly Brazilian population after they have had periodontal treatment.

Who can participate?

Adults older than 60 with periodontal disease.

What does the study involve?

All participants have an initial periodontal clinical examination and have their quality of life (as affected by their oral health) assessed by them answering a set of questions and completing a questionnaire. They are then randomly allocated into one of two groups. Those in group 1 (test group) are given conventional non-surgical periodontal treatment that includes scaling and root planing along with oral hygiene instructions. Those in group 2 (control group) are not given the treatment at this time. Both groups of participants are then invited back for a second clinical examination and quality of life assessment 30 days after the start of the study and after the test group participants have had their treatment. Finally, the participants in the control group are offered the same non-surgical periodontal treatment once the study is complete.

What are the possible benefits and risks of participating?

There is no immediate direct benefit to those taking part. However, there should be benefits to

research related to periodontal health and quality of life in an elderly population. There are no risks to participating because patients are treated with conventional non-surgical periodontal treatment.

Where is the study run from?

Department of Dental Clinics, Division of Graduate Periodontics, Federal University of Rio de Janeiro (Brazil)

When is the study starting and how long is it expected to run for?

July 2004 to March 2005

Who is funding the study?

Federal University of Rio de Janeiro (Brazil)

Who is the main contact?

1. Mrs Flávia Santos

2. Mrs Anna Leão

Contact information

Type(s)

Scientific

Contact name

Mrs Flávia Santos

Contact details

Federal University of Rio de Janeiro
Department of Dental Clinics
Division of Graduate Periodontics
Niterói, RJ
Brazil
24220-121

Type(s)

Scientific

Contact name

Mrs Anna Leão

Contact details

Rio de Janeiro Federal University
Av . Carlos Chagas Filho 373
CCS building block K, 2nd floor
Room 56, University City Island
Rio de Janeiro
Brazil
21.941-902

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

N/A

Study information

Scientific Title

Self-perceived impacts on daily living in an elderly population regarding periodontal treatment.

Study objectives

The periodontal treatment does not cause impacts on the daily living of an elderly Brazilian population.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Clementino Fraga Filho University Hospital Human Research Committee, Federal University of Rio de Janeiro (UFRJ) - RJ, Brazil, 01/07/2004, ref: 089/04

Study design

Interventional single-centre study.

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Community

Study type(s)

Quality of life

Participant information sheet

Not available in web format, please use contact details to request a participant information sheet

Health condition(s) or problem(s) studied

Periodontal disease

Interventions

A sample of 53 patients aged over 60 were randomly allocated into one of 2 groups: control or test.

Participants in both groups had a periodontal clinical examination and had their oral health impact on their quality of life assessed by a set of 22 questions (PHQ-QoL) and through completing a questionnaire (OHIP-14) at baseline and then after the test group have had the therapy described below. Participants in the control group received the same therapy after the study was complete.

Therapy: full-mouth scaling and root planing.

Intervention Type

Other

Primary outcome measure

1. PHQ-QoL - a set of 22 questions related to gingival bleeding, halitosis, sensitivity, aesthetics dissatisfaction, eating and social restriction
2. OHIP-14 questionnaire, assessing quality of life.

Measurements were taken at baseline and after the periodontal treatment period

Secondary outcome measures

1. Probing pocket depth
2. Clinical attachment level
3. Visible plaque
4. Bleeding on probing
5. Tooth mobility
6. Number of Teeth

Measured at baseline and after the periodontal treatment period

Overall study start date

22/07/2004

Completion date

30/03/2005

Eligibility

Key inclusion criteria

1. Patients older than 60
2. Males or females
3. At the minimum of 15 teeth
4. At least 1 site with pocket depth >5 mm
5. Two teeth with clinical attachment level > 6 mm at baseline

Participant type(s)

Patient

Age group

Senior

Sex

Both

Target number of participants

53 patients (27 in control group and 26 in test group)

Key exclusion criteria

1. Total edentulism
2. Presence of acute periodontal diseases
3. Neurologic and auditive problems

Date of first enrolment

22/07/2004

Date of final enrolment

30/01/2005

Locations

Countries of recruitment

Brazil

Study participating centre

Department of Dental Clinics

Division of Graduate Periodontics

Federal University of Rio de Janeiro

Rio de Janeiro

Brazil

21.941-902

Sponsor information

Organisation

Federal University of Rio de Janeiro

Sponsor details

Department of Dental Clinics

Division of Graduate Periodontics

Rio de Janeiro

Brazil

21.941-902

Sponsor type

University/education

Website

www.odontologia.ufrj.br

ROR

<https://ror.org/03490as77>

Funder(s)

Funder type

Not defined

Funder Name

Investigator initiated and funded (Brazil)

Results and Publications

Publication and dissemination plan

I intend to publish this study in an international journal related to dental hygiene, prevention and quality of life in 2015.

Intention to publish date

01/12/2015

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Other