Pain relief for the first stage of labour using an electric massage chair

Submission date 04/07/2022	Recruitment status No longer recruiting	[X] Prospectively registered		
		☐ Protocol		
Registration date 03/08/2022	Overall study status Completed	Statistical analysis plan		
		[X] Results		
Last Edited	Condition category	Individual participant data		
10/09/2025	Pregnancy and Childbirth			

Plain English summary of protocol

Background and study aims

Labour is typically painful, especially during the first labour. Many women require pain relief in labour. Epidural, opiate-type drugs and inhaling gas and air are commonly offered forms of pain relief. These methods have potential undesired effects on the mother and her baby. Massage by a human therapist is an effective pain-relieving technique during labour. However, trained massage therapists are not available in the usual labour ward setting, the service is labour intensive and likely costly.

We plan to evaluate the use of a commercially available electric massage chair capable of providing a full body massage on its pain-relieving potential during the labour of women expecting their first baby as proof of the concept of mechanical massaging being effective.

Who can participate?

Women who have not had a previous pregnancy beyond 20 weeks (nulliparous), who are at full term, and who were admitted to the labour ward in the first stage of labour

What does the study involve?

Participants in one group will receive 30 minutes on a commercially procured electric massage chair that provides a full body massage program with the massaging function switched on and then 30 minutes with the massaging function switched off. Participants in a second group will do the opposite and receive 30 minutes on the electric massage chair with the massaging function switched off and then 30 minutes with the massaging function switched on. The groups will be compared using a Visual Numerical Rating Scale (VNRS) and questionnaire to score perceived pain.

What are the possible benefits and risks of participating?

There is no significant benefit or risk anticipated as all participants are exposed to the massage chair as this trial has a cross-over design. The massage provided by the chair may or may not be effective in providing pain relief. No significant harm is anticipated from the intervention with the massage chair as routine labour monitoring and care are continued during the massage.

Where is the study run from?
University Malaya Medical Centre (Malaysia)

When is the study starting and how long is it expected to run for? January 2022 to March 2023

Who is funding the study? Universiti Malaya (Malaysia)

Who is the main contact? Nadia Mazia (Malaysia) maziatunk@ummc.edu.my

Contact information

Type(s)

Scientific

Contact name

Dr Maziatun Nadia

Contact details

University Malaya Medical Centre (UMMC) Lembah Pantai Kuala Lumpur Malaysia 59100 +60 (0)13-2677656 maziatunk@ummc.edu.my

Type(s)

Principal investigator

Contact name

Prof Peng Chiong Tan

ORCID ID

https://orcid.org/0000-0001-8713-6581

Contact details

University Malaya Medical Centre (UMMC) Lembah Pantai Kuala Lumpur Malaysia 59100 +60 3 7841 4000 pctan@um.edu.my

Additional identifiers

Clinical Trials Information System (CTIS)

Nil known

ClinicalTrials.gov (NCT)

Nil known

Protocol serial number

2.0

Study information

Scientific Title

Massage chair as pain relief in the first stage of labour in nulliparas: A randomised cross-over trial (MASPAR)

Acronym

MASPAR

Study objectives

The objective of this proposed study is to evaluate the effect of massage chair therapy on labour pain

Ethics approval required

Old ethics approval format

Ethics approval(s)

Approved 13/6/2022, Medical Research Ethics Committee of The University of Malaya Medical Centre (Lembah Pantai, 59100 Kuala Lumpur, Malaysia; +60 (0)3-79493209; iresearch@ummc. edu.my), ref: 2022323-11097

Study design

Randomized cross-over trial

Primary study design

Interventional

Study type(s)

Treatment

Health condition(s) or problem(s) studied

Pain relief in labour

Interventions

Nulliparous women at term in the first stage of labour will be randomised to 30 minutes on the electric massage chair with the massaging function switched on then cross over to 30 minutes with the massaging function switched off or 30 minutes on the electric massage chair with the massaging function switched off then cross over to 30 minutes with the massaging function switched on. A commercially procured electric massage chair that provides a full body massage program will be used.

Intervention Type

Device

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Electric massage chair

Primary outcome(s)

Pain score (0 to 10 numerical rating scale) measured using an 11-point visual numerical rating scale (VNRS) (0 scoring as no pain at all to 10 as the worst pain imaginable) immediately after each 30-minute session on the massage chair with the massage function switched on or off

Key secondary outcome(s))

Maternal outcomes:

- 1. Mode of delivery measured by reviewing patient's notes after delivery
- 2. Intervention to delivery interval measured by reviewing patient's notes after delivery
- 3. Epidural anaesthesia in labour measured by reviewing patient's notes after delivery
- 4. Likert scale response to the questionnaire on pain relieving effectiveness of the massage by the massage chair after delivery

Neonatal outcomes:

- 4. Umbilical cord arterial pH measured by reviewing patient's notes at birth
- 5. Apgar score at 1 and 5 minutes measured by reviewing patient's notes at birth
- 6. Neonatal admission measured by reviewing patient's notes at birth

Completion date

02/03/2023

Eligibility

Key inclusion criteria

- 1. Nulliparous (no previous pregnancy beyond 20 weeks)
- 2. Aged 18 years old and over
- 3. 37 weeks gestation and over
- 4. Singleton pregnancy
- 5. Cephalic presentation
- 6. Reassuring cardiotocogram in last 1 hour
- 7. Cervical dilation 4-8 cm
- 8. Contraction ≥ 3 in 10 minutes
- 9. Pain score \geq 5 (0-10 NRS)

Participant type(s)

Patient

Healthy volunteers allowed

No

Age group

Adult

Lower age limit

18 years

Sex

Female

Total final enrolment

208

Key exclusion criteria

- 1. High-risk pregnancies (pre-eclampsia, active hemorrhage)
- 2. No prior pain relief (epidural, opiates, nitrous oxide or pain relief device)
- 3. Meconium-stained liquor
- 4. Oxytocin dose adjustment in the last hour
- 5. Bladder catheter in place
- 6. Delivery anticipated within the next 2 hours

Date of first enrolment

04/08/2022

Date of final enrolment

23/02/2023

Locations

Countries of recruitment

Malaysia

Study participating centre University Malaya Medical Centre (UMMC)

Lembah Pantai Kuala Lumpur Malaysia 59100

Sponsor information

Organisation

University Malaya Medical Centre

ROR

https://ror.org/00vkrxq08

Funder(s)

Funder type

University/education

Funder Name

Universiti Malaya

Alternative Name(s)

University of Malaya, University Malaya, Malayan University, King Edward VII College of Medicine, Raffles College, University of Malaya in Singapore, , , , UM

Funding Body Type

Government organisation

Funding Body Subtype

Universities (academic only)

Location

Malaysia

Results and Publications

Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date

IPD sharing plan summary

Data sharing statement to be made available at a later date

Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article		04/03/2024	10/09/2025	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes