# The iHOLDS Trial: High Or Low Dose Syntocinon for induction of labour

Submission date 13/05/2021	<b>Recruitment status</b> Suspended	[X] Prospectively registered		
		<pre>Protocol</pre>		
Registration date 20/05/2021	Overall study status Suspended	Statistical analysis plan		
		Results		
<b>Last Edited</b> 20/11/2023	<b>Condition category</b> Pregnancy and Childbirth	Individual participant data		
		<ul><li>Record updated in last year</li></ul>		

#### Plain English summary of protocol

Background and study aims

About 34% of (or one in three) first time mothers have their labour induced. Labour induction is the process we use to start labour artificially. This may be required for a variety of reasons, most commonly because a woman has gone past her due date, or because her waters have broken but labour contractions have not yet started.

Labour can be induced in different ways. Usually the cervix (neck of the womb) is softened using vaginal prostaglandins and some women will go into labour at this point but many still require their waters to be broken artificially. If contractions still do not start, a drug is given using a drip in the arm to start them off. The drug used is artificial oxytocin and this is given to around 60% of women (six in ten) who have labour induced.

Oxytocin is a natural hormone produced by the brain which plays a significant role in child birth as it helps to make the muscles in the womb contract during labour. There are man-made drugs almost identical to the natural hormone oxytocin which are widely used across UK Maternity Units and have been since the 1960s. Syntocinon is one of these drugs, which helps the contractions build up to the best strength and speed to encourage vaginal birth. All over the country the same standard dose and speed is used to increase the contractions to the best strength and frequency. Once this happens the dose is not increased any further. The dose and speed of oxytocin are adjusted up or down depending on the number of contractions whilst making sure the baby stays safe.

Many mothers who undergo induction of labour (including those who receive oxytocin) require an unplanned caesarean section to safely deliver their baby. While necessary, caesarean sections (also known as c-sections) generally result in longer hospital stays. C-sections can increase the chance of complications for both mother and baby, and are major surgery that most women would rather avoid. Women whose first baby is delivered by caesarean section are also more likely to require caesarean sections for future babies.

We want to find out if increasing the dose of oxytocin will reduce the chances of requiring a caesarean section. This study will compare the standard dose used routinely in hospitals, with a higher dose. The aim is to achieve regular contractions more quickly, and also allows a higher

maximum dose to be used if necessary. The higher dose is already used in some hospitals but we need to see how well it works in comparison with the standard dose.

#### Who can participate?

First-time mothers aged 16 years or above for whom oxytocin is required for induction of labour.

#### What does the study involve?

Women requiring oxytocin for induction of labour who meet all other study eligibility criteria and are willing to participate will be asked to sign a Consent Form before being randomised to receive either the standard or higher dose of oxytocin. There is an equal chance of being given either dose, and research staff will not know which dose women have been allocated to (this is called a double-blinded study design).

Women will be given the allocated drug dose through a drip in the arm, and mother and baby will be closely monitored. Information will be collected from the medical notes about the labour, birth and care and reported to the iHOLDS Trial Office. The mother will be asked to fill in a short multiple-choice questionnaire at home about two weeks after birth with some simple questions to find out how the birth experience was.

#### What are the possible benefits and risks of participating?

We cannot predict whether women or their babies will benefit directly from taking part in the trial, but the information we get from carrying out this study could potentially change the way oxytocin is administered to women in the future. Women will not be paid for their participation, however, all women who take part in the trial and return the Birth Satisfaction Questionnaire will be sent a £5 Amazon shopping voucher.

Where is the study run from?
Birmingham Clinical Trials Unit (BCTU) at the University of Birmingham (UK)

When is the study starting and how long is it expected to run for? From October 2019 to November 2023

Who is funding the study?
The National Institute for Health Research (UK)

Who is the main contact?

1. Mrs Dee Wherton
iHOLDS@trials.bham.ac.uk

2. Prof. Sara Kenyon
S.Kenyon@bham.ac.uk

## Contact information

## Type(s)

Scientific

#### Contact name

Prof Sara Kenyon

#### **ORCID ID**

https://orcid.org/0000-0002-0756-9617

#### Contact details

Institute of Applied Health Research University of Birmingham Birmingham United Kingdom B15 2TT +44 (0)121 414 9063 S.kenyon@bham.ac.uk

## Additional identifiers

## Clinical Trials Information System (CTIS)

2020-004387-26

#### Integrated Research Application System (IRAS)

278209

#### ClinicalTrials.gov (NCT)

Nil known

#### Protocol serial number

CPMS 47006, IRAS 278209

## Study information

#### Scientific Title

High or low dose Syntocinon for induction of labour in nulliparous women: a double blind, randomised controlled trial

#### Acronym

**iHOLDS** 

#### Study objectives

In nulliparous women who require oxytocin as part of induction of labour, a high dose regimen reduces the rate of caesarean section (CSR) by at least 20% compared with a standard dose regimen.

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Approved 01/03/2021, West Midlands – Edgbaston Research Ethics Committee (HRA Centre – Manchester, 3rd Floor, Barlow House, 4 Minshull Street, Manchester M1 3DZ; +44 (0)207 1048127; edgbaston.rec@hra.nhs.uk), ref: 21/WM/0034

## Study design

Multi-centre double-blind randomized controlled trial

## Primary study design

#### Interventional

#### Study type(s)

Treatment

#### Health condition(s) or problem(s) studied

Induction of labour

#### Interventions

A pilot stage will be conducted to assess continuation of recruitment against predefined criteria using a traffic light system; these consider recruitment, treatment adherence and outcome data. The Trial Steering Committee will meet to assess these criteria and report their recommendations to the Sponsor and Funder regarding continuation of the trial.

The clinical pathway for nulliparous women recruited to iHOLDS is no different to those undergoing induction of labour who require oxytocin as standard care, other than the dose regimen that is given and the completion of the birth satisfaction questionnaire.

A discussion between the woman and the clinical team will take place whenever the induction is booked, and information, including a Participant Information Leaflet (PIL), will be given/sent to women. For some women, induction is booked in advance and for others, the decision is made immediately prior to the process starting; the PIL may therefore be posted to their home address or issued as an outpatient or inpatient in hospital.

Discussion regarding the trial will take place during the period of time that cervical ripening is undertaken, which for most women (either as an inpatient or outpatient) is 24 h or more. During this time, women will not be in early/active labour. Issuing of trial-related information may be carried out by Clinical Midwives, Research Midwives or Obstetricians (who may also be Investigators for the trial). Eligibility will be confirmed by a GCP target trained Obstetrician. After ample time for consideration and the opportunity to ask further questions, eligible women who may require oxytocin for induction of labour will be consented to the trial by any of the above named GCP target trained individuals who have been delegated the duty of obtaining informed consent on the Site Signature and Delegation Log or local Training Log.

Following consent, if oxytocin is indicated the site will telephone the 24 h randomisation line at the University of Aberdeen to perform randomisation and receive treatment pack allocation. The relevant treatment pack will be retrieved from the drug fridge and administered as per the protocol and local policy. Women randomised to the standard dose will receive a solution containing 2 x 5 iu ampoules made up to 50 or 500 ml of intravenous fluid and those randomised to the high dose, a solution containing 2 x 10 iu made up to 50 or 500 ml of intravenous fluid. The participant and her baby will be closely monitored throughout on the Delivery Suite and the rate of the dose increase to establish effective uterine contractions will be titrated in response to the frequency and strength of contractions and the condition of the baby by the midwife caring for the woman, overseen by the Obstetrician. If an additional allocation is required (e.g. due to continued treatment being required or damaged IMP) the site will telephone the randomisation line and select the relevant option to receive a new pack number (the dose of which will match the initial dose).

We have chosen to use regimens that fall within the ranges described within the Cochrane review, which minimise the escalation doses that are outside of the manufacturers recommendations (demonstrated in the Trial Treatment section of the protocol) and that

facilitate blinding as they match the increments of the standard dose regimen. We will therefore compare a high dose regimen of oxytocin (4 mU/min increasing every 30 minutes to a maximum of 64 mU/min) with a standard dose regimen (2 mU/min increasing every 30 minutes to a maximum 32 mU/min). The high dose regimen has a higher starting dose, earlier attainment of conventional maximum doses (at 2 h rather than 4 h) with the aim of the higher dose regimen being to achieve regular contractions more rapidly, rather than simply giving a higher total dose of oxytocin.

Case Report Forms (CRFs) will capture data routinely collected regarding labour, birth and discharge. If the baby is admitted to the Neonatal Unit, details of the admission will also be reported, in addition to any reportable serious adverse events (SAEs) which may be experienced by the mother or baby. Where relevant, deviations from the trial protocol will be reported, in addition to details of withdrawal of consent.

Approximately 14 days after birth women will be sent a questionnaire via their preferred method (collected on the ICF) by email, post or text to collect maternal psychological outcomes which will explore satisfaction with care and the experience of labour and birth. The BCTU will send questionnaires to women following confirmation from the site via the CRFs that the participant and her baby are alive and well.

Interim analyses of safety and efficacy for review by the independent Data Monitoring Committee (DMC) will take place during the study. The committee will meet prior to study commencement to agree the manner and timing of such analyses but this is likely to include the analysis of the primary and major secondary outcomes and full assessment of safety (SAEs) at least at annual intervals. Criteria for stopping or modifying the study based on this information will be ratified by the DMC.

The trial is expected to last 3 years (including setup time): the pilot recruitment phase will last for 8 months followed by 13 months of recruitment to the main trial. For the participants, the end of trial is defined as 8 weeks after birth to allow for the collection of maternal satisfaction data. This will allow sufficient time for the completion of protocol procedures, data collection and data input. The end of trial is defined as 6 months after the date of last data capture and following resolution of all data queries relating to critical data items.

#### Patient and public involvement:

PPI has been a fundamental part of our trial design process. A service user is a co-applicant for the study and has informed many decisions regarding the trial design. We also have PPI input on the Trial Steering Committee (TSC). Both of whom will continue to be involved with the development and oversight of this trial, including engagement with maternity service users at participating sites

#### Intervention Type

Drug

#### Phase

Not Applicable

### Drug/device/biological/vaccine name(s)

oxytocin

#### Primary outcome(s)

Birth by caesarean section collected on the case report form (CRF) at the birth of baby

#### Key secondary outcome(s))

Maternal outcomes:

- 1. Epidural use during labour and birth collected on the CRF at the birth of baby
- 2. Duration of the second stage of labour (between full dilation to the birth of the baby) collected on the CRF at the birth of baby
- 3. Duration of the third stage of labour (between the birth of the baby to the expulsion of the placenta and membranes) collected on the CRF at the expulsion of the placenta and membranes
- 4. Time from randomisation to birth (mins) collected on the CRF at the birth of baby
- 5. Time from induction of labour (induction of labour is defined as the process by which labour is started prior to its spontaneous onset by progressive cervical effacement and dilatation and/or artificial stimulation of uterine contractions, leading to active labour and birth) to birth (mins) collected on the CRF at the birth of baby
- 6. Mode of birth (spontaneous vaginal birth, instrumental or caesarean section) collected on the CRF at the birth of baby
- 7. Degree of perineal trauma collected on the CRF at the birth of baby. Defined as follows:
- 7.1. First degree, injury to skin only
- 7.2. Second degree, injury to the perineal muscles but not the anal sphincter
- 7.3. Third degree, injury to the perineum involving the anal sphincter complex:
- 7.3.1. 3a, less than 50% of external anal sphincter thickness torn
- 7.3.2. 3b, more than 50% of external anal sphincter thickness torn
- 7.3.3. 3c, internal anal sphincter torn
- 4. Fourth degree, injury to the perineum involving the anal sphincter complex (external and internal anal sphincter) and anal epithelium.
- 8. Reason for, and grade of caesarean section (immediate threat to the life of the woman or fetus, maternal or fetal compromise which is not immediately life-threatening, no maternal or fetal compromise but needs early delivery, or delivery timed to suit woman or staff) collected on the CRF at the time of caesarean section
- 9. Confirmed urinary retention requiring catheterization collected on the CRF at discharge from hospital
- 10. Tachysystole (≥5 uterine contractions in 10 mins for a 20 min period) requiring reduction in oxytocin and/or tocolysis collected on the CRF at the birth of baby
- 11. Hyperstimulation (≥5 uterine contractions in 10 mins for a 20 min period resulting in non-reassuring or abnormal fetal heart rate) collected on the CRF at the birth of baby
- 12. Fetal blood sampling (FBS) during labour or significant ST analysis (STAN) event (for those Units that use ST waveform analysis for intrapartum fetal monitoring) collected on the CRF at the birth of baby
- 13. Abnormal cardiotocogram leading to immediate birth without fetal blood sample collected on the CRF at the birth of baby
- 14. Active management of third stage of labour collected on the CRF at the birth of baby
- 15. Length of time after birth in hospital (days) collected on the CRF at discharge from hospital
- 16. Admission to HDU and/or ITU collected on the CRF at discharge from hospital
- 17. Maternal death collected on the CRF at maternal death
- 18. Satisfaction with care and the experience of labour measured using the Birth Satisfaction Scale-Revised Index (Maternal Satisfaction Questionnaire) at 2 weeks post-birth

#### Process outcomes:

- 1. Time from randomisation to commencement of allocation (mins) collected on the CRF between recruitment and allocation
- 2. Total oxytocin dose [IU] collected on the CRF at the birth of baby
- 3. Time to maximum oxytocin rate (mins) collected on the CRF at the birth of baby
- 4. Maximum oxytocin dose reached collected on the CRF at the birth of baby

#### Neonatal outcomes:

- 1. Birthweight collected on the CRF at the birth of baby
- 2. Apgar score at 5 min collected on the CRF at the birth of baby
- 3. Venous and arterial cord blood gases and pH collected on the CRF at the birth of baby
- 4. Need for resuscitation collected on the CRF at the birth of baby
- 5. Breastfeeding on discharge from hospital collected on the CRF at discharge from hospital
- 6. Length of time after birth in hospital (days) collected on the CRF at discharge from hospital
- 7. Birth trauma (brachial plexus injury, fractured clavicle) collected on the CRF at the birth of baby
- 8. Need for neonatal review on ward (excluding routine baby check) collected on the CRF at the birth of baby
- 9. Use of any antibiotics collected on the CRF at discharge from hospital
- 10. Jaundice requiring phototherapy and/or transfusion collected on the CRF at discharge from hospital
- 11. Level of Neonatal Unit care received (level 1,2,3) including Intensive Care collected on the CRF at discharge from hospital
- 12. Duration of respiratory support (days) collected on the CRF at discharge from hospital
- 13. Days to full oral feeds collected on the CRF at discharge from hospital
- 14. Meconium aspiration syndrome collected on the CRF at discharge from hospital
- 15. Seizures collected on the CRF at discharge from hospital
- 16. Neonatal encephalopathy measured using SARNAT grade and collected on the CRF at discharge from hospital
- 17. Therapeutic hypothermia (cooling) collected on the CRF at discharge from hospital
- 18. Intrapartum stillbirth collected on the CRF at the birth of baby
- 19. Early neonatal death (within seven days of birth) collected on the CRF at time of neonatal death

#### Completion date

30/09/2025

## **Eligibility**

#### Key inclusion criteria

- 1. Nulliparous women at >24 +0/40 weeks gestation
- 2. Singleton cephalic pregnancy
- 3. Ruptured membranes undergoing induction of labour
- 4. Prescribed oxytocin is indicated as part of the induction process
- 5. Recieved prostin >6 h ago, or propess >30 min ago (if applicable)
- 6. Clinicians are willing to randomise
- 7. Give written informed consent to participate prior to randomisation
- 8. Aged ≥16 years
- 9. COVID-19 positive participants are eligible for study inclusion in accordance with local Trust /Health Board policy

#### Participant type(s)

**Patient** 

#### Healthy volunteers allowed

No

#### Age group

#### Adult

#### Sex

**Female** 

#### Key exclusion criteria

Nulliparous women who:

- 1. Are in the second stage of labour
- 2. Have any of the following conditions:
- 2.1. Existing cardiac disease
- 2.2. Bleeding disorders
- 2.3. Previous uterine surgery
- 2.4. Significant antepartum haemorrhage
- 3. Have a known contra-indication to oxytocin therapy
- 4. Are participating in other interventional trials of an Investigational Medicinal Product (IMP) or procedure for induction of labour

#### Date of first enrolment

01/09/2021

#### Date of final enrolment

31/05/2023

## Locations

#### Countries of recruitment

United Kingdom

England

Scotland

Wales

## Study participating centre

Birmingham Women's and Children's NHS Foundation Trust

Birmingham Children's Hospital Steelhouse Lane Birmingham United Kingdom B4 6NH

## Study participating centre Guy's and St Thomas' NHS Foundation Trust

St Thomas' Hospital Westminster Bridge Road London United Kingdom SE1 7EH

## Study participating centre South Tyneside And Sunderland NHS Foundation Trust

Sunderland Royal Hospital Kayll Road Sunderland United Kingdom SR4 7TP

### Study participating centre

Chelsea And Westminster Hospital NHS Foundation Trust

Chelsea & Westminster Hospital
369 Fulham Road
London
United Kingdom
SW10 9NH

## Study participating centre Shrewsbury And Telford Hospital NHS Trust

Mytton Oak Road Shrewsbury United Kingdom SY3 8XO

## Study participating centre NHS Greater Glasgow and Clyde

J B Russell House Gartnavel Royal Hospital 1055 Great Western Road Glasgow United Kingdom G12 0XH

## Study participating centre South Tees Hospitals NHS Foundation Trust

The James Cook University Hospital Marton Road

Middlesbrough United Kingdom TS4 3BW

## Study participating centre North Tees And Hartlepool NHS Foundation Trust

University Hospital Of Hartlepool Holdforth Road Hartlepool United Kingdom TS24 9AH

## Study participating centre County Durham And Darlington NHS Foundation Trust

Darlington Memorial Hospital Hollyhurst Road Darlington United Kingdom DL3 6HX

## Study participating centre Great Western Hospitals NHS Foundation Trust

Great Western Hospital Marlborough Road Swindon United Kingdom SN3 6BB

## Study participating centre Cardiff & Vale University LHB

Woodland House Maes-y-coed Road Cardiff United Kingdom CF14 4HH

# Study participating centre East Lancashire Hospitals NHS Trust Royal Blackburn Hospital

Royal Blackburn Hospital Haslingden Road Blackburn United Kingdom BB2 3HH

## Study participating centre Buckinghamshire Healthcare NHS Trust

Amersham Hospital Whielden Street Amersham United Kingdom HP7 0JD

## Study participating centre Aneurin Bevan University LHB

Headquarters - St Cadoc s Hospital Lodge Road Caerleon Newport United Kingdom NP18 3XQ

## Study participating centre Manchester University NHS Foundation Trust

Cobbett House Oxford Road Manchester United Kingdom M13 9WL

## Study participating centre NHS Lothian

Waverley Gate 2-4 Waterloo Place Edinburgh United Kingdom EH1 3EG

## Study participating centre Nottingham University Hospitals NHS Trust

Trust Headquarters Queens Medical Centre Derby Road Nottingham United Kingdom NG7 2UH

## Study participating centre Ashford And St Peter's Hospitals NHS Foundation Trust

St Peters Hospital Guildford Road Chertsey United Kingdom KT16 0PZ

## Study participating centre Royal Cornwall Hospitals NHS Trust

Royal Cornwall Hospital Treliske Truro United Kingdom TR1 3LJ

## Study participating centre Leeds Teaching Hospitals NHS Trust

St. James's University Hospital Beckett Street Leeds United Kingdom LS9 7TF

## Sponsor information

## Organisation

Birmingham Women's and Children's NHS Foundation Trust

#### **ROR**

https://ror.org/056ajev02

## Funder(s)

Funder type

#### **Funder Name**

National Institute for Health Research

#### Alternative Name(s)

National Institute for Health Research, NIHR Research, NIHRresearch, NIHR - National Institute for Health Research, NIHR (The National Institute for Health and Care Research), NIHR

#### Funding Body Type

Government organisation

#### **Funding Body Subtype**

National government

#### Location

United Kingdom

## **Results and Publications**

#### Individual participant data (IPD) sharing plan

The data sharing plans for the current study are unknown and will be made available at a later date

## IPD sharing plan summary

Data sharing statement to be made available at a later date

#### **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
HRA research summary			28/06/2023		No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes
Study website	Study website	11/11/2025	11/11/2025	No	Yes