# The effect of reiki on pain in women undergoing non-emergency caesarean sections

Submission date	Recruitment status  No longer recruiting	<ul><li>Prospectively registered</li></ul>		
14/04/2009		☐ Protocol		
Registration date 27/05/2009	Overall study status Completed	Statistical analysis plan		
		[X] Results		
Last Edited	Condition category	[] Individual participant data		
30/12/2020	Pregnancy and Childbirth			

## Plain English summary of protocol

Not provided at time of registration

# Contact information

## Type(s)

Scientific

#### Contact name

Dr Gideon Koren

#### Contact details

Professor of Pediatrics, Pharmacology, Pharmacy, Medicine and Medical Genetics Clinician-Scientist The Hospital for Sick Children Division of Clinical Pharmacology, Rm 8232 555 University Ave Toronto Canada M5G 1X8 +1 416 813 1500 ext. 5781 gkoren@sickkids.ca

# Additional identifiers

Protocol serial number

REB#-08-030

# Study information

Scientific Title

The effect of reiki on pain in women who will undergo non-emergency caesarean sections and the effect of genetic variation on the use of codeine: a randomised controlled trial

#### Acronym

**REPA** 

#### **Study objectives**

Can reiki reduce pain after caesarean section by more than 25%?

#### Ethics approval required

Old ethics approval format

#### Ethics approval(s)

St. Michael's Hospital Ethics Review Board approved on the 12th May 2008 (ref: 08-030)

#### Study design

Randomised controlled trial

#### Primary study design

Interventional

# Study type(s)

Treatment

# Health condition(s) or problem(s) studied

Pain levels after caesarean section

#### **Interventions**

- 1. Control group: usual care
- 2. Reiki Group: usual care plus 20 minutes of Remote Reiki

Duration: three days while in the hospital.

## Intervention Type

Other

#### Phase

Not Applicable

#### Primary outcome(s)

Visual Analogue Scale for pain in movement, at day 1, day 2 and day 3 (amended as of 10/08/09; initial outcome timepoints were 'after 72 hours in hospital').

#### Key secondary outcome(s))

- 1. Visual Analogue Scale for pain (in rest)
- 2. Pain medication (codeine consumption/kg)
- 3. Return time to normal activities

All measured at day 1, day 2 and day 3 (amended as of 10/08/09; initial outcome timepoints were 'after 72 hours in hospital').

# Completion date

30/04/2009

# Eligibility

## Key inclusion criteria

- 1. Informed consent
- 2. Planned caesarean section
- 3. Pregnant women aged 18 48 years

# Participant type(s)

**Patient** 

# Healthy volunteers allowed

No

# Age group

Adult

#### Lower age limit

18 years

#### Sex

Female

#### Total final enrolment

80

#### Key exclusion criteria

Does not meet with inclusion criteria

#### Date of first enrolment

30/06/2008

#### Date of final enrolment

30/04/2009

# Locations

## Countries of recruitment

Canada

#### Study participating centre

Professor of Pediatrics, Pharmacology, Pharmacy, Medicine and Medical Genetics

Toronto

Canada

M5G 1X8

# Sponsor information

# Organisation

The Hospital for Sick Children (Canada)

#### **ROR**

https://ror.org/057q4rt57

# Funder(s)

# Funder type

Other

#### Funder Name

Investigator initiated and funded (Canada)

# **Results and Publications**

Individual participant data (IPD) sharing plan

# IPD sharing plan summary

Not provided at time of registration

# **Study outputs**

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
Results article	results	26/02/2011	30/12/2020	Yes	No
Participant information sheet	Participant information sheet	11/11/2025	11/11/2025	No	Yes