

# Double Blind Randomised Placebo Controlled Trial of Oral Co-trimoxazole in Pulmonary Fibrosis

<b>Submission date</b> 25/08/2005	<b>Recruitment status</b> No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
<b>Registration date</b> 09/09/2005	<b>Overall study status</b> Completed	<input type="checkbox"/> Statistical analysis plan <input checked="" type="checkbox"/> Results
<b>Last Edited</b> 05/07/2018	<b>Condition category</b> Circulatory System	<input type="checkbox"/> Individual participant data

**Plain English summary of protocol**  
Not provided at time of registration

## Contact information

**Type(s)**  
Scientific

**Contact name**  
Dr Veronica Varney

**Contact details**  
St Helier Hospital  
Wrythe Lane  
Carshalton  
United Kingdom  
SM5 1AA  
+44 (0)20 8296 2401  
vvarney@epsom-sthelier.nhs.uk

## Additional identifiers

**EudraCT/CTIS number**

**IRAS number**

**ClinicalTrials.gov number**

**Secondary identifying numbers**

UK study number for local regional ethical committee (LREC) 64/99

# Study information

## Scientific Title

Double Blind Randomised Placebo Controlled Trial of Oral Co-trimoxazole in Pulmonary Fibrosis

## Acronym

Septrin and CFA

## Study objectives

Oral Co-trimoxazole improves exercise capacity in patients with pulmonary fibrosis

## Ethics approval required

Old ethics approval format

## Ethics approval(s)

Not provided at time of registration

## Study design

Randomised controlled trial

## Primary study design

Interventional

## Secondary study design

Randomised controlled trial

## Study setting(s)

Not specified

## Study type(s)

Treatment

## Participant information sheet

## Health condition(s) or problem(s) studied

Idiopathic pulmonary fibrosis (UIP/NSIP/Mixed)

## Interventions

Investigations prior to selection and randomisation:

All selected patients underwent baseline investigations before entry. This included:

1. Arterial gases at rest (on air)
2. Pulmonary function tests (laboratory measurement of TLC, DLCO)
3. St Georges Hospital respiratory questionnaire (SGHRQ), and MRC 5 Point Dyspnoea Score
4. Two shuttle-walking tests (2 weeks apart) demonstrating oxygen desaturation below 90% (either during or upon cessation of exercise)
5. An echocardiogram and ECG
6. Routine bloods tests including serum samples for cytokine measurements
7. Sputum samples for pneumocystis silver stain at request of Ethics Committee

**Randomisation:**

Patients were randomly allocated to active or placebo treatments by our clinical trials pharmacist, using computer generated random numbers. This assigned patients to study treatment groups, with a patient number. The study was conducted double blind. To preserve the double blind status of the trial, medication was issued by the pharmacist; using identical placebo tablets and labelling to ensure all treatment packs were identical.

**Drug Treatment:**

Co-trimoxazole or identical placebo (480 mg) tablets were supplied with dosage according to body-weight. Patients up to 70 kg received 960 mg bd, those above 70 kg, took 1440 mg (3 x 480 mg) bd. Folic acid 5 mg was given 3 times a week (minimum dose) to protect the bone marrow. Ranitidine 150 mg bd was supplied but optional for any indigestion. Patients on oral daily prednisolone had no adjustments to their prednisolone dose throughout the double blind study, except that permitted during a viral illness.

**Assessment during Study:**

On entry to the study all patients made visits for the following assessments every 2 weeks:

1. Body-weight
2. Resting respiratory rate and chest auscultation
3. FVC
4. Oxygen saturations at rest and during shuttle walking test and recovery recorded by Pulsox 3i wrist watch Minolta range
5. MRC-5 Point Dyspnoea Score
6. Compliance with treatment/side effects/study withdrawal and deaths were assessed and recorded
7. Blood tests (FBC, U & E, LFTs, ESR and CRP)

At 3 months all patients had arterial gases, full pulmonary function tests, serum cytokines and SGHRQ, repeated. Six weeks of pulmonary rehabilitation (rehab) was then commenced. Two weeks post rehabilitation final assessments were made before decoding. These assessments were identical to the regular 2 week assessments during the study.

**Intervention Type**

Drug

**Phase**

Not Specified

**Drug/device/biological/vaccine name(s)**

Co-trimoxazole

**Primary outcome measure**

Exercise capacity (shuttle walking test) with area under the curve oxygen desaturation

**Secondary outcome measures**

1. Respiratory function tests (FVC, TLC, DLCO)
2. Arterial oxygen measurements
3. Quality of life data
4. Benefit of pulmonary rehabilitation

Twenty patients were selected.

**Overall study start date**

21/02/2000

**Completion date**

01/02/2004

## Eligibility

**Key inclusion criteria**

1. Male or female below 85 years old
2. Demonstrated breathlessness with oxygen desaturation below 90% on two baseline shuttle walking tests (SWT)
3. Physical examination, HRCT scan and pulmonary function test results compatible with IPF (with or without histological diagnosis)
4. New diagnosis of IPF without treatment or previous diagnosis on regular daily prednisolone
5. Symptoms of exertional dyspnoea affecting life quality
6. Normal glucose 6-phosphate dehydrogenase levels to avoid drug induced haemolysis. Normal vitamin B12 levels

**Participant type(s)**

Patient

**Age group**

Adult

**Sex**

Both

**Target number of participants**

20

**Key exclusion criteria**

1. Recognised secondary causes of pulmonary fibrosis
2. Co-trimoxazole allergy or severe upper GI symptoms
3. Abnormal liver function tests or concurrent drug treatments that disturb liver function including azathioprine
4. Inability to perform the shuttle test due to musculoskeletal or cardiac causes

**Date of first enrolment**

21/02/2000

**Date of final enrolment**

01/02/2004

## Locations

**Countries of recruitment**

England

United Kingdom

**Study participating centre**  
**St Helier Hospital**  
Carshalton  
United Kingdom  
SM5 1AA

## **Sponsor information**

**Organisation**  
The Peel Trust Fund (UK)

**Sponsor details**  
Sceptre Court  
40 Tower Hill  
Gloucester  
United Kingdom  
GL1 3NN

**Sponsor type**  
Charity

**ROR**  
<https://ror.org/05ag50972>

## **Funder(s)**

**Funder type**  
Charity

**Funder Name**  
Part funded by The Peel Trust Fund for funding of placebo and Co-trimoxazole drugs only

## **Results and Publications**

**Publication and dissemination plan**  
Not provided at time of registration

**Intention to publish date**

**Individual participant data (IPD) sharing plan**

## IPD sharing plan summary

Not provided at time of registration

## Study outputs

Output type	Details	Date created	Date added	Peer reviewed?	Patient-facing?
<a href="#">Results article</a>	results	01/05/2008		Yes	No