

Influence of intravenous versus oral administration, arterial versus venous sampling and gender on pharmacokinetic-pharmacodynamic modelling of morphine and morphine-6-glucuronide-induced pain relief in healthy volunteers

Submission date 20/12/2005	Recruitment status No longer recruiting	<input type="checkbox"/> Prospectively registered <input type="checkbox"/> Protocol
Registration date 20/12/2005	Overall study status Completed	<input type="checkbox"/> Statistical analysis plan <input type="checkbox"/> Results
Last Edited 13/05/2009	Condition category Other	<input type="checkbox"/> Individual participant data <input type="checkbox"/> Record updated in last year

Plain English summary of protocol
Not provided at time of registration

Contact information

Type(s)
Scientific

Contact name
Prof Albert Dahan

Contact details
Leiden University Medical Centre (LUMC)
Department of Anesthesiology
P.O. Box 9600
Leiden
Netherlands
2300 RC
adahan@lumc.nl

Additional identifiers

EudraCT/CTIS number

IRAS number

ClinicalTrials.gov number

Secondary identifying numbers

NTR229

Study information

Scientific Title

Study objectives

This study is designed to get a full pharmacokinetic-pharmacodynamic (PK/PD) characteristic of the opioid analgesic morphine and its active metabolite M6G after oral and intravenous (iv) infusion and to test whether sex differences exist in the analgesic behaviour of both opioids.

Ethics approval required

Old ethics approval format

Ethics approval(s)

Received from the local medical ethics committee

Study design

Randomised double blind active controlled parallel group trial

Primary study design

Interventional

Secondary study design

Randomised controlled trial

Study setting(s)

Hospital

Study type(s)

Treatment

Participant information sheet

Health condition(s) or problem(s) studied

Post-operative pain

Interventions

Drug administration (blinded): morphine or M6G.

Intervention Type

Drug

Phase

Not Applicable

Drug/device/biological/vaccine name(s)

Morphine

Primary outcome measure

Pain relief related parameters (Visual Analogue Scale [VAS] to heat pain) in males versus females.

Secondary outcome measures

PK parameters

Overall study start date

01/11/2005

Completion date

01/11/2006

Eligibility

Key inclusion criteria

Healthy volunteers, aged 18+ years

Participant type(s)

Healthy volunteer

Age group

Adult

Lower age limit

18 Years

Sex

Both

Target number of participants

60

Key exclusion criteria

1. Body mass index (BMI) greater than 30 kg/m²
2. Pregnancy or lactation
3. Presence of medical disease
4. Presence of psychiatric disease
5. Allergy to study medication
6. History of drugs or alcohol abuse

Date of first enrolment

01/11/2005

Date of final enrolment

01/11/2006

Locations

Countries of recruitment

Netherlands

Study participating centre

Leiden University Medical Centre (LUMC)

Leiden

Netherlands

2300 RC

Sponsor information

Organisation

Leiden University Medical Centre (LUMC) (Netherlands)

Sponsor details

Albinusdreef 2

P.O. Box 9600

Leiden

Netherlands

2300 RC

Sponsor type

University/education

Website

<http://www.lumc.nl/>

ROR

<https://ror.org/027bh9e22>

Funder(s)

Funder type

Industry

Funder Name

CeNes Ltd (UK)

Results and Publications

Publication and dissemination plan

Not provided at time of registration

Intention to publish date

Individual participant data (IPD) sharing plan

IPD sharing plan summary

Not provided at time of registration